

## **Review Article**

### **AN EXTENSIVE REVIEW: INDUSTRIALLY IMPORTANT ENZYMES, THEIR CLASSIFICATION AND DEPICTING THE CONSORTIUM OF INDUSTRIAL APPLICATIONS**

#### **ABSTRACT**

Microorganism has been used since the start of human society. Enzyme processing ~~processes~~ processes are quickly gaining the attention because of their short time of processing, cost effective, non-toxic, low energy input and environment friendly characters as well. Moreover, through protein engineering and recombinant DNA technology, a microorganism can be easily manipulated and cultured in large scale to meet increased demand in different sectors. Therapeutic enzymes have a huge variety of selective uses such as anticoagulants or thrombolytics, oncolytics and also as replacement for various metabolic deficiencies. Proteolytic enzymes are excellent anti-inflammatory agents. There are various factors that decrease the potential of microbial source enzymes, once we enter to medical sciences due to high molecular size of catalyst which stop their distribution within somatic cells. In industrial processes the quantity of enzymes should be high, while in therapeutic case the purity and ~~specificity~~ specificity should be excellent, if the quantity level is less no matter. The kinetics of such enzymes is high and low so that it is maximally efficient even at low concentration of substrate and enzymes. The source of such enzymes should be designated with high care to minimize or prevent the chances of undesirable growth by mismatched material and also to enable ready purification.

Comment [1]: specificity

#### **INTRODUCTION**

Microorganism has been used since the start of human society. The first reported microorganism is the uses of yeast to make alcoholic products from barley by the Sumerians and Babylonians as early as 6000 BC ([reference?](#)). Microbes enzymes catch their attention globally for their great

application in distinguish sectors of industries, such as in chemical, energy, food, medicine and agriculture. Enzyme processing ~~processes~~~~process~~ are quickly gaining the attention because of their short time of processing, cost effective, non\_toxic, low energy input and environment friendly characters as well. Moreover through protein engineering and recombinant DNA technology a microorganism can be easily manipulated and cultured in large scale to meet increased demand in different sectors. Related driving factors that<sup>2</sup>s enhance the use of microbes enzyme products in industrial uses are increasing demand of user, natural resources reduction, necessity of cost reduction and eco-friendly. World market for industrial microbial enzymes was projected nearly about \$4.2-4.5 billion in 2014 and predictable to make at a compound AGR of nearly 8% over the time from 2016 to 2020 to influence \$6.4 billion(Kour et al., 2019). Microbial enzymes are biological compounds, protein in nature except catalytic ribonucleic acid molecules and ~~act as a catalystaet-catalyst~~ to speed up the chemical reaction and to support almost all of the chemical reaction needed for the life (Gurung, Ray, Bose, & Rai, 2013). Enzymes are highly specific in nature; speed\_up the chemical reaction by decreasing the activation energy of the chemical reaction with any permanent change in the reaction, and therefore major molecules to support life (Anastas, Bartlett, Kirchhoff, & Williamson, 2000). For enzyme activity they need optimal temperature and pressure for catalyzing reaction, and are uses in substitute toxic chemical pollutant owing to their non\_toxic and biodegradable nature (Anastas et al., 2000).

In addition to advantages of enzymes over conventional methods, there are some drawbacks of using enzymes in healthcare and other industries. For many mammalian enzymes, 37\_°C and 7.4 are the optimal temperature and pH, respectively, and their activity is highly sensitive to any change in these parameters. Higher temperature (>40\_°C), and a large deviation from the

physiological pH (7.4) lead to their denaturation, which limits the use of these macromolecules in non-physiological conditions. Additionally, they are susceptible to substrate or product inhibition and their products may cause allergic reactions. The high cost of isolation and purification of enzymes and their difficult recovery for subsequent reuse may discourage their use (Singh, Kumar, Mittal, & Mehta, 2016).

## ENZYMES

Enzymes are large molecules consist of AA attached through amide NH<sub>2</sub> bonds, range between kilo-Dalton to mega-Dalton in molecular mass. The specificity of enzymes molecules to catalyze reaction between one kinds of chemical compound ~~over the offers~~ the center of its name and classification. In biochemistry area a great advancement achieved after 1940, a huge number of ~~enzymes were isolated~~ ~~enzyme isolate~~ and identified, for this purpose it was mandatory to legalize enzyme nomenclature. IUBMB in discussion with International Union of Pure and Applied Chemistry develop an enzyme Commission ~~to be in charge~~ ~~to in-charge~~ of managing systemic classification and naming to enzymes ([reference?](#)). Enzyme was classified into six different groups on the basis of the reaction they catalyzed. For industrial enzymes, microbes are the best choice because of their easily availability, and fast growth rate of microbes. Through recombinant DNA technology genetic changes can easily be done in microbes for the production of enzymes and scientific development (Liu, [year?](#)). Microbial enzymes production is mandatory event in industrial zone, due to the loftier and great performance of enzymes from various microorganisms, because they performed well in wide range of chemical and physical condition (Anbu [et al.](#), ~~Gopinath, Chaulagain, Tang, & Citartan~~, 2015). Microbial enzymes help in the cure of many disorder linked with the absence of human enzymes caused due to genetic changes. Some enzymes are greater than the substrate they act on, and only a very small part of enzymes

Comment [2]: Not clearly understood

is directly involved in catalysis (Almeida ~~et al., Serra, Silveira, & Moura~~, 2010). Enzymes also have a site that bind to the cofactors, which are required for catalysis of the reaction (Bartlett ~~et al., Porter, Borkakoti, & Thornton~~, 2002). Like protein enzymes have linear chain of AA, long that fold to make a 3D products. Each AA sequence ~~makesmake~~ a selective structure, which has unique properties. ~~MostMostly~~ enzymes are denatured that is outspread and deactivated by chemical or heating, which disrupt 3D structure of the protein. Denaturation may be irreversible or ~~reversible dependentreversibledependent~~ on the enzymes. Enzymes are classified into six groups on the basis of reaction they catalyze (Kotera ~~et al., Okuno, Hattori, Goto, & Kanehisa~~, 2004).

Formatted: Font: Italic

Comment [3]: Repetition from above paragraph

### **Amylase enzyme**

Amylase is an enzyme that break starch molecule into sugars. Amylase enzyme is commonly found in saliva that breakdown the food and help in digestion. Some foods contain high amount of starch and contain slight sugar, such as potato and rice, taste somewhat as they are masticated because amylase cracks a few of their starch into sugar ~~in the humanin-human~~ mouth. Pancreas also produce amylase which hydrolyze dietary starch into disaccharide carbohydrates and trisacchrides which are rehabilitated by other enzymes. Bacteria and some plants also produce amylase. In 1833 amylase was the first enzyme isolated and identified by Anselme Payen. Entirely amylases act on  $\alpha$ -1,4-glycosidic and are glycoside hydrolases. Amylases has nearly 25 % enzyme market ~~and is widelyand widely~~ used in industry (Edner et al., 2007). ~~In the presentIn present~~ area chemical hydrolysis of starch in starch processing industry is almost replaced by amylases. Amylases isolated from microbes are more stable than plants and ~~animalsanimal~~ and for ~~that reasonthat's reason~~ they ~~have a broadhave broad~~ spectrum of industrial uses. Microbes are easy to change their nature and desired enzymes, so the major benefit of microbes for the

production of enzymes is the mass production and economical. In biotechnology sector starch hydrolyzing amylolytic enzymes are of high advantages ranging from paper, textile, fermentation and food industries. Amylases can be extracted from several sources such animals and plants but enzymes obtained from microbes generally satisfied industrial burdens and had made greater contribution to the beravges and food industry(Souza, 2010).

Comment [4]: Repetition

## Types

Amylase has three sub-classes — $\alpha$ -  $\beta$ -  $\gamma$ -amylase.

### $\alpha$ -Amylase

Alpha amylase enzymes help in the hydrolysis or breakdown of the internal  $\alpha$ -1,4-glycosidic linkages in low molecular weight ~~compound~~ such as maltotriose, maltose and glucose (Sundarram & Murthy, 2014).  $\alpha$ -Amylase are important form of amylase found in human, cow, sheep, and in other mammals (Butterworth ~~et al.~~, Warren, & Ellis, 2011). It is also present in those seeds of plants that comprise of starch as food reserve and is released by numerous ~~fungifungus~~. Amylyse are also present in high amount in saliva and pancreatic juice, each of which has itself isoform of human amylase. In human amylase are connected to 1p21 chromosome.

### Uses of $\alpha$ -Amylase

$\alpha$ -Amylase ~~is used~~ in production of ethyl alcohol by break down the starch in grains into sugars. The intital phase in the making of high fructose corn syrup is the treatment of cornstarch

with  $\alpha$ -amylase, ~~making a smaller~~making smaller chain of sugars known as oligosaccharides.

Some microbial source alpha amylases help in removing starch detergent (Butterworth et al., 2011).

Comment [5]: Rephrase

### **$\beta$ -Amylase**

$\beta$ -amylase is another form of amylases synthesized by plants, fungi and bacteria. The alternative name for  $\beta$ -amylase is 1, 4- $\alpha$ -D-glucan maltohydrolase; saccharogen amylase; glycogenase).  ~~$\beta$ -amylase, working~~~~from a non-reducing~~from non-reducing end, catalyzes the breakdown of the 2<sup>nd</sup>  $\alpha$ -1,4 glycosidic bond, ~~breaking two~~breaking of two glucose ~~units~~unit at the same time.  $\beta$ -amylase breaks the starch into maltose during the ripening of fruits, resulting in the sweet flavor of ripe fruits.  $\alpha$ -amylase and  $\beta$ -amylase are found in seeds,  $\beta$ -amylase are inactive before the germination while proteases and  $\alpha$ -amylase seen once germination has started. For the production of malt cereal grain amylase is the key. Different ~~microorganisms~~microorganism also produce amylases that help in the degradation of extracellular starches. Animal ~~cell~~cell and ~~tissue~~tissue do not have  $\beta$ -amylase, although ~~the~~ microorganisms present in ~~the~~ gut of animal produce  $\beta$ -amylase. The best pH for  $\beta$ -amylase is 4-5.

Comment [6]: Whole paragraph written without any reference

### **Use of $\beta$ -amylase**

Both alpha and beta amylases are significant in ~~making liquor~~making liquor and beer prepared from sugars consequent from starch. In microbial fermentation process, ~~mushrooms~~mushroom ingest sugars and ~~produce ethyl~~defecateethyl alcohol. In liquors and beer, ~~the sugars that~~ ~~exists~~sugars exist at the start of fermentation have been formed by "squashing" grains or other starch ~~cradles~~.

Comment [7]: Rephrase

Comment [8]: Need to rephrase this section

### **$\gamma$ -Amylase**

Gamma-amylase is an amylase which break  $\alpha(1-4)$  glycosidic and  $\alpha(1-6)$  glycosidic linkages at the non-reducing end of amylopectin and amylose, resulting in glucose. The  $\gamma$ -amylase has high acidic pH because they are highly active at pH 3. The alternative name for  $\gamma$ -amylase is 1, 4- $\alpha$ -glycosidase.

### Uses $\gamma$ -Amylase

$\gamma$ -Amylase are used in chemical, pharmacological, drugs distribution and food industry, as well as environmental and agriculture engineering. Hydroxypropyl beta cyclodextrin is the main compound present in Gamble's and [Procter Freshening](#) ~~Procter freshening~~ [products](#) ~~product~~.

Comment [9]: Reference?

### Bacterial spp amylase

Amylase can be extracted from various species of microbes, but for commercial use,  $\alpha$ -amylase derived from *Bacillus amyloliquefaciens*, *Bacillus licheniformis* and *Bacillus stearothermophilus*, has wide range of application in various industries such as in paper, textile, fermentation and food industries (Souza, 2010). Industrial enzymes has major and desirable characters known as thermostability. Temperature stable enzymes have found huge number of commercial significance due to their stability in high temperature. The production of valuable industrial products such as [glucose](#) ~~glucose~~, dextrose syrup, maltose, crystalline dextrose and maltodextrins are improving by using [thermo stable](#) ~~thermos table~~ amylolytic enzymes (Guzmán- Maldonado [et al.](#), ~~Paredes López,~~ & Biliaderis, 1995). *Bacillus amyloliquefaciens*, *Bacillus subtilis*, *Bacillus licheniformis* and *Bacillus stearothermophilus* are widely used for industrial level production of enzymes for huge application. In present temperature resistance [enzymes](#) ~~enzyme~~ of *Bacillus licheniformis* or *Bacillus stearothermophilus* are used in starch degrading/processing industry (Prakash & Jaiswal, 2010). In some [industries](#) ~~industry~~ high

~~levels~~ level of harsh processes occur and contain huge ~~amounts of amount~~ salt solution which stop/ inhibit the many enzymatic ~~reactions~~ reaction. For this purpose halophilic microbes have excellent activity at high salt concentration and enzymes ~~produced could produced them could~~ be used in such processes (Ventosa & Nieto, 1995). ~~Besides~~ Beside such a good ~~character~~, ~~halobacterial~~ character halobacterial enzyme are temperature stable and can survive for longer period of time at high temperature. Halophilic amylases have been resulting from bacteria such as and *Bacillus dipsosauri*, *Chromohalobacter* sp., *Haloarcula hispanica* (Caton et al., 2004) and *Halomonas meridian* (Edbeib et al., Wahab, & Huyop, 2016).

### Fungal Fungus Amylase

Many ~~researchers~~ researcher work on ~~fungi~~ fungus to produce amylases and ~~to~~ choose the best strains to produce amylase commercially. Previously, mesophilic ~~fungi were reported~~ fungi ~~reported~~ that produce alpha-amylase. Terrestrial isolates of fungi such as *Penicillium* and *Aspergillus* ~~produce~~ *Aspergillus* produce various kinds of enzymes (Kango et al., Jana, & Choukade, 2019). *Aspergillus* spp mainly make a different varieties of extracellular enzymes, and amylases are the ones with most important industrial value. *Aspergillus niger*, and *Aspergillus oryzae* are filamentous ~~fungi that produce~~ fungi produce huge amount of enzymes that can be greatly use in industry. *A. oryzae* is well-thought-out to be the satisfactory host for the manufacture of heterologous protein as it has capability to conceal a huge amount of high valuable proteins and industrial products (Machida et al., 2005). *A. oryzae* has been greatly used in commercial enzymes containing  $\alpha$ -amylase and food such as soy sauce and organic acid such as acetic and citric acid (Hajar-Azhari et al., Wan, Ab Kadir, Abd Rahim, & Saari, 2018). *A. niger* has a significant hydrolytic capacities in the  $\alpha$ -amylase production, because of low pH/ acid tolerant, and they also avoid bacterial growth/contamination (Djekrif-Dakhmouche et al., e,

Gheribi Aoulmi, Meraihi, & Bennamoun, 2006). Fungal enzymes have generally safe status and that's why they are preferred over other microbial enzymes. The high temperature resistance fungus *Thermomyces lanuginosus* is a ~~great~~ ~~great~~ producer of amylase (Tiwari et al., 2015).

## LIPASE

Lipases are enzymes that breakdown the fats. Lipases are the sub-class of the esterase's (Arreaza et al., 1997). These enzymes ~~play a~~ ~~plays~~ significant role in processing of dietary lipids, digestion and transport. Certain viruses also contain genes that ~~encode~~ ~~encodes~~ lipases. Especially ~~in the small~~ ~~small~~ ~~intestine,~~ ~~most~~ ~~intestine~~ ~~most~~ lipases enzymes act ~~in a precise~~ ~~precise~~ position ~~on the glycerol~~ ~~glycerol~~ backbone of lipid substrate. (Rahman & Basri, 2006). Several other kinds ~~of lipase activities~~ ~~of activities of lipase~~ present in nature, such as sphingomyelinases and phospholipase, however, these are frequently treated distinctly from orthodox lipases. Some pathogenic microbes also produce lipases during infection, especially *Candida albicans* ~~which contain a huge~~ ~~contain~~ ~~albicans~~ ~~contain~~ ~~huge~~ number of lipases. ~~In the biotechnology~~ ~~In biotechnology~~ sector lipases ~~play a~~ ~~plays~~ significant role ~~and are valuable~~ ~~and a valuable~~ enzyme in these sectors (Saini ~~et al.,~~ ~~Saini,~~ & Dahiya, 2017). Microbial enzymes ~~have a wide~~ ~~have wide~~ range of ~~applications~~ ~~application~~ at industrial level, because of their differentiated enzymatic properties and substrate specificity (Goodman, 2010). Bacterial lipases enzymes are more stable than plant and animal lipases (Burhan et al., 2003).

## Types

There ~~is~~ ~~are~~ no specific kinds of lipases, but mainly ~~they are classified~~ ~~they classified~~ according to ~~their~~ ~~its~~ use, namely: hepatic lipase, human digestive lipase and pancreatic lipase.

## Uses of Lipase

Formatted: Font: Italic

Lipases ~~carry~~<sup>carried</sup> various biological processes ranging ~~from the daily~~<sup>from daily</sup> metabolism process of dietary triglycerides to inflammation and cell signaling (Pascoal ~~et al.~~<sup>et al.</sup>, ~~Estevinho, Martins, & Choupina,~~ 2018). Some lipases have specific selective function, while ~~others~~<sup>other</sup> ~~play~~<sup>plays</sup> significant role in extracellular ~~compartment~~<sup>compartment</sup>. In lysosome lipase enzyme is present in lysosomes, while other lipases ~~such as pancreatic~~<sup>such as pancreatic</sup> lipases, they veiled into extracellular ~~compartment~~<sup>compartment</sup> where they ~~do~~<sup>done</sup> the breakdown of dietary lipids into simpler forms that can be easily ~~transported~~<sup>transport</sup> throughout the body and absorbed. Bacteria and fungus may also conceal lipases to ease nutrient preoccupation from external medium. Certain bees and ~~wasps~~<sup>wasp</sup> ~~contain~~<sup>contains</sup> phospholipases that increase biological processes of inflammation and wound delivered by sting. Dandruff in ~~humans~~<sup>human</sup> is caused by a fungus known as *Malassezia globosa*, this fungus usually uses lipase to do the breakdown of sebum into oleic acid and enhance skin cell production causing dandruff (~~reference?~~). Lipases ~~play~~<sup>plays</sup> significant role in cheese and yogurt fermentation. In modern area lipases ~~are~~<sup>are</sup> used ~~as a versatile~~<sup>as versatile</sup> and cheap source to degrade lipid. ~~In~~<sup>In</sup> biotechnology sector, recombinant lipases are used in laundry, baking and even as ~~catalyst~~<sup>catalyst</sup>, also used as alternative approaches to change vegetable oil into fuel. Lipases are safer and ~~environmentally~~<sup>environmental</sup> friendly and also ~~replace~~<sup>replaced</sup> traditional ~~catalyst~~<sup>catalyst</sup> in biodiesel processing (Gurung et al., 2013).

### **Bacterial Lipase**

The lipase producing bacterial genera comprises ~~of~~<sup>of</sup>; *Burkholderia*, *Pseudomonas* and *Bacillus*. The bulk production of bacterial lipase are ~~easy~~<sup>easily</sup> and they are commercially very ~~significant~~<sup>significant</sup> ~~because~~<sup>significant of because</sup> of extracellular production (Gupta ~~et al.~~<sup>et al.</sup>, ~~Gupta, & Rathi,~~ 2004). Only a few bacterial species are used commercially for the production of lipases. The bacterial

genera used ~~commercially~~~~commercially~~ include, *Pseudomonas*, *Achromobacter*, *Corynebacterium*, *Alcaligenes*, *Enterococcus*, *Arthrobacter*, *Chromobacterium*, *Burkholderia* and *Bacillus*, and in various biotechnology applications lipases from *Pseudomonas* are greatly used. Different ~~products launched~~~~products are launched in the market~~~~in market~~ are bacterial lipases based ~~in a few~~~~in few~~ years, such as Lipomax and lumafast extracted from *Pseudomonas*, and their major application in detergent (Verma, 2019).

### **Fungal Lipase**

Fungi that produce lipases are present in different habitats, such as dairy byproducts, soil contaminated with waste of vegetables, deteriorated food and seeds (Mehta ~~et al.~~, ~~Bodh,~~ & ~~Gupta~~, 2017). One of the important fungus known as *Candida rugosa lipases*~~*rugosa*~~ have been greatly use in biotechnology sector, because of their potential ~~(which potential?)~~ (Akoh, Lee, & Shaw, 2004). It is previously reported that *Thermomyces lanuginosus* lipases has wide range of applications in the field biotechnology and detergents. Other major lipase making fungi includes, ~~—————~~*Rhizopus*, *Mucor*, *Humicola*, *Candida*, *Geotrichum*, *Aspergillus*, *Penicillium*, *Rhizomucor* and *Rhizopus* (Gurung et al., 2013). The temperature stable lipase is produced by thermophilic *Aspergillus terreus*, *Mucor pusillus* and *Rhizopus homothallic*. A few ~~studies~~~~study~~ reported about molds with thermostable and alkaliphilic lipase (Mehta et al., 2017).

### **Applications of other different enzymes**

Therapeutic enzymes have a huge variety of selective uses such as anticoagulants or thrombolytics, oncolytics and also as replacement for various metabolic deficiencies. Proteolytic enzymes are excellent anti-inflammatory agents. There are various factors that decrease the potential of microbial source enzymes, once we enter ~~the medical~~ ~~to medical~~ sciences due ~~to the~~ ~~high~~ ~~to high~~ molecular size of ~~catalyst~~ ~~seacatalyst~~ which stop their distribution within somatic cells. In industrial processes the quantity of enzymes should be high, while in therapeutic case the purity and ~~specificity~~ ~~specify~~ should be excellent, if the quantity level is less no matter. The kinetics of such enzymes is high and low so that it is maximally efficient even at low concentration of substrate and enzymes. The source of such enzymes should be designated with high care to minimize or prevent the chances of undesirable growth by mismatched material and also to enable ready purification. In market therapeutic enzymes ~~are usually~~ ~~in usually~~ in lyophilized ~~form~~ ~~from~~ with biocompatible mannitol diluent and buffering salts. Cost of such enzymes are usually high in ~~comparison~~ ~~compare~~ to treatment or therapeutic agents. Urokinase is an ~~enzyme~~ ~~enzymes that~~ that is extracted from urine and used for blood clots as ~~an~~ ~~example~~ ~~sample~~ of such enzymes. ~~In the current~~ ~~in current~~ research area therapeutic enzymes are used in cancer treatment such as acute lymphocytic leukemia, ~~example such as~~ asparaginase enzyme has proved an excellent activity. The activity of such enzymes depend ~~upon the~~ ~~upon on~~ ~~the~~ fact that tumour cells lack aspartate-ammonia ligase activity, which inhibit/stop the synthesis of non-essential amino ~~acids~~ ~~acid~~. The asparaginase does not ~~have an impact~~ ~~have impact~~ on normal cells which are ~~capable~~ ~~accomplished~~ of manufacturing sufficient for their own necessities, but they reduce the free exogenous attentiveness, so in tumor cells they ~~cause a~~ ~~state~~ ~~cause state~~ of fatal starvation. The enzymes can be directed intravenously and ~~are only~~ ~~is~~ ~~only~~ operational in decreasing asparagine levels ~~inside the bloodstream~~ ~~inside bloodstream~~,

showing a half-life of about a day. This half-life can be enhanced by 20-fold with use of polyethylene glycol-modified asparaginase (Vellard, 2003).

[Conclusion???](#)

## REFERENCES

- Akoh, C. C., Lee, G.-C., & Shaw, J.-F. (2004). Protein engineering and applications of *Candida rugosa* lipase isoforms. *Lipids*, *39*(6), 513-526.
- Almeida, M. G., Serra, A., Silveira, C. M., & Moura, J. J. (2010). Nitrite biosensing via selective enzymes—a long but promising route. *Sensors*, *10*(12), 11530-11555.
- Anastas, P. T., Bartlett, L. B., Kirchoff, M. M., & Williamson, T. C. (2000). The role of catalysis in the design, development, and implementation of green chemistry. *Catalysis Today*, *55*(1-2), 11-22.

- Anbu, P., Gopinath, S. C., Chaulagain, B. P., Tang, T.-H., & Citartan, M. (2015). Microbial enzymes and their applications in industries and medicine 2014. In: Hindawi.
- Arreaza, G., Devane, W. A., Omeir, R. L., Sajnani, G., Kunz, J., Cravatt, B. F., & Deutsch, D. G. (1997). The cloned rat hydrolytic enzyme responsible for the breakdown of anandamide also catalyzes its formation via the condensation of arachidonic acid and ethanolamine. *Neuroscience letters*, 234(1), 59-62.
- Bartlett, G. J., Porter, C. T., Borkakoti, N., & Thornton, J. M. (2002). Analysis of catalytic residues in enzyme active sites. *Journal of molecular biology*, 324(1), 105-121.
- Burhan, A., Nisa, U., Gökhan, C., Ömer, C., Ashabil, A., & Osman, G. (2003). Enzymatic properties of a novel thermostable, thermophilic, alkaline and chelator resistant amylase from an alkaliphilic *Bacillus* sp. isolate ANT-6. *Process Biochemistry*, 38(10), 1397-1403.
- Butterworth, P. J., Warren, F. J., & Ellis, P. R. (2011). Human  $\alpha$ - amylase and starch digestion: An interesting marriage. *Starch- Stärke*, 63(7), 395-405.
- Caton, T. M., Witte, L. R., Ngyuen, H., Buchheim, J. A., Buchheim, M. A., & Schneegurt, M. A. (2004). Halotolerant aerobic heterotrophic bacteria from the Great Salt Plains of Oklahoma. *Microbial Ecology*, 48(4), 449-462.
- Djekrif-Dakhmouche, S., Gheribi-Aoulmi, Z., Meraihi, Z., & Bennamoun, L. (2006). Application of a statistical design to the optimization of culture medium for  $\alpha$ -amylase production by *Aspergillus niger* ATCC 16404 grown on orange waste powder. *Journal of Food Engineering*, 73(2), 190-197.

- Edbeib, M. F., Wahab, R. A., & Huyop, F. (2016). Halophiles: biology, adaptation, and their role in decontamination of hypersaline environments. *World Journal of Microbiology and Biotechnology*, 32(8), 1-23.
- Edner, C., Li, J., Albrecht, T., Mahlow, S., Hejazi, M., Hussain, H., . . . Steup, M. (2007). Glucan, water dikinase activity stimulates breakdown of starch granules by plastidial  $\beta$ -amylases. *Plant physiology*, 145(1), 17-28.
- Goodman, B. E. (2010). Insights into digestion and absorption of major nutrients in humans. *Advances in physiology education*, 34(2), 44-53.
- Gupta, R., Gupta, N., & Rathi, P. (2004). Bacterial lipases: an overview of production, purification and biochemical properties. *Applied microbiology and biotechnology*, 64(6), 763-781.
- Gurung, N., Ray, S., Bose, S., & Rai, V. (2013). A broader view: microbial enzymes and their relevance in industries, medicine, and beyond. *BioMed research international*, 2013.
- Guzmán- Maldonado, H., Paredes- López, O., & Biliaderis, C. G. (1995). Amylolytic enzymes and products derived from starch: a review. *Critical Reviews in Food Science & Nutrition*, 35(5), 373-403.
- Hajar-Azhari, S., Wan, W. A. A. Q. I., Ab Kadir, S., Abd Rahim, M. H., & Saari, N. (2018). Evaluation of a Malaysian soy sauce koji strain *Aspergillus oryzae* NSK for  $\gamma$ -aminobutyric acid (GABA) production using different native sugars. *Food science and biotechnology*, 27(2), 479-488.
- Kango, N., Jana, U. K., & Choukade, R. (2019). Fungal Enzymes: Sources and Biotechnological Applications. In *Advancing Frontiers in Mycology & Mycotechnology* (pp. 515-538): Springer.

Kotera, M., Okuno, Y., Hattori, M., Goto, S., & Kanehisa, M. (2004). Computational assignment of the EC numbers for genomic-scale analysis of enzymatic reactions. *Journal of the American Chemical Society*, 126(50), 16487-16498.

Kour, D., Rana, K. L., Yadav, N., Yadav, A. N., Singh, J., Rastegari, A. A., & Saxena, A. K. (2019). Agriculturally and industrially important fungi: current developments and potential biotechnological applications. In *Recent advancement in white biotechnology through fungi* (pp. 1-64): Springer.

Liu, X. Mass Spectrometry-Based Strategies in Protein Higher Order Structure Analysis: Fast Photochemical Oxidation of Proteins and Hydrogen Deuterium Exchange. *St. Louis, Missouri, 1001*, 1.

Comment [10]: Year??

Machida, M., Asai, K., Sano, M., Tanaka, T., Kumagai, T., Terai, G., . . . Kashiwagi, Y. (2005). Genome sequencing and analysis of *Aspergillus oryzae*. *Nature*, 438(7071), 1157-1161.

Mehta, A., Bodh, U., & Gupta, R. (2017). Fungal lipases: a review. *Journal of Biotech Research*, 8.

Pascoal, A., Estevinho, L. M., Martins, I. M., & Choupina, A. B. (2018). Novel sources and functions of microbial lipases and their role in the infection mechanisms. *Physiological and molecular plant pathology*, 104, 119-126.

Prakash, O., & Jaiswal, N. (2010).  $\alpha$ -Amylase: an ideal representative of thermostable enzymes. *Applied biochemistry and biotechnology*, 160(8), 2401-2414.

Rahman, N. Z. R. A., & Basri, M. (2006). *New lipases and proteases*: Nova Publishers.

Saini, R., Saini, H. S., & Dahiya, A. (2017). Amylases: Characteristics and industrial applications. *Journal of Pharmacognosy and Phytochemistry*, 6(4), 1865-1871.

- Singh, R., Kumar, M., Mittal, A., & Mehta, P. K. (2016). Microbial enzymes: industrial progress in 21st century. *3 Biotech*, *6*(2), 1-15.
- Souza, P. M. d. (2010). Application of microbial  $\alpha$ -amylase in industry-A review. *Brazilian journal of microbiology*, *41*(4), 850-861.
- Sundarram, A., & Murthy, T. P. K. (2014).  $\alpha$ -amylase production and applications: a review. *Journal of Applied & Environmental Microbiology*, *2*(4), 166-175.
- Tiwari, S., Srivastava, R., Singh, C., Shukla, K., Singh, R., Singh, P., . . . Sharma, R. (2015). Amylases: an overview with special reference to alpha amylase. *J Global Biosci*, *4*, 1886-1901.
- Vellard, M. (2003). The enzyme as drug: application of enzymes as pharmaceuticals. *Current opinion in biotechnology*, *14*(4), 444-450.
- Ventosa, A., & Nieto, J. (1995). Biotechnological applications and potentialities of halophilic microorganisms. *World Journal of Microbiology and Biotechnology*, *11*(1), 85-94.
- Verma, M. L. (2019). Biotechnological applications of lipases in flavour and fragrance ester production. In *Microbial Technology for the Welfare of Society* (pp. 1-24): Springer.