

## Minireview Article

### A mini- review : Effect of *Phyllanthus niruri* L. For Fish Growth and Health

**Comment [jt1]:** The title is not congruent with what was expressed in the work.

#### ABSTRACT

In aquaculture activities, disease attack is one of the factors causing the decline in fish production. Several types of antibiotics have been widely used to treat fish diseases. The negative impacts of the use of antibiotics include resistance to pathogenic bacteria, accumulation of antibiotics in the aquatic environment and fish bodies which are certainly harmful to human health and other aquatic biota. The use of herbal ingredients is relatively safer and cheaper to treat disease. *Phyllanthus niruri* L is one of the herbs that has been widely used to treat various diseases in humans such as intestinal infections, kidney stones, chronic liver disease, diabetes, hepatitis B, asthma, gonorrhea, bronchitis, syphilis and boosting the immune system. The ability of *P niruri* in tackling various diseases is based on its phytochemical content and pharmacological properties possessed by *P. niruri*. Several secondary metabolites and philanthine as specific compounds contained in *P. niruri* have effects as antibacterial, antioxidant, for growth, treatment and immunostimulant. So the purpose of this article is to explain *P niruri* to maintain fish health. Based on the above studies, *P. niruri* has the potential to be used as an alternative supplement for the health of various fish, because it has antibacterial, antioxidant, treatment, growth and immunostimulant effects.

Keywords : *P niruri*, disease, treatment, prevention, immunostimulant.

#### 1. INTRODUCTION

The use of drugs derived from herbal ingredients can be used as an alternative to replace antimicrobial agents that have a toxic effect and cause microbial resistance to the antibiotics used [1]. Plants contain several compounds that have pharmacological properties, namely as antimicrobials [2,3]. Apart from being an antimicrobial compound, the compounds contained in plants also have an antifungal, antioxidant, antiviral [4] and immunostimulant [5, 6] effect which are of course very useful for human and animal health. In general, plants contain several secondary metabolites, namely alkaloids, flavonoids, steroids, saponins, terpenoids and tannins that can play a role in fighting various diseases [5,7].

*Phyllanthus niruri* L. belongs to the Euphorbiaceae family, is a tropical plant spread throughout the world, known as a medicinal plant with many benefits [8,9]. In Brazil *P. niruri* is known as Chanca Piedra, in southern India it is called Bhumyamalaki, in Malay it is called Dukong Anak and in Chinese it is called zhu zi cao [9]. Traditionally this plant has been widely used by people in tropical countries to treat various diseases including intestinal infections, kidney stones [10, 11], chronic liver disease, diabetes, and prevent hepatitis B virus infection [11 12], asthma, gonorrhea, bronchitis, and syphilis [13,14] and enhances the immune system [15]. *P. niruri*'s ability to treat various diseases is due to its pharmacological

properties, such as immunomodulator, antiviral, antibacterial, diuretic, anti-hyperglycemic and hepatoprotector [9, 16]. In Indonesia, this plant is used to boost the immune system. This plant extract has been shown to be an immunomodulator that can increase peripheral blood proliferative activity and macrophage phagocytic activity [15].

Based on the phytochemical content and pharmacological properties of *P niruri*, from several research results, the plant can not only be used for the treatment of diseases in humans, but also for the treatment of diseases in animals including fish. Therefore, the purpose of writing this article is to describe the *P. niruri* plant in an effort to help manage fish health, by looking at the phytochemical content of the plant, its antibacterial, antioxidant and growth effects as well as the immunostimulant effect on cultured fish that has been carried out by several researchers.

## 2. PHYTOCHEMICAL SCREENING OF *Phyllanthus niruri* L.

Many studies have been carried out on the chemical components of the plant *Phyllanthus niruri* L. Based on the results of qualitative phytochemical tests on *P. niruri* L. extract using different solvents produced several different types of compounds. But overall the compounds contained in the extract of *Phyllanthus niruri* L are alkaloids, flavonoids, tannins, saponins, steroids, phenols and terpenoids (Table 1).

Table 1. Phytochemical Screening of *Phyllanthus niruri* L.

solvents extract	Alkaloids	Flavonoids	Tannins	Saponins	Steroids	Phenols	Terpenoids	Ref
Ethanol 70%	+	+	+		+			[17]
Methanol 80%	+	+		+		+	+	[18]
Ethanol 96%	+	+	+	+				[19]
Ethanol 95%		+	+	+	+	+		[20]

According to several studies, *P. niruri* L. in addition to containing several bioactive molecules of flavonoids, tannins, alkaloids, steroids, also contains lignans, phyllanthin, hypophyllanthin, glycosides, ellagitannins, triterpenes, phenyl propanoids, ricinolic acid, niruriside and phyltetralin [21, 22, 23]. According to Sudarsono (1996) [25] the *P. niruri* L. plant also contains essential oils, anthraquinones and arbutin. The leaves contain components of flavonoid compounds such as quercetin. The stem contains components of niruri, niruritenin, rutin. Lignin components such as phyllanthine, hypophilantin are present in all parts of the plant [26]. Triterpenoid compounds contain components of lupeol acetate and betasitosterol [27]. Based on the results of quantitative phytochemical tests, the total phenol content of the plant *P. niruri* L is equivalent to 28.05 g of gallic acid in 1 mg of plant extracts, while the flavonoid content in 1 mg of plant extracts is equivalent to 61.41 g of quercetin, while the flavonoid content in 1 mg of plant extracts is equivalent to 61.41 g of quercetin.

**Comment [jt2]:** The information presented is very good and pertinent to the subject of *P.nururi*, but I do not see where it relates to use in fish.

70% ethanol extract of *P. niruri* was 0.864% [17]. The Phytochemical analysis of the methanol extract revealed that *P. niruri* contains mainly flavonoids, alkaloids, tannins, saponins, coumarins, polyphenols, terpenoids and steroids compounds. [28, 29].

The type of solvent used for extraction affects the levels of secondary metabolites produced. *P. niruri* extracted using different solvents, namely 60% methanol, 60% ethanolacetone and 60% isopropanol produced different levels of secondary metabolites. The highest total flavonoids were obtained using 60% ethanol extract solution, which was 269.26±1.21 mg QE/g, while those using 60% methanol, 60% acetone and 60% Isopropanol solvents each produced 209.05±0.75, 247.60±1.23 and 212.70±1.14 mg QE/g. 60% acetone solvent resulted in the highest total phenol, tannin and antioxidant activity, respectively 188.77±1.05 mg GAE/g, 297.51±1.20 mg TAE/g, and 71.99±0.42 % [30]. The results of this study indicate that acetone is effectively used to extract *P. niruri* so as to produce the highest total phenol. This indicates that the polarity of the phenolic compounds of *P. niruri* extract corresponds to the polarity of acetone. Compounds obtained from extraction are based on the similarity of polarity to the solvent used [31].

### 3. ANTIBACTERIAL EFFECT OF *Phyllanthus niruri* L.

The phenolic compounds, flavonoids, saponins, alkaloids, and terpenoids contained in *Phyllanthus niruri* have antibacterial activity against several bacteria that can infect cultured fish, such as *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Aeromonas hydrophila* and *Edwardsiella tarda* [18, 32,19 ,33].

The results of the inhibitory test of *P. niruri* solution using paper discs against the bacteria *Edwardsiella tarda*, showed a concentration of 50,000 ppm produced the highest average inhibition, which was 14.3 mm, while at concentrations of 25,000 ppm, 10,000 ppm and 5,000 ppm each produced average drag of 12 mm, 8.7 mm and 6 mm [34]. Methanol extract 80% *P. niruri* at a concentration of 50µg/ml (100µl) can inhibit the growth of *Staphylococcus aureus* bacteria with an inhibition zone diameter of 13 mm, while for *Escherichia coli* and *Pseudomonas aeruginosa* bacteria the diameter of the inhibition zone is 10 mm and 12 mm, respectively. [18]. The concentration of 50µL of *P. niruri* extract using various solvents can inhibit the growth of *Aeromonas hydrophila* bacteria. Chloroform solvent produced an inhibition zone diameter of 12 cm, while with ethanol, ethyl acetate and petroleum ether the same inhibitory zone was 16 mm [32] (Table 2). From this, it can be seen that 50µL *P. niruri* extract with ethanol, ethyl acetate and petroleum ether as solvents was more effective in inhibiting the growth of *A. hydrophila* bacteria. Meanwhile, 80% methanol extract of *P. niruri* at a concentration of 50µg/ml (100µl) was more effective in inhibiting the growth of *Staphylococcus aureus*.

Table 2. Antibacterial activity of *Phyllanthus niruri*

extract solvents	Concentration Extract	Type of bacteria	Inhibition zone (mm)	Ref.
boiled solution	50.000 ppm	<i>Edwardsiella tarda</i>	14,3 mm	[34]
Methanol 80%	50µg/ml (100µl)	<i>Staphylococcus aureus</i>	13	[18]

Methanol 80%	50µg/ml (100µl)	<i>Escherichia coli</i>	10	[18]
Methanol 80%	50µg/ml (100µl)	<i>Pseudomonas aeruginosa</i>	12	[18]
Chloroform	50µL	<i>Aeromonas hydrophila</i>	12	[32]
Ethanol	50µL	<i>Aeromonas hydrophila</i>	16	[32]
Ethyl Acetate	50µL	<i>Aeromonas hydrophila</i>	16	[32]
Petroleum Ether	50µL	<i>Aeromonas hydrophila</i>	16	[32]
boiled solution	500.000 ppm	<i>Aeromonas hydrophila</i>	14,66	[35]
boiled solution	50.000 ppm	<i>Aeromonas hydrophila</i>	12	[35]
boiled solution	5.000 ppm	<i>Aeromonas hydrophila</i>	9	[35]

Thin layer chromatography results showed that *P. niruri* ethanol extract was able to inhibit the growth of *Staphylococcus aureus* bacteria with an Rf value of 0.46. Analysis with infra red spectrophotometer showed that the ethanol 96% extract has hydroxyl (-OH) and carbonyl (C=O) functional groups. This shows that the compounds that play a role in inhibiting *S. aureus* are alkaloids and tannins [19]. The results of the antibacterial test of *P. niruri* extract against *E. tarda* bacteria using the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) test showed that the extract concentration of 0.0313 g/ml was the best concentration, because it was the lowest concentration that could inhibit growth of *E. tarda* bacteria [25]. Quercetin is a flavonoid compound of the flavonol type which has antimicrobial and antiviral roles [36]. Phenols, flavonoids, and tannins are the main class of phytochemicals that have antimicrobial activity [37].

#### 4. TREATMENT EFFECTS OF *Phyllanthus niruri* L. ON FISH

The results of the LD<sub>50</sub> test of *P. niruri* solution on *Clarias gariepinus* fish showed that at concentrations of 40,000 and 60,000 ppm with a 24-hour immersion time it caused 10% death. When tested to treat *Clarias gariepinus* fish infected with *Edwardsiella tarda* with a density of 1 ml x 10<sup>-9</sup> bacterial cells/ml/100 g of fish weight through immersion for 30 minutes, it turned out that a concentration of 7000 ppm was quite effective, with a survival rate of 86, 67% [34]. Based on the in vivo test results, a solution of *P. niruri* with a concentration of 5,000 mg/L can treat *Cyprinus carpio* fish infected with *Aeromonas hydrophila* bacteria through immersion. The effective immersion time was 5 hours, with the highest survival rate of 76.67% [35].

## 5. ANTIOXIDANT EFFECT OF *Phyllanthus niruri* L.

Several secondary metabolites contained in the *P. niruri* plant, apart from being very useful for treating various diseases, can also be used as antioxidants [38]. Flavonoid compounds are one of the secondary metabolite compounds contained in the *P. niruri* plant, besides having antithrombotic and anti-inflammatory effects, they also have antioxidant and anti-radical effects [39]. Phenols and flavonoids have a significant effect as antioxidants.

Quercetin is a flavonoid compound of the flavonol type [36] which is a strong anti-radical, able to fight free radical damage, is also a strong anti-radical against peroxy radicals, superoxide anions, and hydroxyl and is able to scavenge free radicals directly [40]. The results of the total antioxidant activity test on *P. niruri* extract showed significant results. A total of 1 mg/ml of *P. niruri* extract showed total antioxidant with antioxidant activity equivalent to 216.28 g of ascorbic acid which was further confirmed by the DPPH radical scavenging test. The antioxidants contained in *P. niruri* extract were able to reduce DPPH. The IC<sub>50</sub> value of the growing extract *P. niruri* produced for scavenging activity of DPPH free radicals was 10.53 g, while for ascorbic acid it was 8.90 g. As much as 1 mg of *P. niruri* for its antioxidant activity its strength is comparable to 152 ascorbic acid. Here, *P. niruri* extract shows high antioxidant activity and is able to reduce ferric chloride and scavenge DPPH radicals. The ability of this plant extract is based on several secondary metabolite compounds contained in these plants including phenolic compounds and flavonoids. Based on the results of the study showed the higher the amount of phenols and flavonoids, the higher the antioxidant activity and scavenge free radicals [18].

## GROWTH EFFECT OF *P. niruri* L

Research has shown that several medicinal plants have been successfully used as supplements to trigger growth and feed conversion in fish and shrimp, including garlic [41], turmeric [42], black cumin [43], Aloe vera [44], *Andrographis paniculata* [45] and so on. Likewise with *P. niruri* apart from being an antibacterial, antioxidant and immunostimulant, it can also trigger fish growth. As research conducted by Sunitha (2017) [46] showed carp (*Cyprinus carpio*) that had been fed a diet mixed with *P. niruri* powder for 60 days showed better growth compared to controls. Among the treatments given, feeding mixed with *P. niruri* powder as much as 2% (10 g/ 500 g) showed the highest fish growth rate in terms of weight parameters (Weight :  $1.15 \pm 0.16$  g, growth :  $0.55 \pm 0.04$  g and SGR :  $0.92 \pm 0.07\%$  and length ( $3.75 \pm 0.24$  cm). While the fish fed with *P. niruri* powder at 1% (5g/500 g), resulted in lower weight and length, namely weight :  $1.05 \pm 0.09$ , growth :  $0.49 \pm 0.03$  and SGR :  $0.82 \pm 0.07\%$  and the length is  $3.53 \pm 0.25$  cm. *Phyllanthus niruri* as an additive mixed into feed, with a duration of administration for 60 days can increase the growth of *Carassius auratus* fish. Experimental results show a concentration of 1.5 percent produces the highest average weight gain, which is 1.769 g [47].

## 5. EFFECT OF *Phyllanthus niruri* L. AS AN IMMUNOSTIMULANT

Research on the effects of *P. niruri* as an immunostimulant that can increase the body's resistance in both humans and animals to disease attacks has been carried out. Observation of

antibody titer is one of the parameters to determine the effectiveness of a substance that acts as an immunostimulant, by looking at the ability of serum proteins containing antibodies to collect and destroy antigens that enter the body [48]. Based on the results of the study, *Oreochromis mossambicus* fish which had been treated with aqueous extract of *P. niruri* leaves at various doses, then challenged using sheep red blood cells (SRBC) as an antigen, showed an increase in the response of the primary antibody titer.

The highest significant increase occurred at a dose of 20 mg. On day 15 the secondary antibody response was significantly at all doses (20, 2, and 0.002 mg) resulting in a higher value when compared to the control. As for the neutrophil activation test using hot aggregated serum albumin (HA BSA), the results obtained showed that all doses could increase neutrophil activity, but the highest number of activated neutrophils was in fish given *P. niruri* leaf aqueous extract of 0.02 mg, followed by 0.02 mg. 0.2 and the lowest was given the extract at a dose of 0.002 mg [49]. Neutrophils are part of the white blood cells that are part of the immune system. In addition to neutrophils, the white blood cells that are part of the immune system are monocytes, lymphocytes and macrophages. While cytokines, antibodies, and complement proteins are included in immune molecules. Synchronous interactions between the immune system and immune molecules can induce a good immune response, which can lead to resistance to pathogenic infections [50]. The results of research conducted by Ma'at et al. (1996) [51] showed *P. niruri* in vitro to have an immunomodulating effect. *P. niruri* extract (50–200 mg/kg) can elicit both cellular and humoral immune responses. As the research results of Nworu et al. (2010) [52] showed an aqueous extract of *P. niruri* (12.5–200 g/ml) could induce lymphocytes and macrophages in experimental animals.

Tilapia (*Oreochromis niloticus*) which had been given *P. niruri* leaf extract by injection was able to survive the attack of *Aeromonas hydrophila* bacteria. concentration of 50 mg was significantly the best concentration by producing the highest tilapia survival, which was 76.7% [53].

*Cyprinus carpio* L. after being given feed mixed with *P. niruri* powder for 60 days then challenged with *A. hydrophila* bacteria with a density of  $10^3$  CFU, able to withstand the attack of these bacteria. Giving *P. niruri* powder as much as 2% (10 g/500 g of feed) showed the highest resistance of fish to attack by *A. hydrophila* bacteria, which was 100% [46].

The POM (2006) [54] stated that *P. niruri* is one of the immunomodulatory drugs, namely drugs that can improve or enhance the immune system. Compounds that play a role in this are the flavonoid group which is the main component in *P. niruri*. Flavonoids act on body cells by sending intracellular signals to cell receptors, so that cells work optimally. Compounds contained in meniran, including phyllanthin and hyphophyllanthin are two compounds that have anti-inflammatory activity so that they can strengthen immunity.

*P. niruri* has an immunomodulatory effect through activation and augmentation of the cellular immune system. Specifically, *P. niruri* can activate neutrophils, macrophages, monocytes, T and B lymphocytes. The active process of phagocytosis by neutrophils indicates an acceleration of the process of eradicating pathogenic microbes, especially for extracellular pathogens, such as viruses, bacteria, or fungi. *P. niruri* can induce an increase in monocytes and macrophages, which are able to lyse infecting intracellular pathogenic cells and expose these pathogenic cells to other immune components in the extracellular compartment. In addition, *P. niruri* can modulate cytokine secretion, including stimulating IFN- $\gamma$ , TNF- $\alpha$ , IL-4,

**Comment [jt3]:** With what is expressed in this part of the work, the study of *P. niruri* is very good information, but its application in fish is very poor and has no sustenance.

IL-6, IL-12, and suppression of IL-10. This suggests that *P. niruri* can inhibit the body's defenses, by engaging the cellular immune system against foreign pathogens [55] .

## CONCLUSION

*P. niruri* has the potential to be used as an alternative supplement for the health of various fish, because it has antibacterial, antioxidant, treatment, growth and immunostimulant effects

## NOTE:

The study highlights the efficacy of "Herbal" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

## REFERENCES

1. Rashid S, Rather MA, Shah WA, Bhat BA. Chemical composition, antimicrobial, cytotoxic and antioxidant activities of the essential oil of *Artemisia indica* Willd. *Food Chem.* 2013;138(1): 693-700.
2. Mydeen KP, Haniffa MA. Evaluation of antibacterial activity of medicinal plants on fish pathogen, *Aeromonas hydrophila*. *Journal of Research on Biology.* 2011; 1:1-5.
3. Turker, H., A. B. Yildirim and F. P. Karakas (2009). Sensitivity of bacteria isolated from fish to some medicinal plants, *Turkish J. Fish. Aqu. Sci.*, 9 : 181-186.
4. Kolkovski, S. and J. Kolkovski (2011). Herbal medicine in aquaculture. *International Aquafeed*, 14(2) : 28-31.
5. Pandey, G. and S. Madhuri, 2010. Pharmacological activities of *Ocimum sanctum*: A review. *Int. J. Pharm. Sci. Rev. Res.*, 5: 61-66.
6. Yin G, Cao L, Xu P, Jency G, Nakao M. Hepatoprotective and antioxidant effects of *Hibiscus sabdariffa* extract against carbon tetrachloride - induced hepatocyte damage in *Cyprinus carpio* In vitro *Cell. Dev. Biol. Anim.* 2011; 47(1):10-15.
7. Ravikumar S, Selvan GP, Gracelin AA. Antimicrobial activity of medicinal plants along Kanyakumari coast, Tamil Nadu, India. *African Journal of Basic Applied Sciences.* 2010; 2:153-157.
8. Dirjomuljono M, Tjandrawinata RR. Clinical trials involving *Phyllanthus* species. In: Kuttan R, Harikumar KB, editors. *Phyllanthus species: scientific evaluation and medicinal applications.* Boca Raton: CRC Press; 2011, p. 289-313.
9. Lee NY, Khoo WK, Adnan MA, Mahalingam TP, Fernandez AR, Jeevaratnam K. The pharmacological potential of *Phyllanthus niruri*. *J Pharm Pharmacol* 2016; 68: 953-69.

**Comment [jt4]:** t must homogenize the wording of the references.

**Comment [jt5]:** The references on the subject of fish is almost nil, when the work is referred to these organisms.

10. Bieski IG, Leonti M, Arnason JT, Ferrier J, Rapinski M, Violante IM, Balogun SO, Pereira JF, Figueiredo Rde C, Lopes CR, da Silva DR, Pacini A, Albuquerque UP, Martins DT. Ethnobotanical study of medicinal plants by population of Valley of Juruena Region, Legal Amazon, Mato Grosso, Brazil. *Journal Ethnopharmacol.* 2015; 173: 383-423.
11. Patel JR, Tripathi P, Sharma V, Chauhan NS, Dixit VK. *Phyllanthus amarus*: ethnomedicinal uses, phytochemistry and pharmacology: a review. *Journal Ethnopharmacol.* 2011; 138: 286-313.
12. Qi FH, Wang ZX, Cai PP, Zhao L, Gao JJ, Kokudo N, Li AY, Han JQ, Tang W. Traditional Chinese medicine and related active compounds: A review of their role on hepatitis B virus infection. *Drug Discov. Ther.* 2013; 7(6): 212- 224.
13. Chopra, R. N., Nayar, S. L., Chopra, I. I., Glossary of Indian medicinal plants. Ranchi: Catholic Press, 1986.
14. Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN, Ray C. Screening of Indian plants for biological activity: I. *Indian Journal of Experimental Biology.* 1968,. 6(4), 232–247.
15. Putri DU, Rintiswati N, Soesatyo MH, Haryana SM. Immune modulation properties of herbal plant leaves: *Phyllanthus niruri* aqueous extract on immune cells of tuberculosis patient-in vitro study. *Natural product Research.* 2018; 32: 463–467.
16. Tropical Plant Database. Chanca piedra (*Phyllanthus niruri*). 2013. [Online] Available from: <http://www.rain-tree.com/chanca.htm#>. V3nOBfI97IU [Accessed on 4th July, 2016]
17. Alegantina S, Setyorini HA, Triwahyuni. Health Research and Development Agency, Ministry of Health RI in Health Research Bulletin Health Research Bulletin. 2015; 43.
18. Ramandeep K, Nahid A, Neelabh C, Navneet K. Phytochemical Screening of *Phyllanthus niruri* collected from Kerala Region and its Antioxidant and Antimicrobial Potentials. *Journal Pharmaceutical Sciences & Research.* 2017; 9(8) :1312-1316.
19. Mangunwardoyo W, Cahyaningsih E, Usia T. Extraction and Identification of Meniran Herb Antimicrobial Compounds (*Phyllanthus niruri* L.). *Indonesian Journal of Pharmaceutical Sciences.* 2009; 7(2) : 57-63.
20. Rivai, H., Refilia S., Agusri, B. Characterization of Meniran Herb Extract (*Phyllanthus niruri* Linn) by Fluorescence Analysis. *Higea Pharmaceutical Journal.* 2013; 5(2) : 15-22.
21. Calixto JB, Santos AR, Cechinel Filho V, Yunes RA (1998) A review of the genus *Phyllanthus*: their chemistry, pharmacology, and therapeutic potential. *Med Res Rev* 18: 225.

22. Unander DW, Webster GL, Blumberg BS. Usage and bioassays in *Phyllanthus* (Euphorbiaceae). IV. Clustering of antiviral uses and other effects. *J Ethnopharmacol.* 1995;45(1):1–18. doi:10.1016/0378- 8741(94)01189-7.
23. Li Xiang-rong. Chemical components and bioactivities of *Phyllanthus niruri* L. Tianran Chanwu Yanjiu Yu Kaifa. 2007;19:890.
24. Van Dau, N. Ha TTT. Chemical composition of *Phyllanthus niruri* L., Euphorbiaceae. *Tap Chi Duoc Hoc -Saigon Then Hanoi*,2007;47:15-18.
25. Sudarsono, P. A., Gunawan, D., Wahyuono, S., Donatus, I. A., & Drajad, M. (1996). *Medicinal Plants*. Yogyakarta: Research Center for Traditional Medicine, University of Gajah Mada.
26. Gupta, and Ahmed B and Shoyakugaku.Z. A new flavones Glycoside from *phyllanthus niruri* . *J. Nat. Prod.* 1984; 4: 213-215.
27. Sinh, S.K.P. Agarawal and dogra J. V, Variotionis the level of vitamin C. Total Phenolic and Protein in *Phyllantus niruri* L, During leaf mutarationn. *Natl. Acad. Sel. Latt* . 1989; 4 (12): 467-469.
28. Mamta S. Uparkar, Sunil H. Ganatra. Qualitative phytochemical screening and identification of phytoconstituents from *Phyllanthus niruri* Linn. By GC-MS. *Research Journal Pharmasi and Technology.* 2020; 13 (8):
29. Bagalkotkar G, Sagineedu SR, Saad MS, Stanslas J. Phytochemicals from *Phyllanthus niruri* Linn. and their pharmacological properties: a review. *J Pharm Pharmacol.* 2006;58(12):1559–70.
30. Rachmawati RA, Wisaniyasa NW and Suter IK. The Effect of Different Solvents on The Antioxidant Activity of Gale of The Wind Extract (*Phyllanthus niruri* L.). *Jurnal Itepa.* 2020; 9 (4) : 458-467.
31. Harborne, J. B. (1982). *hytochemical Methods Guiding Modern Ways of Analyzing Plants*. Second Edition. Padmawinata Translation. K., and Soediro. I. ITB Bandung Publisher.
32. Thiyagarajan P, Lakshmi AB, M.G. Ebbie, G .Chandra. A study on the control of *Aeromonas hydrophila* infection in the cat fish by medicinal plants. *Scholars Academic Journal of Biosciences* 2014; 2(2): 144-150.
33. Sudarno, Setiorini FA dan Suprpto H. Effectivity Of Meniran (*Phyllanthus niruri*) Extract as Edwardsiella tarda Antibacterial According In Vitro. *Scientific Journal of Fisheries and Marine.* 2011; 3 (1) : 103-108.
34. Setiaji J, Johandan TI and Pramujiono A.. Test Solution *Phyllanthus niruri* for the Treatment of *Clarias gariepinus* the infected Bacteria Edwardsiellatarda. *Jurnal Dinamika Pertanian.* 2013; 28 (2) : 161-166.
35. Ifitah D. 2006. The effectiveness of meniran simplicia (*Phyllanthus niruri* L.) for the treatment of diseases caused by *Aeromonas hydrophila* bacterial infection in carp (*Cyprinus carpio*) through immersion. Final Practical Scientific Work. Diploma 4

Program in Aquaculture Technology Studies. Department of Aquatic Resources Management Technology. Fisheries College. Jakarta. 87 pp.

36. Robinson, T. (1995). High Plant Organic Content. Bandung. ITB Publisher.
37. Cowan, M. M. Plant Products as Antimicrobial Agents. Clin Microbiol Rev. 1999 Oct, 12(4), 564–582. Kaur Ramandeep et al /J. Pharm. Sci. & Res. Vol. 9(8), 2017, 1312-1316.
38. Dalimartha, dr. S. (2000). Atlas of Indonesian Medicinal Plants Volume 2. (E. Priyatini, Ed.). Jakarta: Trubus Agriwidya.
39. Winarsi, H. (2007). Natural Antioxidants and Free Radicals. Yogyakarta: Kanisius. Journal of Mathematics and Science Network. 2019; 1 (1).
40. Robak, J., Gryglewski. Flavanoids are Scavengers of Superoxide Anions. Biochemical Pharmacology. 1988; 37(5).
41. Marentek GA, Manoppo H, Longdong SNJ. Evaluation of The Use of Garlic (*Allium sativum*) in Enhancing Nonspecific Immune Response and Growth of Nile Tilapia (*Oreochromis niloticus*). Budidaya Perairan. 2013 ;1 (1) : 1-7.
42. Santika L, Diniarti N, Astriana BH. The Effect Of Addition The Turmeric Extracton Pellet Feed to Growth And Feed Utilization Efficiency of White Barramundi (*Lates calcarifer*). Marine Journal. 2021; 14 (2): 48-57.
43. Lei S and Xiao-En DC. Effect of *Nigella sativa* on growth and survival rate of *Penaeus vannamei*. International Journal of Fisheries and Aquatic Studies 2019; 7(4): 406-410.
44. Khanal M, Lamichhane S, Bhattarai A, Kayastha BL, and Labh SN. Extract of Aloe vera (*Aloe barbadensis* Miller) Enhances the Growth, Protein Contents, and Gastroscopic Index (GaSI) of Common Carp *Cyprinus carpio*. Hindawi Journal of Nutrition and Metabolism. Volume 2021. 14 pages.
45. Maiti S, Saha S, Jana P, Chowdhury A, Khatua S & Ghosh TK. Effect of dietary *Andrographis paniculata* leaf extract on growth, immunity, and disease resistance against *Aeromonas hydrophila* in *Pangasianodon hypophthalmus*. Journal of Applied Aquaculture. 2021; 33( 3).
46. Sunitha C, Mettilda S and Vinoliya J. Effect of dietary intake of *Phyllanthus niruri* L. on fingerlings of freshwater fish, *Cyprinus carpio* L. International Journal of Fisheries and Aquatic Studies. 2017; 5(1): 352-359.
47. Ahilan, B., A. Nithiyapriyatharshini and K. Ravaneshwaran. Influence of certain herbal additives on the growth, survival and disease resistance of goldfish, *Carassius auratus* (Linnaeus). Tamilnadu J. Vet. Ani. Sci. 2010. 6(1): 5-11.
48. Subowo. Immunobiologi. Immunobiologi. Edisi 2. Angkasa Publisher. Bandung 2009. Sagung Seto. Jakarta.
49. Muthulakshmi M, Subramani PA and Michael RD. Immunostimulatory effect of the aqueous leaf extract of *Phyllanthus niruri* on the specific and nonspecific immune

- responses of *Oreochromis mossambicus* Peters. Iranian Journal of Veterinary Research. 2016; 17(3) Ser. 56 : 200-202.
50. Chaplin, D. D. (2010). Overview of the immune response. J. Allerg. Clin. Immunol. 125, S3–S23. doi: 10.1016/j.jaci.2009.12.980.
51. Ma'at, S. (1996). *Phyllanthus niruri* L. as an immunostimulator in mice. Diss., University of Airlangga, Surabaya.
52. Nworu C, Akah P, Okoye F, Proksch P and Esimone C. The effects of *Phyllanthus niruri* aqueous extract on the activation of murine lymphocytes and bone marrow-derived macrophages. Immunol. Invest. 2010; 39: 245–267.
53. Wulandari R. Effectiveness of Meniran (*Phyllanthus niruri*) Leaf Extract by injection for the prevention of bacterial infection (*Aeromonas hydrophila*) in Tilapia (*Oreochromis niloticus*).2014; Bachelor Thesis, Universitas Muhammadiyah Purwokerto.
54. Food and Drug Supervisory Agency of the Republic of Indonesia. Meniran; The Latest Scientific Data Series on Medicinal Plants. Directorate of Native Indonesian Medicines Deputy for Supervision of Traditional Medicines, Cosmetics and Complementary Products. Jakarta. 2006. Tjandrawinata RR, Maat S dan Noviaranya D. Effect Of Standardized *Phyllanthus niruri*. L Extract on Changes In Immunologic Parameter : Correlation Between Preclinical and Clinical Studies. Medika. 2005; 31 (6) : 367-371.