

Original Research Article

Increasing Growth Performance and Omega-3 of Pangasius Catfish with Fish Oil Administration

ABSTRACT

This research aims to examine the effect of adding fish oil on the growth performance and content of Omega-3 fatty acids in Pangasius catfish (*Pangasius* sp). Pangasius catfish with an average weight of 11.9-12-71 g were reared in a fiber tank with a water volume of 240 liters stocked filled with 15 individual fish, with giving a stocking density of 10 fish⁻¹ m². The fish were reared -reared for 28 days. This study used a completely randomized design with five treatments of adding fish with fish oil supplemented to the feed at 0%, 3%, 4%, 5%, and 6% with each treatment consisting of three replications. The fish were Feed was given fed twice a day at 08:00 am and 16:00 pm West Indonesian time, with a dose of at a feeding level of 3% of the fish biomass per day. The results showed that the application of fish oil in the feed had a different effect (P<0.05) on the specific growth rate (1.69-2.07%), and the efficiency of feed (69.66-85.85%). Feeding with a concentration of 4% fish oil increased the fat content of the treatment without the addition of fish oil from DHA 24 mg⁻¹ 100 and EPA 7.3 mg⁻¹ 100 gr to EPA 30.7 mg⁻¹ 100 and DHA 10.6 mg⁻¹ 100 gr. The water quality parameter values during the research were in optimal condition for the growth and survival of Pangasius catfish.

Keywords: Pangasius Catfish; Fish Oil; Specific Growth Rate; Feed Efficiency; Omega-3.

1. INTRODUCTION

Pangasius catfish is a type of freshwater consumption fish that has good prospects because of its high selling value and easy cultivation. Pangasius catfish processing can use the skin, meat and bones such as fillets, meatballs, dumplings, nuggets, sausages, bone nuggets, shredded, and skin crackers. The demand for processed pangasius catfish is also quite large in Indonesia. This is triggered by the large and widespread and development of aquaculture [35]. Pangasius catfish have benefits as a source of a high supply of animal protein and as an ornamental fish. The protein content of pangasius catfish in 159 g fillet is 24.7 g, the value of meat protein reaches 14.53%, fat 1.03%, ash 0.74%, and water 82.22%, and is safer for health because it has higher cholesterol levels, lower (21-39 mg/100 g) compared to livestock meat, so it is good for a diet program [35].

Fatty acids are classified based on their structure, carbon-chain length (short, medium, or long), degree of saturation (number of double bonds), and location of double bonds [21]. Classification of the double bond, fatty acids are divided into two parts, namely saturated fatty acids and unsaturated. Omega-3 fatty acids are polyunsaturated fatty acids that have many double bonds and are derivatives of their precursors, namely the essential fatty acids linoleic and linolenic. Essential fatty acids cannot be formed by the body, so they must be supplied from food. The precursor enters the elongation and desaturation process which

Comment [ma1]: Correct this weight

Formatted: Not Highlight

Comment [ma2]: This statement needs to be better phrased

Comment [ma3]: The authors needs to make this statement clear, precise and a true reflection of the results.

Comment [ma4]: The water quality was not among the factors to be investigated (please see the title). It is only growth performance and omega 3 concentrations that were planned to be studies. Therefore, the authors should exclude water quality from the abstract

Formatted: Not Highlight

Comment [ma5]: The grammar needs to be corrected

Comment [ma6]: Gramatical error here also

Comment [ma7]: There is need for more references

produces three forms of omega-3 fatty acids, namely LNA/ALA (alpha-linoleic acid (C1:3, n-3)), EPA (eicosapentaenoic acid (C20:5,n-3)), and DHA (docosahexaenoic (C22:6, n-3) [6].

Apart from being used as an energy source, fat also functions as a source of essential fatty acids [10]. Essential fatty acids are fatty acids that cannot be synthesized by the body and come from the fish food source itself, so they need to be added to the feed. Fish oil is a source of essential fatty acids. Fatty acids play an important role in supporting metabolic activities, precursors of several prostanoids, substrates for the formation of lipoygenase, membrane components, and main precursors for the formation of leukotrienes [15]. Fat can also be used as a protein-sparing *effect* in forming tissue so that the use of protein can be optimized as a source of growth for fish [48].

One of the main components of food that has a positive impact on health is fat that has multifunctionality, namely as the largest energy contributor (30% or more of the total energy needed by the body) and is a source of essential fatty acids linoleic and linolenic [10]. In addition, it is used as an energy source, part of cell membranes, mediator of biological activity between cells, insulator in maintaining the balance of body temperature, protecting body organs and solvents for vitamins A, D, E, and K. Omega-3 unsaturated fatty acids, It plays an important role in the morphological, biochemical, and molecular development of the brain and other organs. The deficiency of omega-3 fatty acids caused by inadequate intake or due to diseases that reduce absorption can inhibit brain development, physical health, and environmental interactions and have a strong effect on the formation of cognitive development [14].

2. METHODS

This research was carried out for 28 days starting in October 2021 at the Faculty of Fisheries, Padjadjaran University. The research used was experimental with a Completely Randomized Design (CRD). Parameters *specific growth rate* (SGR) and feed efficiency were tested using analysis of variance/F test and further tests using Duncan's test to see the effect of treatment on each parameter to be tested, at a 95% confidence level. Water quality parameters were analyzed descriptively.

Treatment A: Control (without the addition of fish oil), Treatment B: Addition of 3% fish oil, Treatment C: Addition of 4% fish oil, Treatment D: Addition of 5% fish oil, and Treatment E: Addition of 6% fish oil

2.1 Test Feed and

Preparation of the test feed was started by mixing the feed with fish oil according to the treatment, namely 0%, 3%, 4%, 5% and 6%. The feed used was Hi Provite PF 781 brand. The feed mixed with fish oil was added with 5 g of progol which functions as a feed adhesive and fish oil. Then the feed ~~is was~~ dried until ~~the feed is ensured that~~ the fish oil ~~is was~~ glued and ~~does did~~ not come off the feed. After the feed ~~is was~~ dry, the feed is put in a plastic ziplock and stored in a dry place. ~~The fish were fed on~~ ~~Feed was given as much as~~ 3% of ~~fish their~~ biomass ~~during the study~~ daily.

Comment [ma8]: The introduction should end with a statement on what is the objective of the study. In addition, the hypothesis to be tested in this study needs to be stated

Comment [ma9]: Use past tense when describing the materials and method

Comment [ma10]: 28 days is a short period of time to enable long term effects to be noted in growth performance and omega 3 accretion.

Comment [ma11]: The authors have only specific growth rate and feed efficiency, these are too few parameters. To make the manuscript stronger, I would recommend that the authors include more growth and nutrient utilisation efficiency parameters.

Comment [ma12]: Data analysis should be stated at the end of the materials and methods section

Comment [ma13]: This should be trasfered to the section on test feed

Comment [ma14]: State the fish species that the Hi Provite is meant for?

2.2 Pisciculture

The test fish used in this study was Pangasius catfish from Purwakarta, West Java. Pangasius catfish sized 9-11 cm with an average weight of 12.41 ± 0.43 g/head and the stocking density used was 1 m²10. The fish are adapted to the new environment first and the feed will be given in a fiber tank with a volume of 450 liters. Then transferred to the rearing container as many as 15 test fish per fiber tub with a volume of 240 liters which had previously been weighed. Feeding as much as 3% of fish biomass was carried out at 08:00 West Indonesian time and 16:00 West Indonesian time. Siphoning is carried out every day, while water changes are carried out 4 times a day as much as 35% of water with water that has been deposited first. Maintenance was carried out for 28 days of observation. At the beginning and end of rearing several fish samples were taken for proximate analysis and fatty acid omega-3 analysis.

Comment [ma15]: There are grammatical errors in this section. Please correct them. There are also redundant words that need to be eliminated. For example as much as.....

Comment [ma16]: Correct this

Comment [ma17]: The weight of the fish used in the experiment needs to be mentioned

Comment [ma18]: You needed to have also analysed the concentration of omega 3 fatty acids and lipid/fat in the feed before adding any fish oil to it. This could give a good baseline for making comparisons with the experimental fish

In addition, the concentration of omega 3 fatty acids in the fish oil also needed to be analysed

2.3 Observation Parameters

2.3.1 Specific Grow Rate (SGR)

The test fish were calculated using the following equation [46]:

$$SGR = \frac{\ln W_t - \ln W_o}{t} \times 100\%$$

Information:

W_t = Average weight of fish at the end of rearing (g)
W_o = Average weight of fish at the beginning of rearing (g)
t = Length of time of rearing

2.3.2 Feed Efficiency

The formula used to calculate feed efficiency is [29]:

$$FCR = \frac{F}{(W_t + D) - W_o}$$

Information:

EP = Feed efficiency (%)
W_t = Final fish weight (g)
W_o = Initial fish weight (g)
D = Dead fish weight (g)
F = Amount of feed consumed (g)

2.3.3 Proximate Analysis and Omega-3 Fatty Acids Proximate

Proximate analysis of pangasius catfish meat was carried out at the Laboratory of ruminant nutrition and animal feed chemistry, Faculty of Animal Science, Padjadjaran University. Omega-3 fatty acids were tested using the Gas Chromatography method at the Indo Genetech Laboratory.

Comment [ma19]: Flesh

Comment [ma20]: Provide a reference (s) for the methods that were followed in this study. The authors are also advised to explain the methods use for omega 3 fatty acid analysis.

3. RESULTS AND DISCUSSION

3.1 Specific Growth Rate

The biomass data for *Pangasius catfish* during the study is shown in Figure 1. Based on these data, each treatment showed a different average increase in biomass. The average initial biomass of *Pangasius catfish* was 178.48 g – 190.62 g and at the last observation, the biomass reached 294.17 g – 238.89 g. The highest increase in average growth value was in treatment C with an average final biomass of 252.87 g.

Comment [ma21]: I advise that the graph should have error bars





Fig 1. Diagram of Pangasius Catfish Biomass

The SGR yield of pangasius catfish was still relatively good at >0.72% [2] and the highest yield of 2.11% was comparable to the biomass value obtained during the study. This shows that the application of fish oil to commercial feed is still in a good range for the specific growth rate, where the results obtained are 1.69%-2.07%. The difference in average growth in each treatment was caused by the addition of fish oil which is fat into the feed, causing a larger source of energy to be used in activities such as movement or metabolism, while the energy source from protein can be optimized in the utilization of fish growth. The addition of fat content increases energy in the given feed and increases the growth rate of tilapia [38]. The addition of fish oil increased the growth and survival of tilapia [11]. The addition of fish oil to *Pangasius catfish* feed showed a positive impact on its growth by considering the limit

Comment [ma22]: The authors need to phrase these sentences better, linking them to the results of the study using *Pangasius*.



of fish oil content given.

Comment [ma23]: Remove the word value from the graph

Fig 2. Diagram of Pangasius Catfish Specific Growth Rate

Comment [ma24]: Be consistent with the words used. Pangas is appearing here only. Remember, in the introduction, you did not state anything about Pangas but you have been consistently using Pangasius

Fat that enters the body becomes one of the energy contributors to support metabolic activities so that most of the protein from feed can be utilized to support growth. However, the high-fat content due to the addition of fish oil causes the activity of lipogenic enzymes to decrease, thereby inhibiting fatty acid synthesis [48]. The high-fat content in the feed will increase the chance of fat peroxidase and affect the sensory attributes of the muscles [3]. This can result in lower growth rates and increased feed conversion. The low-fat content in feed results in less than optimal growth because the fat content in the feed given is only used for body maintenance and replacing damaged cells [22].

3.2 Feed Efficiency

Comment [ma25]: The introduction needs to state the known information on effects of fish oil on growth performance, and feed efficiency in farmed fish species. Then state what information gaps are there that this study is addressing

The addition of fish oil to the feed given to pangasius catfish showed an increase in protein, fat content, carbohydrates, and energy in fish at the beginning of the study and fish at the end of the study in treatments A and C, which are presented in Table 2. The increase in fat content of fish meat at the beginning of the study to treatment A at the end of the study reached 26% while compared to treatment C at the end of the study it reached 50.82%.




Fig. 3 Diagram of Feed Efficiency

Comment [ma26]: Remove the word value from the graph

The value of feed efficiency in each treatment during the study was classified as good (69.66%-85.85%), feed can be said to be good if the feed efficiency value is more than 50% or even close to 100% [5]. High feed efficiency indicates efficient use of feed so that only a small amount of protein is broken down by the body to meet energy needs and the rest is used for growth [27]. This supports the results of the specific growth rate of pangasius catfish during maintenance which is classified as good in each treatment.

The efficiency of feed utilization by pangasius catfish increased from every week of rearing in treatment C (4% fish oil), this showed that the adaptation of feed with the addition of fish oil was better by pangasius catfish during rearing. Treatment with the added fish oil more than 4% showed an increase and decrease in several weeks of rearing, which was directly proportional to the specific growth rate which was smaller than treatment C (4% fish oil). The difference in feed efficiency values is thought to be due to the digestibility of different feeds.

Comment [ma27]: The authors do not provide any data that they collected to support this statement

One of the indicators used to assess the level of efficiency of feed given to fish is feed digestibility [8].

UNDER PEER REVIEW

3.3 Proximate Analysis and Fatty Acid Omega-3

The following is the result of the proximate analysis of pangasius catfish at the beginning of the study, the treatment without the addition of fish oil, and the addition of 4% fish oil.

Table 1. Results of Proximate Analysis Pangasius Catfish

Perlakuan	Water %	Ash %	Protein %	Fat %	Carbohydrate %	Gross Energy Kcal/kg
Fish initial	81,90	3.37	18,59	6,10	66,41	3801
Fish without addition fish oil	44,05	5.19	21,13	7,71	67,79	4173
Fish with addition 4% fish oil	34,08	1.97	22,42	9,20	70.12	4333

Comment [ma28]: It would have been better if the authors did proximate analysis for each of the treatments in this study

Fat content in fish at the beginning of rearing, treatment A and treatment C of 6.10%, 7.71% and 9.20. Pangas catfish contain a fat content of 5.49% [50]. The fat content in fish is obtained from the consumption of available feed which is then absorbed or synthesized by the liver. The fat anabolism process goes through several stages, namely the formation of glycerol, the formation of fatty acid molecules and the condensation of fatty acids with glycerol to form fats [41]. Fat catabolism occurs by the breakdown of fats into fatty acids and glycerol. Fatty acids will go through the beta-oxidation process to produce acetyl CoA which enters the Krebs cycle and produces energy [47]. Through the process of fat metabolism, higher fat content which is one of the energy sources has the potential to increase the amount of energy in fish. In addition, the results of fatty acids derived from fat in the feed will produce omega-3 which are essential fatty acids in the body.

Pangas catfish contains a water content of 77.43% [34] which indicates that the water content of Pangas catfish in nature contains a high water content equal to the water content of the control treatment. The water content has the opposite relationship with the fat content, namely the higher the water content in the fish, the lower the fat content [44]. The entry of water into the spaces between cells and plasma. Fish with red meat has a low protein content, but a higher water content [17]. White fish meat has a high protein content and low water content. Pangas catfish have a flesh color that tends to be whiter in color compared to other fish that have a higher protein content.

The protein content of the catfish pangas increased from the beginning, the control, and treatment C at the end of the rearing with values of 18.59%, 21.13%, and 22.42%, respectively, when compared to the cultured catfish pangas containing 18.9% protein content while in nature it was 20.2% [18], the results of this study showed that the difference in numbers was not significant but there was an increase in treatment C (4% fish oil). The level of digestibility of each content in the feed source will affect the content of the fish. The level of protein and fat digestibility in fish will be directly proportional, if there is an increase in protein digestibility, there will be an increase in fat digestibility and vice versa [24]. This happens because the fatty acids contained in the fat are used to contribute fish metabolism which can affect protein digestibility. So that the addition of fish oil in feed containing fatty acids has an effect on protein levels in fish, feeding with fatty acid content in Kuwe fish

affects protein digestibility [33]. One of the functions of protein, namely as a source of energy, has been fully fulfilled through the existing fat.

Carbohydrates increased in the initial, control, and treatment C fish with values of 66.41%, 67.79%, and 70.12%, respectively, this increase indicates that the pangasius catfish can utilize nutrients in the feed. Carbohydrates and fats have a big role in meeting the needs of fish intake. The increase in carbohydrates was in line with the increase in fat in Pangasius catfish. Carbohydrates will be converted into glucose which is used for energy needs, some will be stored as glycogen in the liver and fish meat and some will be converted into fat and will be stored in fat tissue [1]. The ash content of Pangasius catfish meat during the study was still relatively low. Ash content is influenced by the high and low levels of minerals in fish meat, minerals play an important role in body functions, at every level of cells, tissues, organs, and even the function of the body as a whole [39].

Pangas catfish have an energy value of 2070 to 2210 Kcal/kg [16]. The increase in energy until the end of rearing indicates the utilization of energy from the feed given. Fat in the body is more likely to be stored in the body as an energy source that is utilized for growth [25]. The feed with fish oil added had a higher energy content than the feed without fish oil added. The higher energy will support the activity of the pangasius catfish, which will be utilized in the process of metabolism, movement, growth, and reproduction. Optimal growth for fish requires an energy of 3320 Kcal so growth and energy utilization in this study are classified as optimal [9].

The effect of adding fish oil to the feed on the levels of EPA and DHA in treatments A and C can be seen in Table 2.

Table 2. Results of analysis of fat, EPA, and DHA Content of Pangasius Catfish and Fish Oil Used

Treatment	EPA (mg ⁻¹ 100 gr)	DHA (mg ⁻¹ 100 gr)
Fish Oil Used	1698.5	11175.8
Fish without addition fish oil	7.3	24
Fish with addition 4% fish oil	10.6	30.7

The addition of fish oil showed an increase in EPA levels in treatment C reaching 27.91% and DHA reaching 31.13%. Pangasius catfish contains EPA 8.5-23.8 mg⁻¹ and DHA 2.55-11.05 mg⁻¹ 100 gr [12]. Catfish also experienced an increase in EPA 115.56% and DHA 114.47% with the addition of 4% fish oil to the feed. Feeding a feed with a higher omega-3 content will further increase the omega-3 content in fish, but if the feed is given a lower omega-3 content it will produce higher omega-9 and vice versa [47]. The addition of 5% fish oil to eel increased EPA 0.51 mg⁻¹ 100g and decreased omega-3 yield with feeding >5% fish oil [1]. The difference in the results obtained is thought to be due to differences in the type of fish, the fish oil used, and the time of maintenance carried out [26].

The process of fat metabolism including omega-3 fatty acids occurs in the liver, especially in the cytoplasm [47]. EPA and DHA anabolism begins with the breakdown of fat from feed into glycerol which will eventually produce palmitate CoA which will synthesize long and short-chain unsaturated fatty acids. The formation of EPA and DHA is assisted by elongase and desaturase enzymes [6]. However, omega-3 fatty acids are essential fatty acids that must be supplied from the given feed source [15]. The addition of fish oil with a high content of unsaturated fatty acids is the main source of ingredients for the formation of EPA and DHA in pangasius catfish.

Comment [ma29]: The authors needed to have analysed samples from each of the treatments separately

3.5 Water Quality

Water quality during the study for each treatment is presented in the table. 3. Water quality is one of the factors in the environment of fish living media that affects the survival and growth of fish. Measurement of water quality aims to see if some parameters are still within tolerance and support the growth of good pangasius catfish. The results of water quality measurements in each treatment were relatively the same, this shows that feeding with the addition of fish oil did not change the water quality of the rearing media.

Table 3. Results of Water Quality

Parameter	Unit	Range Value
Treatment	°C	23-30
pH	-	7.1-8.2
Dissolved Oxygen	mg/L	5.9-7.5
Ammonia	mg/L	0-0.3

Temperature water that is too high or too low can result in most of the energy stored in the fish's body being used for adaptation to an unfavorable environment so that it can damage the metabolic system or exchange of substances [19]. The temperature range of 25-32°C is the optimal temperature for supporting growth for tropical aquatic organisms [28]. In most fish species, the metabolic rate above the optimum temperature will increase and energy begins to be diverted from growth to compensate for the high metabolic rate so that the growth rate decreases [42].

The pH content during the study did not change significantly in each treatment and was still in the concentration range that supported the growth of catfish heat. The pH suitable for the life of the pangasius catfish ranges from 6.5 to 8.0 [32]. Freshwater pond fish have an acidic dead point of 4.0 and an alkaline dead point of 11.0 [45]. Dissolved oxygen content ranged from 6-7.4 mg/L, this content was relatively constant until the end of the study because each maintenance container was equipped with aeration equipment to supply oxygen. The standard dissolved oxygen content for the growth of pangasius catfish is DO 3 mg/L [23]. The dissolved oxygen content is also affected by temperature, the higher the temperature, the lower the oxygen content and vice versa [31].

Ammonia levels during the study were quite low due to regular siphoning and water changes. Fish excrete 80-90% of ammonia (N-inorganic) through the process of osmoregulation, while from feces and urine about 10-20% of total nitrogen [43]. Ammonia levels are toxic compounds that can harm health and inhibit fish growth [7]. The higher the oxygen concentration, pH, and water temperature, the higher the ammonia content. Ammonia accumulation in aquaculture media is one of the causes of decreased water quality which can result fail in of fish culture production.

4. CONCLUSION

Based on the results obtained, it can be concluded that the addition of fish oil in pangasius catfish feed can be done by adding as much as 4% to increase daily growth, feed utilization efficiency, omega-3 fat content, and optimal water quality.

REFERENCES

Comment [ma30]: Water quality was not one of the parameters that the authors set out to investigate. Note: The title and introduction of this manuscript do not state anything on water quality.

In this study, just like many other studies, water quality was monitored in order to keep it within optimal ranges for maximum growth and development of the fish. So that only the feed is the factor influencing the observed growth and feed efficiency.

What the authors need to include under this section is the recommended optimal ranges (temperature, oxygen, pH, ammonia) for growing Pangasius, and then compare them with what the authors recorded during the 28-day study.

Comment [ma31]: Probably this should be temperature

Formatted: Highlight

Formatted: Highlight

Comment [ma32]: Double check the references for conformity to the journal guidelines. Also ensure that what is listed here is in the text and what is in the text above is included in the reference list.

1. Almatsier S. Basic principles of nutrition. Main Library Gramedia. 2004.
2. Bokings L, Koniyo Y, Juliana. Growth and survival of Siamese catfish fry with artificial feed and silk worms. *Scientific Journal of Fisheries and Marine Affairs*. 2016;4(3): 81–88.
3. Bureau DP, Hua K, Harris AM. The effect of dietary lipid and long-chain n-3 PUFA levels on growth, energy utilization, carcass quality, and immune function of rainbow trout, *Oncorhynchus mykiss*. *Journal of the World Aquaculture Society*. 2005;29:1–21.
4. Cholik FR, Jauzi A, Poernomo P. Aquaculture: hope for the nation's future. Archipelago Fisheries Society and Freshwater Aquarium Park – TMII. 2005.
5. Craig S, Helfrich LA. 2002. Understanding Fish Nutrition, Feeds and Feeding. Cooperative Extension Service Publication. Virginia State University, USA.
6. Diana FM. OMEGA 3. *Journal of Public Health*. 2012;6(2):113–117.
7. Floyd RF, Watson C. Ammonia florida cooperative extension service. University of Florida. 2005.
8. Gunadi B, Febrianti R and Lamanto. Digestibility performance of sinking and floating feed for Dumbo catfish (*Clarias gariepinus*) with and without aeration. *Proceedings of the Aquaculture Technology Innovation Forum*. 2010;823-829.
9. Haetami, 2007. Consumption and feed efficiency of Jambal Siam fish fed with different protein energy levels. *Journal of Aquatics*. 2007;3(2):146-158.
10. Halver JE, Hardy RW. Fish nutrition. Academic Press, New York. 2003
11. Hasan U, Siswoyo BH, Manullang HM, Irwanmay. Effect of adding fish oil to artificial feed on growth and survival of tilapia (*Oreochromis niloticus*) fry. *Journal of Aquaculture Indonesia*. 2002;1(1):38–46.
12. Hashim RB, Jamil EF, Zulkipli FH, Daud JM. Fatty acid compositions of Silver Catfish, *Pangasius sp.* farmed in several rivers of Pahang, Malaysia. *Journal of OleoScience*. 2015;209(2):205–209.
13. Hidayaturrehman, Santoso, HB, Nurley. Profile of liver glycogen levels in hyperglycemic rats after administration of catfish (*Pangasius hypophthalmus*) fish oil extract. *Borneo Journal Pharmascientech*. 2017;01(02).
14. Innis SM. The Role of dietary n-6 and n-9 fatty acids in the developing brain. *Development Neuroscience*. 2008;22:21-28.
15. Izquierdo M. 2005. Essential fatty acid requirements in mediterranean fish species. *Cahiers Options Mediterranee*. 2005;63:91-102.
16. Jahan S, Habib AHS, Islam S, Begum M, Bardhan S. Comparative study on proximate and mineral composition of native and hybrid Pangas (*Pangasius pangasius*, *P. hypophthalmus*) at raw and fried stages. *Asiat Journal. soc. Bangladesh*. 2021;47(1):13–22.
17. Kantun W, Malik AA, Harianti. The feasibility of solid waste Tuna Loin Maddihang *Thunnus albacares* as raw material for diversified products. *Indonesian Fishery Product Processing Society*. 2015;18(3):303–314.
18. Karl MM, Lehmann I, Ostermeyer U, Schroder U. Natural chemical composition of commercial fish species: characterization of pangasius, wild and farmed turbot and barramundi. *Foods*. 2016;5(58):1–14.
19. Kelabora, Dominggas M. Effect of temperature on survival and growth of carp (*Cyprinus carpio*) larvae. *Terubuk Fisheries Periodic*. 2010;38(1):71-81.
20. Ministry of Marine Affairs and Fisheries. 2019 Performance Report
21. Klek S. Omega-3 fatty acids in modern parenteral nutrition: A review of the current evidence. *Journal of Clinical Medicine*, 2016;5(3):34.
22. Komariyah, Setiawan I. The effect of adding different doses of fish oil to artificial feeds on the growth of catfish (*Pangasius pangasius*) fry. 2009; 1(1):19-29.
23. Manunggal A, Hidayat R, Mahmudah S, Sudinno S, Kasmawijaya A. Water quality and growth of catfish using biopore technology on peatlands. *Journal of Fisheries and Marine counseling*. 2018;12(1):11-19.

24. Marzuqi M, Anjusary N. Digestibility of feed nutrients with different protein and fat levels in juvenile sand grouper (*Epinephelus corallicola*). *Journal of Tropical Marine Science and Technology*. 2013;5(2):311–324.
25. Meyer G, Fracossi DM. Protein requirement of judia fingerlings *Rhamida quelen* at two dietary energy concentrations. *Aquaculture*. 2004;19:74-78.
26. Mukti RC, Bambang N, Utomo P, Affandi R. Addition of fish oil to commercial feed on the growth of *Anguilla bicolor bicolor*. *Indonesian Journal of Aquaculture*. 2014;13(1):54–60.
27. Munisa Q, Subandiyono, Pinandoyo. Effect of different fat and energy content in feed on feed utilization and growth of catfish (*Pangasius pangasius*). *Journal of Aquaculture Management and Technology*. 2015;4(3):12–21.
28. Murtiono LH, Yuniarto D, Nuraini W. Analysis of land suitability for grouper cultivation with floating net cage systems with the application of geographic information systems in the waters of Ambon Dalam Bay. *Journal of Marine Aquaculture Technology*. 2016;6(1):1-16.
29. NRC. Nutrient Requirements of warm water fishes and shellfish. *Nutrional Academy of Sciences*. 1997;102.
30. National Research Council. 2011. Nutrient requirements of fish and shrimp. National Academy Press. 2011.
31. Nugroho A. Bioindicators of water quality. Jakarta, Trisakti University. 2006
32. Nurhamidah D. Effect of stocking density on growth performance of catfish (*Pangasius hypophthalmus*) with a recirculation system. Bogor Agricultural Institute. 2017
33. Most N, Rachmansyah, Usman. Effect of different fat sources on the growth of Kuwe fish, *Caranx sexfasciatus*. *Indonesian Research Journal*. 2002;8(3):25-29.
34. Rathod NB, Pagarkar AU, Pujari KH, Shingare PE, Satam SB, Phadke GG, Gaikwad BV. Status of valuable components from *Pangasius*: A review. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(4):2106-2120.
35. Rukmana HR, Yudirachman HH. The success of intensive catfish farming. Lily Publisher. 2016.
36. Sadekarpawar S, Parikh. P. Gonadosomatic and hepatosomatic indices of freshwater fish *Oreochromis mossambicus* in response to a plant nutrient. *World Journal of Zoology*. 2013;8(1):110-118.
37. Sari IP, Pontoh J. Chemical composition of fatty acids in anchovy meat. *Chem. Prog*, 2018;11(2), 2–7.
38. Setiawati M. Use of catfish fat in the diet of tilapia *Oreochromis niloticus*. Faculty of Fisheries and Marine Science. Bogor Agricultural Institute. Bogor. *Indonesian Journal of Aquaculture*. 2007; 6(1): 89-95.
39. RMST preparation. Making concentrate of catfish meat (*Pangasius hypophthalmus*) using ethanol solvent with different concentrates. Riau University. 2019;1-10.
40. Sipahutar RB. Morphoanatomy of catfish (*Pangasius* sp) reared by photoperiod manipulation. *Student Online Journal*. 2019;4(1), 1–11.
41. Siregar FA, Makmur T. Lipid metabolism in the body. *Journal of Public Health Innovation*, 2020;1(2).
42. Stickney RR. Principle of warm water aquaculture. John Wiley and Sons. 1979.
43. Sumoharjo. Nitrogen Waste Removal in Tilapia (*Oreochromis niloticus*) rearing in aquaponics system: bioreactor design configuration. Thesis. Postgraduate Program, Bogor Agricultural University. 2010.
44. Suzuki T. Fish and krill protein: processing technology. Applied Science. London : Publishers Ltd. 1991
45. Syahrizal S, Arifin MY. Analysis of mercury (Hg) content in Siamese catfish (*Pangasius hypophthalmus*) in the Lake Sipin KJA Jambi. *Journal of River and Lake Aquaculture*. 2017;2(1):9-17.
46. Takeuchi T. Laboratory work-chemical evaluation of dietary nutrients. In:Watanabe, T. (Ed.). fish nutrition and mariculture. JICA, Tokyo University Fish. 1988;179-229.

47. Tocher DR. Fatty Acids Requirements in Ontogeny of Marine and Freshwater Fish. *Aquaculture Research*. 2010;41:417-732.
48. Wang JT, Liu YJ, Tian LX, Mai KS, Du ZY, Wang Y, Yang HJ. Effect of dietary lipid level on growth performance, lipid deposition, hepatic lipogenesis in juvenile cobia *Rachycentron canadum*. *Aquaculture*. 2005;249:439-447.
49. Yandes Z, Affandi R, Mokiginta I. Effect of cellulose in feed on the biological condition of gouramy (*Osphronemus gourami* Lac) fry. *Indonesian Journal of Ichtiology*, 2003;3(1):27-33.
50. Zulkiflee SZ, Yusof MF, Rahman NAA, Rostam MA. Moisture, ash and fat composition of *Pangasianodon Hypophthalmus* (Sauvage, 1878) and *Pangasius Nasustus* (Bleeker, 1863). *International Journal of Allied Health Sciences*, 2020;4:1486-1495.

UNDER PEER REVIEW