

**Variation of hemogram parameters of refrigerated therapeutic whole blood at the blood transfusion center of the Douala General Hospital Cameroon. Why not consider;**

**“Evaluation of haemogram parameters in refrigerated therapeutic whole blood at the blood transfusion center of the Douala General Hospital Cameroon”**

**ABSTRACT**

A brief introduction of the study before the aim. Rewrite the aim of the study to capture the title of the study. The aim of this work was to evaluate the stability of hemogram parameters under the influence of cold chain on whole blood bags seronegative from infections transmissible by blood transfusion (ITT).

**Methodology:** A cohort of 200 bags of whole blood collected consecutively at the blood transfusion center of the Douala General Hospital constituted the population of the study conducted from March 1<sup>st</sup> to September 30<sup>th</sup>, 2018. Blood bag containing citrate phosphate adenine dextrose (preferably, citrate phosphate dextrose adenine) (CPDA) was collected and 5ml of this blood from each donor was collected directly into a dry tube (for the blood count using URIT 3000 Plus machine) and the rest was stored in a BIOBASE brand refrigerator at a temperature between 2 and 6°. (Statement is not clear. Is it the CPDA you are testing or the blood collected into the CPDA blood bag. Also is the 5ml of blood directly from the donor or from the donor's blood bag. On the day of delivery, a second sample was taken by section of the tubing of the bag for the analysis of a second hemogram of the same blood bags. Temperature, refrigerator opening frequency and blood were collected. While not use the word noted or documented in lieu of collected. Include the statistical package used and the level of significance set.

**Results:** The storage temperature significantly ( $p \leq 0.05$ ) decrease the rate of leukocytes, erythrocytes and hemoglobin. When the shelf life and the frequency of opening the refrigerator increased, hemoglobin, hematocrit, MCV and lymphocyte decreased significantly while leukocyte, MCHC, thrombocytes and granulocytes increase significantly.

**Conclusion:** How do you reconcile this result with the concluding statement? Temperature significantly ( $p \leq 0.05$ ) decrease the rate of leukocytes. While the shelf life and the frequency of opening the refrigerator increased leukocyte significantly

This study showed that, the decrease in leukocytes, red blood cells and hemoglobin levels was significantly related to either temperature, (please delete either) shelf life and frequency of opening the refrigerator. Knowledge on this variation could be very useful in the selection of blood donors HOW or do you mean to say selection of refrigerated blood or pint and the efficiency of transfusion.

**Keywords:** Cold Chain, Storage, Blood Transfusion, Variation.

## INTRODUCTION

Blood transfusion consists of administering blood or one of its components (red blood cells, platelets, granulocytes, plasma, proteins) from one or more subjects called (donors), to one or more sick subjects called (recipients) [1]. It is a substitute therapy which occupies a place of choice in the therapeutic arsenal of many countries in sub-Saharan Africa hence the existence of a cold chain for the conservation of blood. The latter is defined as a systematic process intended to ensure the safe storage and transport of blood from its collection from the donor until its administration to the patient who needs transfusion. According to WHO, hematological components are thermolabile and therefore retain their stability if the cold chain is maintained between 2 and 6°C for a period between 21 and 42 days depending on the type of anticoagulant [2]. However, during this storage process, several events may occur, such as voltage fluctuations, repeated opening of the refrigerator. These events associated with the long shelf life of more than 21 days, can be the cause of a number of variations in hematological parameters in the blood bag [3]. It is in view of these factors of variation that we proposed to evaluate the profile of hemogram parameters of blood bags after collection and at the time of delivery to the Douala General Hospital-Cameroon.

## MATERIALS AND METHODS

An evaluative study of 200 bags of whole blood was conducted in the blood transfusion service of the Douala General Hospital between the months of March 1<sup>st</sup> to September 30<sup>th</sup>, 2018. After collecting samples of blood bags (containing 63ml of CPDA anticoagulant to collect 450ml of blood) by frank venipuncture at the bend of the donor's elbow with asepsis containing 70° alcohols, a quantity of approximately 5ml was poured (used dispensed instead) into a dry tube for the first blood count and the tubing of each blood bag was cut after welding with an electric sealer, without breaking the safety of the blood product and stored in a

BIOBASE brand blood bank refrigerator at a temperature between 2 and 6°C. On the day of delivery, part of the tubing was cut and then its contents poured into a dry tube for a second blood count, using the same URIT 3000 plus hematological counter. Donor data (age, sex) were collected from the donor candidate selection form and the donor register. From the day of collection to the day of delivery of the blood bags, the temperatures and the frequency of opening (of before the refrigerators) the refrigerators were noted. The data collected during this study was entered using Microsoft Office Excel 2013 software and analyzed by SPSS 20.0 for Windows software. The data was presented in the form of frequency distribution tables. The comparison of frequencies and the tests of association were carried out using the Chi 2 test. The results were considered significant at  $p < 0.05$  NOT  $p \leq 0.05$  you need to be specific

## RESULTS

(Please this should be Table 1 with a caption of what its represent) Of the 200 donors selected, 89% were male with an average age of  $28.38 \pm 7.8$  years. The average storage time was  $7.32 \pm 6.79$  days with a minimum of 2 days and a maximum of 18 days. Regarding the average temperature, it was  $3.44 \pm 0.85^\circ\text{C}$  with a maximum of  $4.88^\circ\text{C}$  and a minimum of  $0.50^\circ\text{C}$ . The daily refrigerator opening frequency obtained was  $12 \pm 4$  times. (Please a table is needed for the result above as Table 1) (Delete “The other results are compiled in the following tables”.)

Table 2 not Table I presents the comparison of the means of the blood count parameters at collection and at delivery for the entire study population. According to this table, we noted a general variation of all the parameters of the hemogram between the day of collection of the blood bags (please be specific. Is it the blood bag or the pint of blood) and its delivery. A significant decrease in leukocytes, granulocytes and thrombocytes count was observed. However, a significant increase in lymphocytes was observed on the delivery blood count.

**Table 2 not Table I:** Comparison of means of blood count parameters at collection and at delivery

Parameters	COLLECTION		DELIVERANCE		VARIATIONS	
	Mean	Standard deviation	Mean	Standard deviation	Mean	Standard deviation
Leucocytes	4.16	1.257	3.043	1.38	1.1261	1.099
Erythrocytes	4.43	0.51	4.42	0.54335	0.0966	0.11729

<b>Hemoglobine</b>	11.911	1.51	11.9	1.5349	0.0879	0.15294
<b>Hematocrite</b>	36.15	4.57	36.453	5.0348	1.0432	1.41494
<b>MCV</b>	81.58	8.04	82.235	8.3653	1.0243	2.11685
<b>MCH</b>	27.171	5.2598	27.223	5.1676	0.4804	0.45623
<b>MCHC</b>	32.873	0.9171	32.556	1.1479	0.7965	1.00797
<b>Thrombocytes</b>	204.347	45.9037	117.265	54.4165	87.1558	46.38542
<b>Granulocytes</b>	42.076	9.8033	22.302	14.0259	20.9272	13.48464
<b>Lymphocytes</b>	46.077	10.1625	64.617	15.9613	20.0923	14.93952

**KEY:** MCHC : Mean Corpuscular of Hemoglobin Concentration; MCH: Mean Corpuscular of Hemoglobin Content; MCV: Mean Cell Volume (delete the of)

**Table 3 not Table II** summarizes the correlations between the variation of blood count parameters and the storage temperature. A negative correlation coefficient was observed in leucocyte ( $r = -0.232$ ,  $p = 0.001$ ), erythrocytes ( $r = -0.172$ ,  $p = 0.015$ ), and haemoglobin ( $r = -0.219$ ,  $p = 0.002$ ) and not a drop in leukocytes ( $p=0.001$ ), erythrocytes ( $p=0.015$ ) and hemoglobin ( $p=0.002$ ) was observed with the drop in temperature (a negative correlation).

**Table 3 not Table II:** correlation between the variation of blood count parameters and storage temperature

<b>PARAMETERS</b>	<b>R</b>	<b>P-VALUE</b>
<b>variation of leucocytes</b>	- 0.232	0.001
<b>variation of erythrocytes</b>	- 0.172	0.015
<b>variation of hemoglobin</b>	- 0.219	0.002
<b>variation of hematocrite</b>	- 0.028	0.693
<b>variation of MCV</b>	- 0.020	0.776
<b>variation of MCH</b>	- 0.050	0.479
<b>variation of MCHC</b>	- 0.027	0.710
<b>variation of thrombocytes</b>	- 0.057	0.426
<b>variation of granulocytes</b>	- 0.136	0.056
<b>variation of lymphocytes</b>	- 0.038	0.600

**KEYS:** MCHC : Mean Corpuscular of Hemoglobin Concentration; MCH: Mean Corpuscular of Hemoglobin Content; MCV: Mean Cell Volume; R: correlation factor

Table 4 not The tables below and it's a correlation table please present the variations of the means of complete blood count parameters with respect to the duration of storage (delete (table III) ) and write out the results Table 5 represent Delete and also it's a correlation table. the frequency of opening of the refrigerator (delete (table IV).) write out the results for each tables. Thanks. We noted from these tables a significant increased ( $p < 0.05$ ) not ( $p < 0.5$ ) in leukocytes, granulocytes, MCHC, thrombocytes and a significant decrease ( $p > 0.05$ ) not ( $p < 0.5$ ) in hemoglobin, hematocrit, MCV and lymphocytes (note; was observed) with increased in the number of days of storage and the frequency of opening the refrigerator.

Please rewrite the results (using r and p as a correlation table as corrected in table 2 above) for Tables 4 and 5 with the right tittles separately not combined

**Table 4 not Table III:** Correlation between the variation of blood count parameters and storage time

PARAMETERS	R	P-VALUE
variation of leucocytes	0.276	0.0001
variation of erythrocytes	- 0.121	0.089
variation of hemoglobin	- 0.326	0.0001
variation of hematocrite	- 0.568	0.0001
variation of MCV	- 0.803	0.0001
variation of MCH	0.041	0.565
variation du MCHC	0,547	0.0001
variation of thrombocytes	0.374	0.0001

<b>variation of granulocytes</b>	0.327	0.0001
<b>variation of lymphocytes</b>	- 0.282	0.0001

**KEYS:** MCHC : Mean Corpuscular of Hemoglobin Concentration; MCH: Mean Corpuscular of Hemoglobin Content; MCV: Mean Cell Volume; R: correlation factor

**Table 5 not Tableau IV** : correlation between the variation of hemogram parameters and the frequency of opening of the refrigerator.

<b>PARAMETERS</b>	<b>R</b>	<b>P-VALUE</b>
<b>variation of leucocytes</b>	0.254	0.0001
<b>variation of erythrocytes</b>	-0.147	0.046
<b>variation of hemoglobin</b>	-0.334	0.0001
<b>variation of hematocrit</b>	-0.595	0.0001
<b>variation of MCV</b>	-0.821	0.0001
<b>variation of MCH</b>	0.073	0.325
<b>variation of MCHC</b>	0.598	0.0001
<b>variation of thrombocytes</b>	0.334	0.0001
<b>variation of granulocytes</b>	0.298	0.0001
<b>variation of lymphocytes</b>	-0.280	0.0001

**KEYS:** MCHC: Mean Corpuscular of Hemoglobin Concentration; MCH: Mean Corpuscular of Hemoglobin Content; MCV: Mean Cell Volume; R: correlation factor

## DISCUSSION

This study aimed to evaluate the stability of blood count parameters during storage, (full stop after storage and then begin the next sentence with This study revealed) revealed a decrease in erythrocytes which was significantly correlated with the increase in the frequency of opening the refrigerator, the increasing number of days of storage and the drop in the storage temperature of the blood bag in the cold chain. Indeed, a red blood cell has a lifespan of about 120 days and can be eliminated by two mechanisms, namely extravascular hemolysis which is predominant and intravascular hemolysis [4, 5]. During storage, erythrocytes are subject to lesions resulting from the alteration of energy metabolism (decrease in ATP, acidosis, increase in calcium concentration) as well as oxidative stress (oxidative attacks on structural proteins) thus causing modifications to delete both metabolic, functional, morphological and

rheological erythrocytes. These lesions appear gradually from the first days of storage and accumulate rapidly from the second week, hence their decrease observed over time [6,7,8,9]. In addition, greater hemolysis is reflected with the long stay of erythrocytes in the cold chain by a decrease in the level of hemoglobin. These results are comparable to those of (10) in lieu of Jung-Min Lee and Jin Seok Kang (you need to consistent using figure in place of authors name) who observed a variation in hemoglobin in dogs delete this[10].

The number of thrombocytes decreases upon delivery. According to some studies, when standard platelet concentrates are prepared and well preserved, they can only last a maximum of five days (in vitro), a period which guarantees their viability for a satisfactory platelet transfusion yield [3]. However, delete an increased thrombocytes “was” in place of “is” observed when they are varied with respect to increased in the frequency of opening the refrigerator and the shelf life. This increase can be attributed to pseudothrombocytosis due to fragments of hemolyzed red blood cells, proteins, etc. which can be read as platelets by automaton [11]. Storage is accompanied by a decrease in antioxidant defenses and an increase in oxidative attacks with the formation of Reactive Oxygen Species (ROS) [6, 7-8]. Causing fragmentation of structural proteins, reduced deformability, osmotic fragility and increased mechanical fragility that can be read as platelets [12,13]

Leukocytes decreased on the day of delivery compared to the day of donation. This result could also be due to oxidative stress during storage. This result is in agreement with those of several authors [14,15]. Granulocytes are neutrophils, eosinophils and basophils. The Urit write out in capital letters 300Plus machine counts them and associates them with the monocytes. With the influence of storage, in the long run between 2-6 degrees Celsius, the probable morphological changes are read as lymphocytes by the automaton; hence the increase in the rate of lymphocytes at the detriment of granulocytes with the increase in the number of days of storage and the frequency of opening the refrigerator. Quote the corroborator(s)

## CONCLUSION

The results of this study show a decrease in leukocytes, granulocytes, thrombocytes and an increase in lymphocytes on the day of delivery was correlated with the increase in frequency of opening the refrigerator, number of days of storage and a significant variation of storage temperature of the blood bags in the cold chain.

## **ETHICAL CONSIDERATIONS**

For this study, we received a research certificate from the Dean of the Faculty of Medicine and Pharmaceutical Sciences of the University of Douala: the administrative authorization of the Director of the General Hospital of Douala and the ethical clearance issued by the institutional ethics committee for human health research of the University of Douala.

### **COMPETING INTERESTS DISCLAIMER:**

**Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.**

### **REFERENCES (should be organized in accordance with the journal rules)**

- [1] WHO. Burkina Faso, summary country profi/2le for HIV/AIDS treatment scaleup. Available at [www.who.int/countries/bfa/](http://www.who.int/countries/bfa/) (consulted on 14/02/2022).
- [2] World Health Organization. Blood Transfusion, Labile Blood Products. Available: <http://www.medix.free.fr>. copyright 2014©medix.
- [3] WHO. 2008 Blood cold chain: Guide for the selection and acquisition of equipment and accessories. Geneva, p6.7.
- [4] Hematocell.fr Hematology laboratory of the University Hospital of Angers. Erythropoiesis [Internet], available at <http://www.hematocell.fr/index.php/enseignement-de-lhematologieculaire/globules-rouges-et-leur-pathologie/20-erythropoiese> (consulted on 15/08/19).
- [5] DAllard D. (2013). Normal and pathological erythropoiesis, c-Kit internalization and morphology of the nucleolus. Doctoral thesis in hematology and oncology. Paris:Paris 5 University, 160 p.

- [6] Orlov D, Karkouti K. (2015).The pathophysiology and consequences of red blood cell storage. *Anaesthesia* ,70(1):29-37.
- [7] Epps DE, Knechtel TJ, Bacznyskj O.(1994). Tirilazad mesylate protects stored erythrocytes against osmotic fragility. *Chem Phys Lipids*;74(2):163-74.
- [8] Riebardis AG, Antonelou MH, Stamoulis KE.(2007). Progressive oxidation of cytoskeletal proteins and accumulation of denatured hemoglobin in stored red cells. *J Cell Mol Med*; 11(1):148-55.
- [9] Hess JR. (2010).Red cell changes during storage. *Transfus Apher Sci*;43(1):51-59
- [10] Jung-Min Lee, Jin Seok Kang. (2007). Changes of hematological references depends on storage period and temperature condition in rat and dog; 32 (4):p7
- [11] RFL. The pitfalls of automates platelet counting. Available at [WWW.labovialle.Com/index.php/](http://WWW.labovialle.Com/index.php/) achives /54 articles –paus (consulted on119 /01/2022)
- [12] Roussel C, Dussiot M, Marin M. (2017). Spherocytic shift of red blood cells during storage provides a quantitative whole cell-based marker of the storage lesion: spherocytic shift of RBCs during. *Transfusion* ;57(4):1007-1018.
- [13] D'Alessandro A, Kriebardis AG, Rinalducci S, et al. (2015).An update on red blood cell storage lesions, as gleaned through biochemistry and omics technologies: An omics update on RBC storage. *Transfusion*;55(1):205-219.
- [14] SillimanCC,ThurmanG,Ambruso DR.(1992). Stored blood components contain agents that prime the neutrophilie NADDPH oxidative through the plateactivating-factor receptor. *Vox sang*;63(2):133-136.
- [15] BehroozG, Azita A, Ali AP, Mohammadreza D, Alireza G. (2018).Comparative evaluation of biochemical and hematological parameters of pre-storage leukoreduction during RGC Storage. *Int J hematomol oncol cell Res*; 12:35-42.