

BIVARIATE COMPOUND EXPONENTIATED SURVIVAL FUNCTION OF THE LOMAX DISTRIBUTION: ESTIMATION AND PREDICTION

Abstract:

In this paper, bivariate compound exponentiated survival function of the Lomax distribution is constructed based on the technique considered by AL-Hussaini (2011). Some properties of the distribution are derived. Maximum likelihood estimation and prediction of the future observations are considered. Also, Bayesian estimation and prediction are studied under squared error loss function. The performance of the proposed bivariate distribution is examined using a simulation study. Finally, a real data set is analyzed under the proposed distribution to illustrate its flexibility for real-life application.

Keywords: Lomax ^d Distribution; Bivariate distributions; Compound exponentiated survival functions; Maximum likelihood estimators; Prediction; Bayes ^e Estimators; Monte Carlo simulations.

1. Introduction

Although bivariate extensions of univariate distributions are useful, it has not been applied in practice due to shortage of inferential procedures caused by numerical complexity. Moreover, generalization of univariate models is not straightforward in the sense that certain desirable properties may hold for more than one multivariate model.

One of the objectives of this paper is to construct a *bivariate compound exponentiated survival function of the Lomax* (BCESFLO) distribution; based on the technique considered by AL-Hussaini (2011) who constructed a class of multivariate distributions. It could be useful in studying reliability maintainability of complicated systems.

This paper consists of six sections. In Section 2, construction of BCESFLO distribution based on the technique considered by AL-Hussaini (2011), also some properties of the distribution are obtained. Maximum likelihood estimation and prediction are considered in Section 3. In Section 4, simulation study and an example data set are presented to illustrate the theoretical results derived for ML estimation and prediction. In Section 5, Bayesian estimation; for the unknown parameters, rf and hrf of BCESFLO distribution, is derived, also Bayesian prediction is considered. Numerical

illustration: simulation study and an example data set for the results of Bayesian estimation and prediction is introduced in Section 6.

2. Construction of a Compound Exponentiated Survival of the Lomax Distribution

Recently, in the statistical literature several methodologies of constructing bivariate and multivariate distributions based on marginal and conditional distributions have been proposed; see Arnold *et al.* (1999, 2001), Kotz *et al.* (2000), Sarabia and Gomez-Deniz (2008) and Balakrishnan and Lai (2009) among others.

Bivariate survival data arise when each study subject experiences two events. Examples include failure times of paired human organs; (e.g., kidneys, eyes, lungs, breasts and others), first and second occurrences of given disease. Bivariate survival data may consist of time to diagnosis or hospitalization and the time to eventual death from a fatal disease. The medical literature considered paired organs of an individual as a two-component system, which work under interdependency circumstances. In industrial applications these data types may come from system whose survival depends on the survival of two similar components. For an example, the breakdown times of dual generators in a power plant or failure times of twin engines in a 2-engine airplane are illustrations of bivariate survival data. In fact, there are many bivariate distributions [see Kotz *et al.* (2000)] that can be employed for the analysis of paired data.

Two bivariate Pareto distributions were suggested by Mardia (1962) which are called bivariate Pareto of the first kind and bivariate Pareto of the second kind. Arnold (1983) suggested a bivariate Pareto of the fourth kind and presented three methods to derive this model. Muliere and Scarini (1987) proposed a bivariate Pareto survival function which was characterized by Padamadan and Nair (1994) using the survival function of the marginal distributions. In this paper, a BCESFLO is introduced.

In this section, two cases of the construction of the CESFLO distribution (univariate and bivariate) are introduced.

2.1 Construction of the univariate compound exponentiated survival of the Lomax distribution

AL-Hussaini (2011) introduced a theorem to construct a class of distributions by compounding the exponentiated survival function (*sf*) with the gamma probability density function

(pdf). The resulting class includes all distributions with positive domain. Such domain could be the whole positive half of the real line or subset of it. Suppose that T has *Lomax* (LO) distribution with the following pdf and *cumulative distribution function* (cdf), respectively, as follows:

$$g(t, \alpha) = \alpha(1+t)^{-(\alpha+1)}, \quad t > 0, \alpha > 0, \quad (1)$$

and

$$G(t, \alpha) = 1 - (1+t)^{-\alpha}, \quad t > 0, \alpha > 0. \quad (2)$$

If $G(t)$ is an absolutely continuous cdf and β is a positive real number, then

$$F(t|\beta) = 1 - [\bar{G}(t)]^\beta = 1 - [(1+t)^{-\alpha}]^\beta, \quad (3)$$

where $F(t|\beta)$ is an absolutely continuous cdf, hence the sf can be written as

$$s(t|\beta) = [\bar{G}(t)]^\beta = [(1+t)^{-\alpha}]^\beta. \quad (4)$$

The corresponding pdf is given by

$$\begin{aligned} f(t|\beta) &= \frac{d}{dt} [1 - s(t|\beta)] = \beta [\bar{G}(t)]^{\beta-1} g(t) \\ &= \alpha\beta [(1+t)^{-\alpha}]^{\beta-1} (1+t)^{-(\alpha+1)}, \quad t > 0, (\alpha, \beta > 0), \end{aligned} \quad (5)$$

where $g(t)$ is the pdf corresponding to $G(t)$, and β is a positive random variable with pdf $\eta(\beta)$ which is given by

$$\eta(\beta) = \frac{b^a}{\Gamma(a)} \beta^{a-1} \exp(-b\beta), \quad \beta > 0, (a, b > 0), \quad (6)$$

and

$$f(t) = \int_0^\infty f(t|\beta) \eta(\beta) d\beta, \quad (7)$$

where $f(t|\beta)$ is given by (5),

hence

$$f(t) = \frac{a}{b} \lambda_G(t) \left[1 - \frac{1}{b} \ln \bar{G}(t) \right]^{-(a+1)}, \quad (8)$$

applying (8), where $\bar{G}(t) = (1+t)^{-\alpha} \Rightarrow g(t) = \alpha(1+t)^{-(\alpha+1)}$, and $\lambda_G(t) = \frac{g(t)}{\bar{G}(t)} = \frac{\alpha}{1+t}$, then the pdf of the CESFLO distribution is given, for $t > 0, \alpha > 0, \beta > 0$, by

$$f(t) = \frac{a}{b} \left[\frac{\alpha}{1+t} \right] \left[1 + \frac{\alpha}{b} \ln(1+t) \right]^{-(a+1)}, t > 0, \alpha, \beta > 0, \tag{9}$$

from (9) the cdf of the CESFLO distribution can be written as

$$F(t) = 1 - \left[1 + \frac{\alpha}{b} \ln(1+t) \right]^{-a}. \tag{10}$$

The hazard rate function (hrf) corresponding to $H(t)$ is

$$\begin{aligned} \lambda_H(t) &= \frac{f(t)}{1 - F(t)} \\ &= \frac{a}{b} \frac{\alpha}{(1+t)} \left[1 + \frac{\alpha}{b} \ln(1+t) \right]^{-1}. \end{aligned} \tag{11}$$

The reversed hazard rate function (rhrf) is given by

$$\lambda_H^*(t) = \frac{\frac{a}{b} \frac{\alpha}{(1+t)} \left[1 + \frac{\alpha}{b} \ln(1+t) \right]^{-(a+1)}}{1 - \left[1 + \frac{\alpha}{b} \ln(1+t) \right]^{-a}}.$$

2.2 Construction of the bivariate compound exponentiated survival of the Lomax distribution

Suppose that $\underline{t} = (t_1, t_2)$ is a random vector such that $F(t_i|\beta)$ and $f(t_i|\beta)$ are given by

$$F(t_i|\beta) = 1 - [\bar{G}(t)]^{\theta_i \beta} = 1 - [(1+t_i)^{-\alpha_i}]^{\theta_i \beta},$$

and

$$f(t_i|\beta) = \theta_i \beta \alpha_i [(1+t_i)^{-\alpha_i}]^{\theta_i \beta - 1} (1+t_i)^{-(\alpha_i + 1)}, \quad i = 1, 2, \tag{12}$$

for $i=1, 2$, respectively, where $t_i, \beta, \alpha_i, \theta_i > 0$, where β is a positive random variable following the gamma distribution with pdf given by (6).

The bivariate pdf of the random vector \underline{t} can be constructed by compounding $\varphi(\underline{t}; \beta)$ with $\eta(\beta)$ to yield

$$f(\underline{t}) = \int_{\theta}^{\infty} \varphi(\underline{t}; \beta) \eta(\beta) d\beta,$$

where

$$\varphi(\underline{t}; \beta) = \prod_{i=1}^2 f(t_i|\beta), \quad \text{and} \quad \underline{t} = (t_1, t_2).$$

Then the bivariate pdf of the random vector \underline{t} can be constructed by compounding $\varphi(\underline{t}; \beta)$ with $\eta(\beta)$, then the BCESFLO can be written as follows:

$$f(\underline{t}) = f(t_1, t_2) = \frac{\Gamma(a+2)}{\Gamma(a)} \left[\prod_{i=1}^2 \gamma_i \lambda_G(t_i) \right] \left[1 - \sum_{i=1}^2 \gamma_i \ln \bar{G}(t_i) \right]^{-(a+2)},$$

where $\gamma_i = \frac{\theta_i}{b}$ and $\lambda_G(t_i) = \frac{\alpha_i}{1+t_i}$, then

$$f(\underline{t}) = a(a+1) \left[\prod_{i=1}^2 \gamma_i \left(\frac{\alpha_i}{1+t_i} \right) \right] \left[1 + \sum_{i=1}^2 \alpha_i \gamma_i \ln(1+t_i) \right]^{-(a+2)}, t_i > 0, i=1,2 \tag{13}$$

Proof

Next, we will derive the function f. Observe that

$$f(\underline{t}) = \int_0^\infty \varphi(\underline{t}; \beta) \eta(\beta) d\beta, \tag{14}$$

where

$$\begin{aligned} \varphi(\underline{t}; \beta) &= \prod_{i=1}^2 f(t_i | \beta) = \prod_{i=1}^2 \theta_i [\bar{G}(t_i)]^{\theta_i \beta} \frac{g(t_i)}{\bar{G}(t_i)} \\ &= \beta^2 \prod_{i=1}^2 \theta_i \lambda_G(t_i) \exp \left[\sum_{i=1}^2 \theta_i \beta \ln(\bar{G}(t_i)) \right]. \end{aligned} \tag{7} \tag{15}$$

Substituting (15) and (6) in (14), one obtains

$$\begin{aligned} f(\underline{t}) &= \prod_{i=1}^2 \theta_i \lambda_G(t_i) \frac{b^a}{\Gamma(a)} \int_0^\infty \beta^{a+1} e^{-\beta [b - \sum_{i=1}^2 \theta_i \ln(\bar{G}(t_i))]} d\beta \\ &= \frac{\Gamma(a+2)}{\Gamma(a)} \left[\prod_{i=1}^2 \gamma_i \lambda_G(t_i) \right] \left[1 - \sum_{i=1}^2 \gamma_i \ln \bar{G}(t_i) \right]^{-(a+2)}. \end{aligned}$$


where $\gamma_i = \frac{\theta_i}{b}$, $\lambda_G(t_i) = \frac{g(t_i)}{\bar{G}(t_i)}$,

Then the pdf in the bivariate case can be rewritten as


$$\begin{aligned} f(t_1, t_2) &= a(a+1) \left[\left(\frac{\theta_1}{b} \left(\frac{\alpha_1}{1+t_1} \right) \right) \left(\frac{\theta_2}{b} \left(\frac{\alpha_2}{1+t_2} \right) \right) \right] \\ &\quad \times \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) + \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-(a+2)}, \end{aligned} \tag{8} \tag{16}$$

where, $t_i, \alpha_i, \theta_i > 0$, for $i=1,2$.


The contour plots of the joint pdf of the BCESFLO distribution for different parameter values are presented in Figure 1.




(1.a)



(1.b)



(1.c)



(1.d)

Figure 1. The contour plots of the joint pdf of the BCESFLO distribution for different parameter values

(1.a) ($a = 0.8, b = 0.8, \alpha_1 = 0.8, \alpha_2 = 0.8, \theta_1 = 0.8, \theta_2 = 0.8$),

(1.b) ($a = 1.5, b = 0.5, \alpha_1 = 2, \alpha_2 = 1, \theta_1 = 1, \theta_2 = 0.5$),

(1.c) ($a = 1, b = 1, \alpha_1 = 1, \alpha_2 = 1, \theta_1 = 1, \theta_2 = 0.5$)

and (1.d) ($a = 1, b = 1, \alpha_1 = 1, \alpha_2 = 1, \theta_1 = 1, \theta_2 = 1$).

The distribution function can be obtained from (13) as follows:

$$F(t) = \int_0^{t_1} \int_0^{t_2} \alpha(a+1) \left[\prod_{i=1}^2 \gamma_i \left(\frac{\alpha_i}{1+t_i} \right) \right] \left[1 + \sum_{i=1}^2 \alpha_i \gamma_i \ln(1+t_i) \right]^{-(a+2)} dt_2 dt_1.$$

On the other hand, the cdf of BCESFLO distribution is given by

In the arguments of the following functions add the corresponding parameters α_i, θ_i .

$$\begin{aligned}
 F(t_1, t_2) &= \int_0^{t_1} \int_0^{t_2} f(t_1, t_2) dt_2 dt_1 \\
 &= \int_0^{t_1} \int_0^{t_2} a(a+1) \left[\left(\frac{\theta_1}{b} \left(\frac{\alpha_1}{1+t_1} \right) \right) \left(\frac{\theta_2}{b} \left(\frac{\alpha_2}{1+t_2} \right) \right) \right] \\
 &\quad \times \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) + \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-(a+2)} dt_2 dt_1 \\
 &= 1 + \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) + \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-a} - \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) \right\} \right]^{-a} \\
 &\quad - \left[1 + \left\{ \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-a}, \tag{9} \tag{17}
 \end{aligned}$$

where, $t_i, \alpha_i, \theta_i > 0$, for $i=1,2$. Moreover, the marginal's -----

The marginal's pdf and cdf of the BCESFLO distribution can be written, respectively, as

$$f(t_i) = \frac{a\theta_i}{b} \left(\frac{\alpha_i}{1+t_i} \right) \left[1 + \frac{\theta_i \alpha_i}{b} \ln(1+t_i) \right]^{-(a+1)}, \quad i=1, 2,$$

and

$$F(t_i) = 1 - \left[1 + \frac{\theta_i \alpha_i}{b} \ln(1+t_i) \right]^{-a}, \quad i=1, 2, \tag{18}$$

where, $t_i, \alpha_i, \theta_i > 0$, for $i=1,2$. Besides, the joint -----

The joint reliability function (rf) of the BCESFLO distribution is given by:

$$\begin{aligned}
 R(t_1, t_2) &= p(T_1 > t_1, T_2 > t_2) = 1 - F(t_1) - F(t_2) + F(t_1, t_2) \\
 &= \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) + \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-a}, \quad i=1, 2. \tag{10} \tag{19}
 \end{aligned}$$


where, $t_i, \alpha_i, \theta_i > 0$, for $i=1,2$. Also, -----

Also, the joint hrf of the BCESFLO distribution can be defined as


$$\begin{aligned}
 h(t_1, t_2) &= \frac{f(t_1, t_2)}{R(t_1, t_2)} = a(a+1) \left[\left(\frac{\theta_1}{b} \left(\frac{\alpha_1}{1+t_1} \right) \right) \left(\frac{\theta_2}{b} \left(\frac{\alpha_2}{1+t_2} \right) \right) \right] \\
 &\quad \times \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1+t_1) + \frac{\theta_2 \alpha_2}{b} \ln(1+t_2) \right\} \right]^{-2}, \quad i=1, 2, \tag{11} \tag{20}
 \end{aligned}$$

where, $t_i, \alpha_i, \theta_i > 0$, for $i=1,2$.


If $(T_1, T_2) \sim$ BCESFLO, then for all values of $t_1 > 0, t_2 > 0$ both components of $h(t_1, t_2)$ are decreasing functions of t_1 and t_2 . The contour plots of the joint hrf of the BCESFLO distribution for different parameter values are presented in Figure 2.




(2.b)



(2.b)



(2.c)



(2.d)

Figure 2: The contour plots of the joint hazard of the BCESFLO distribution for different parameter values

(2.a) $(a = 1.1, b = 2, \alpha_1 = 1.5, \alpha_2 = 2.2, \theta_1 = 1.9, \theta_2 = 3.1)$,

(2.b) $(a = 1, b = 1, \alpha_1 = 1, \alpha_2 = 0.5, \theta_1 = 0.5, \theta_2 = 0.5)$,

(2.c) $(a = 0.8, b = 0.8, \alpha_1 = 0.8, \alpha_2 = 0.4, \theta_1 = 0.4, \theta_2 = 0.4)$

and (2.d) $(a = 1, b = 1, \alpha_1 = 1, \alpha_2 = 1, \theta_1 = 1, \theta_2 = 1)$.

3. Maximum Likelihood Estimation

In this section the ML estimation and prediction for the vector of the parameters

$\underline{\omega} = (a, b, \alpha_1, \alpha_2, \theta_1, \theta_2)$ for the BCESFLO distribution will be considered.

Throughout this section, highlight the sign of all the parameters

3.1 Maximum likelihood estimation of the parameters

(8)

The likelihood function of the BCESFLO distribution can be derived using the pdf in (16) directly, but compounding of $\prod_{i=1}^k f(t_i|\underline{\omega})$ and $\eta(\beta)$ can be applied to make the ML estimation easier, hence

In the arguments of the following functions add the corresponding parameters .

$$L(\underline{\omega}; t_1, t_2, \beta) = \prod_{j=1}^n f(t_{1j}, t_{2j}; \beta_j) = \prod_{j=1}^n \left[\prod_{i=1}^2 f(t_{ij}|\beta_j) \eta(\beta_j) \right]$$

$$= \frac{b^{an}}{[\Gamma(a)]^n} \left(\prod_{j=1}^n \beta_j \right)^{a+1} \theta_1^n \theta_2^n \alpha_1^n \alpha_2^n \prod_{j=1}^n (1+t_{1j})^{-1} \prod_{j=1}^n (1+t_{2j})^{-1}$$

$$\times \exp \left\{ - \sum_{j=1}^n \beta_j [\theta_1 \alpha_1 \ln(1+t_{1j}) + \theta_2 \alpha_2 \ln(1+t_{2j}) + b] \right\}, \quad (21)$$

where $\underline{\omega} = (a, b, \alpha_1, \alpha_2, \theta_1, \theta_2)$. The log likelihood function is given by

$$\ell(\underline{\omega}; t_1, t_2, \beta) = n a \ln b - n \ln \Gamma(a) + (a+1) \left(\sum_{j=1}^n \ln \beta_j \right) + n \ln \theta_1 + n \ln \theta_2 + n \ln \alpha_1$$

$$+ n \ln \alpha_2 - \sum_{j=1}^n \ln(1+t_{1j}) - \sum_{j=1}^n \ln(1+t_{2j})$$

Explain why the densities are Gamma

$$- \left\{ \sum_{j=1}^n \beta_j [\theta_1 \alpha_1 \ln(1+t_{1j}) + \theta_2 \alpha_2 \ln(1+t_{2j}) + b] \right\}, \quad (12) \quad (22)$$

The ML estimates of the parameters are obtained by differentiating (22) with respect to the parameters, setting to zero and then solving the resulting non-linear system of likelihood equations. Hence

$$\partial \ell / \partial a = n \ln \hat{b} - n \psi(\hat{a}) + \sum_{j=1}^n \ln \beta_j = 0, \quad \text{replace } \ln \text{ with } \ln(\cdot). \hat{b} \hat{a}, \text{ and } \psi ?$$

$$\partial \ell / \partial b = n \hat{a} / \hat{b} - \sum_{j=1}^n \beta_j = 0,$$

$$\partial \ell / \partial \alpha_1 = n / \hat{\alpha}_1 - \sum_{j=1}^n \beta_j \hat{\theta}_1 \ln(1+t_{1j}) = 0, \quad \hat{\alpha}_1 ? \hat{\theta}_1 ?$$

$$\partial \ell / \partial \theta_1 = n / \hat{\theta}_1 - \sum_{j=1}^n \beta_j \hat{\alpha}_1 \ln(1+t_{1j}) = 0,$$

$$\partial \ell / \partial \alpha_2 = n / \hat{\alpha}_2 - \sum_{j=1}^n \beta_j \hat{\theta}_2 \ln(1+t_{2j}) = 0, \quad \hat{\alpha}_2 ? \hat{\theta}_2 ?$$

and

$$\partial \ell / \partial \theta_2 = n / \hat{\theta}_2 - \sum_{j=1}^n \beta_j \hat{\alpha}_2 \ln(1 + t_{2j}) = 0,$$

where $\psi(\hat{a}) = \dot{\Gamma}(\hat{a}) / \Gamma(\hat{a})$. \grave{\Gamma}\{\Gamma\}?

The invariance property of the ML estimators can be applied to obtain the ML estimators for the $R(t_1, t_2)$ and $h(t_1, t_2)$ by replacing the parameters in (19) and (20) by their ML estimators as given below

$$\hat{R}(t_1, t_2) = \left[1 + \left\{ \frac{\hat{\theta}_1 \hat{\alpha}_1}{\hat{b}} \ln(1 + t_1) + \frac{\hat{\theta}_2 \hat{\alpha}_2}{\hat{b}} \ln(1 + t_2) \right\} \right]^{-\hat{a}}, \quad (23)$$

and

$$\hat{h}(t_1, t_2) = \hat{a}(\hat{a} + 1) \left[\left(\frac{\hat{\theta}_1}{\hat{b}} \left(\frac{\hat{\alpha}_1}{1 + t_1} \right) \right) \left(\frac{\hat{\theta}_2}{\hat{b}} \left(\frac{\hat{\alpha}_2}{1 + t_2} \right) \right) \right] \times \left[1 + \left\{ \frac{\hat{\theta}_1 \hat{\alpha}_1}{\hat{b}} \ln(1 + t_1) + \frac{\hat{\theta}_2 \hat{\alpha}_2}{\hat{b}} \ln(1 + t_2) \right\} \right]^{-2}. \quad (24)$$

Here the notation clashes. Now, t_1 and t_2 are not vectors. Replace t_1 by l_1 and t_2 by l_2 .

and hence the $\hat{R}(t_1, t_2)$ and $\hat{h}(t_1, t_2)$ can be calculated numerically.

3.2 Two-sample maximum likelihood prediction

Considering two- sample prediction, the two samples are assumed to be independent and drawn from the same distribution. In univariate case, the density of the s -th order statistic in the future sample is used to obtain the predictive pdf of the s -th ordered statistic. The first variable in the vector of bivariate distribution is the ordered observation and the second variable is its concomitants, therefore the joint pdf of the ordered observations and the concomitants is needed to obtain the joint predictive density function of future ordered observations and their concomitants.

For a future bivariate sample of size m , the joint pdf of future s -th ordered observation and its s -th concomitant denoted by $(y_{1(s:m)}, y_{2(s:m)})$, $s = 1, 2, \dots, m$, has the joint pdf which is given by (16) after replacing t_1 by $y_{1(s:m)}$ and t_2 by $y_{2(s:m)}$. For simplicity, it can be written as $(y_{1(s)}, y_{2(s)})$ instead of $(y_{1(s:m)}, y_{2(s:m)})$. Then the joint pdf of $(y_{1(s)}, y_{2(s)})$ can be derived as follows:

$$f_{s:m}(y_{1(s)}, y_{2(s)}; \underline{\omega}) = \frac{m!}{(s-1)!(m-s)!} f(y_{1(s)}, y_{2(s)}; \underline{\omega}) [F(y_{1(s)}, y_{2(s)})]^{s-1} [1 - F(y_{1(s)}, y_{2(s)})]^{m-s},$$

using binomial expansion to simplify the last term in the previous equation, one gets

$$[1 - F(y_{1(s)}, y_{2(s)})]^{m-s} = \sum_{j=0}^{m-s} \binom{m-s}{j} (-1)^j [F(y_{1(s)}, y_{2(s)})]^j$$

Thus, the joint probability density of $(y_{1(s)}, y_{2(s)})$ is

$$f_{s:m}(y_{1(s)}, y_{2(s)}; \underline{\omega}) = f(y_{1(s)}, y_{2(s)}; \underline{\omega}) \sum_{j=0}^{m-s} C_{m,s,j} [F(y_{1(s)}, y_{2(s)})]^{s+j-1}, \quad (25)$$

where $\underline{\omega} = (\dots)$, and $y_{1(s)}, y_{2(s)}, a, b, \dots, \theta_2 > 0$, with

$$C_{m,s,j} = \frac{m!}{(s-1)!(m-s-j)!j!} (-1)^j. \quad (26)$$

Substituting $f(t_1, t_2)$ given in (16) and $F(t_1, t_2)$ in (17) after replacing t_1 by $y_{1(s)}$ and t_2 by $y_{2(s)}$ then, the joint ML predictive pdf of the ordered observations and their concomitants is given by

$$\begin{aligned} f_{s:m}(y_{1(s)}, y_{2(s)}; \underline{\hat{\omega}}_{ML}) = & \hat{a}(\hat{a} + 1) \left[\left(\frac{\hat{\theta}_1}{\hat{b}} \left(\frac{\hat{\alpha}_1}{1 + y_{1(s)}} \right) \right) \left(\frac{\hat{\theta}_2}{\hat{b}} \left(\frac{\hat{\alpha}_2}{1 + y_{2(s)}} \right) \right) \right] \\ & \times \left[1 + \left\{ \frac{\hat{\theta}_1 \hat{\alpha}_1}{\hat{b}} \ln(1 + y_{1(s)}) + \frac{\hat{\theta}_2 \hat{\alpha}_2}{\hat{b}} \ln(1 + y_{2(s)}) \right\} \right]^{-(\hat{a}+2)} \\ & \times \sum_{j=0}^{m-s} C_{m,s,j} \left\{ 1 + \left[1 + \left\{ \frac{\hat{\theta}_1 \hat{\alpha}_1}{\hat{b}} \ln(1 + y_{1(s)}) + \frac{\hat{\theta}_2 \hat{\alpha}_2}{\hat{b}} \ln(1 + y_{2(s)}) \right\} \right]^{-\hat{a}} \right. \\ & \left. - \left[1 + \left\{ \frac{\hat{\theta}_1 \hat{\alpha}_1}{\hat{b}} \ln(1 + y_{1(s)}) \right\} \right]^{-\hat{a}} - \left[1 + \left\{ \frac{\hat{\theta}_2 \hat{\alpha}_2}{\hat{b}} \ln(1 + y_{2(s)}) \right\} \right]^{-\hat{a}} \right\}^{s+j-1}, \end{aligned} \quad (27)$$

$(y_{1(s)}, y_{2(s)}) > 0, \underline{\omega} > 0.$

where $y_{1(s)}, y_{2(s)}, a, b, \dots, \theta_2 > 0$

The point predictors of the future ordered observations and their concomitants $(Y_{1(s)}, Y_{2(s)})$, $s = 1, 2, \dots, m$, can be obtained as follows:

$$\hat{Y}_1 = E(y_{1(s)}; \underline{\hat{\omega}}_{ML}) = \int_{y_{1(s)}=0}^{\infty} y_{1(s)} \int_{y_{2(s)}=0}^{\infty} f(y_{1(s)}, y_{2(s)}; \underline{\hat{\omega}}_{ML}) dy_{2(s)} dy_{1(s)}, \quad (13) \quad (28)$$

and

$\hat{\underline{\omega}}_{ML}$?

$$\hat{Y}_2 = E(y_{2(s)}; \underline{\hat{\omega}}_{ML}) = \int_{y_{2(s)}=0}^{\infty} y_{2(s)} \int_{y_{1(s)}=0}^{\infty} f(y_{1(s)}, y_{2(s)}; \underline{\hat{\omega}}_{ML}) dy_{1(s)} dy_{2(s)}, \quad (14) \quad (29)$$

From (28) and (29), the point predictors Y_1 and Y_2 cannot be obtained in closed form. Then, the joint point predictors of the future ordered observations is

$$\hat{Y}_1, \hat{Y}_2 = E(y_{1(s)}, y_{2(s)}; \underline{\hat{\omega}}_{ML}) = \int_0^{\infty} \int_0^{\infty} y_{1(s)} y_{2(s)} f(y_{1(s)}, y_{2(s)}; \underline{\hat{\omega}}_{ML}) dy_{2(s)} dy_{1(s)}, \quad (30)$$

which can be evaluated numerically.

$\hat{Y}_1, \hat{Y}_2 = ?$

4. Numerical Illustration

This section aims to clarify the theoretical results for both estimation and prediction on the basis of simulated and real data set.

4.1 Simulation study

In this subsection, a simulation study is conducted to illustrate the performance of the presented ML estimates on the basis of the generated data from the BCESFLO distribution. The ML averages of the estimates of the parameters, rf and hrf are computed. Moreover, *confidence intervals* (CIs) of the parameters, rf and hrf are calculated. Simulation studies are performed using Mathematica 11 for illustrating the obtained results.

The steps of the simulation procedure are as follows:

- a) For given values of $\underline{\omega}$ (where $\underline{\omega} = (a, b, \alpha_1, \alpha_2, \theta_1, \theta_2)$), random samples of size n are generated from the BCESFLO distribution.
- b) For each sample size sort t_{ij} s, such that $(t_{11}, t_{21}), (t_{12}, t_{22}), \dots, (t_{1n}, t_{2n})$.
- c) Repeat the previous two steps N times, where N represents a fixed number of simulated samples.
- For the number of the population parameter values the Newton-Raphson method can be used, the ML averages and the CIs of the parameters are obtained. Also, the rf, hrf and their CIs are calculated using the ML averages of the parameters.
- Evaluating the performance of the estimates is considered through some measurements of accuracy. To study the precision and variation of the estimates, it is convenient to use the **estimated risk (ER)**

$$\text{estimated risk (ER)} = \frac{\sum_{i=1}^N (\text{estimator} - \text{true value})^2}{N}$$

- Simulation results of the ML estimates are displayed in Tables 1, 2, where N = 10000 is the number of repetitions and samples of size ($n=30, 50, 100$), and the population parameter values are $(a = 1.1, b = 2, \alpha_1 = 1.5, \alpha_2 = 2.2, \theta_1 = 1.9, \theta_2 = 3.1)$, and $(a = 0.6, b = 1.2, \alpha_1 = 0.79, \alpha_2 = 1.1, \theta_1 = 0.95, \theta_2 = 1.55)$.
- Tables 1 and 2 present the ML averages, ERs, and CIs of the unknown parameters based. While Tables 3 and 4 display the ML averages, ERs and CIs of the rf and hrf for different values of time t_{01}, t_{02} . The ML two-sample predictors are presented in Table 8.

4.2 Example data set

In this example, a data set is analyzed from a Sankaran-Nair bivariate Pareto distribution [see Sankaran-Nair (1994) and Sankaran and Kundu (2014)]. The generated data set for $n=30$ is: (0.252, 8.400), (1.105, 0.458), (0.427, 1.602), (12.491, 2.383), (0.260, 0.106), (0.240, 1.769), (4.888, 0.758), (0.870, 0.572), (0.036, 0.254), (1.537, 0.023), (1.508, 0.535), (0.239, 1.4120), (0.173, 0.011), (1.090,

1.278), (6.002, 0.017), (0.897, 2.032), (0.690, 0.138), (1.883, 0.398), (0.960, 0.257), (0.561, 0.573), (5.370, 0.325), (0.167, 0.260), (13.602, 0.364), (3.922, 0.938), (0.132, 0.547), (0.603, 0.102), (0.226, 0.481), (0.143, 0.779), (0.643, 0.071), (0.349, 1.586).

The Kolmogorov–Smirnov goodness of fit test is applied to check the validity of the fitted model. The p values are given, respectively 0.808 and 0.393. The p values showed that the model fits the data very well. Table 5 displays the ML estimates and *standard errors* (Se) of the unknown parameters for the real data set. While Tables 6 and 7 present the ML estimates, Se and CIs of the rf and hrf for different values of time t_{01}, t_{02} . Table 8 gives the ML two-sample predictors for the future observation.

Table 1
ML averages, variance, estimated risks and 95% confidence intervals of the parameters
($N = 10000, a = 1.1, b = 2, \alpha_1 = 1.5, \alpha_2 = 2.2, \theta_1 = 1.9, \theta_2 = 3.1$)

n	Parameters	Averages	Var	ER	UL	LL	Length
30	a	0.9764	0.0010	0.0163	1.0393	0.9134	0.1259
	b	1.7883	0.0030	0.0478	1.8963	1.6804	0.2159
	α_1	1.5112	9.6547e-06	0.0001	1.5173	1.5051	0.0122
	α_2	2.2207	0.00003	0.0005	2.2319	2.2095	0.0224
	θ_1	1.9142	0.00002	0.0002	1.9219	1.9065	0.0154
	θ_2	3.1292	0.00007	0.0009	3.1450	3.1134	0.0315
50	a	0.9951	0.00002	0.0110	1.0040	0.9861	0.0179
	b	1.8199	0.00007	0.0325	1.8359	1.8038	0.0321
	α_1	1.5086	1.4879e-08	0.0001	1.5089	1.5084	0.0005
	α_2	2.2168	9.5819e-08	0.0003	2.2174	2.2162	0.0012
	θ_1	1.9109	2.3872e-08	0.0001	1.9113	1.9106	0.0006
	θ_2	3.1237	1.9025e-07	0.0006	3.1245	3.1228	0.0017
100	a	0.9969	1.7630e-06	0.0106	0.9996	0.9944	0.0052
	b	1.8235	5.2811e-06	0.0312	1.8279	1.8189	0.0090
	α_1	1.5089	1.4657e-9	0.0001	1.5089	1.5088	0.0002
	α_2	2.2167	1.0896e-08	0.0003	2.2169	2.2166	0.0004
	θ_1	1.9112	2.3517e-09	0.0001	1.9113	1.9111	0.0002
	θ_2	3.1236	2.1634e-08	0.0005	3.1239	3.1233	0.0006

Table 2
ML averages, variance, estimated risks and 95% confidence intervals
of the parameters
($N = 10000, a = 0.6, b = 1.2, \alpha_1 = 0.79, \alpha_2 = 1.1, \theta_1 = 0.95, \theta_2 = 1.55$)

n	Parameters	Averages	Var	ER	UL	LL	Length
30	a	0.9567	0.0005	0.1277	1.0020	0.9113	0.0908
	b	1.7614	0.0018	0.3169	1.5254	1.6791	0.1647
	α_1	1.5235	9.5511e-07	0.5983	1.5254	1.5216	0.0038
	α_2	2.2370	7.6043e-06	1.29277	2.2424	2.2316	0.0108
	θ_1	1.9298	1.5324e-06	0.9599	1.9322	1.9273	0.0049
	θ_2	3.1521	0.00002	2.5668	3.1598	3.1445	0.0152
50	a	0.9425	1.2119e-06	0.1173	0.9447	0.9404	0.0043
	b	1.7359	3.3451e-06	0.2872	1.7395	1.7323	0.0072
	α_1	1.5232	9.6456e-07	0.5978	1.5251	1.5212	0.0038
	α_2	2.2376	3.2349e-08	1.2941	2.2379	2.2372	0.0007
	θ_1	1.9293	1.5476e-06	0.9591	1.9318	1.9269	0.0049
	θ_2	3.1529	6.4230e-08	2.5694	3.1534	3.1525	0.0009
100	a	0.9411	5.4548e-07	0.1164	0.9426	0.9397	0.0029
	b	1.7335	1.5400e-06	0.2846	1.7359	1.7311	0.0049
	α_1	1.5238	4.0737e-07	0.5988	1.5251	1.5226	0.0025
	α_2	2.2382	1.0435e-09	1.2955	2.2383	2.2381	0.0001
	θ_1	1.9302	6.5359e-07	0.9608	1.9317	1.9286	0.0032
	θ_2	3.1538	2.0718e-09	2.5723	3.1539	3.1537	0.0002

Table 3
ML averages, relative absolute biases, variance, estimated risks and 95% confidence
intervals of the reliability and hazard rate functions
($N = 10000, a = 1.1, b = 2, \alpha_1 = 1.5, \alpha_2 = 2.2, \theta_1 = 1.9, \theta_2 = 3.1, t_{01} = 2, t_{02} = 3$)

n	rf and hrf	Averages	RAB	Var	ER	UL	LL	Length
30	$R(t_{01}, t_{02})$	0.0756	0.2194	0.00002	0.0001	0.0831	0.0681	0.0150
	$h(t_{01}, t_{02})$	0.0051	0.1669	5.7242e-08	0.0016	0.0055	0.0046	0.0009
50	$R(t_{01}, t_{02})$	0.0761	0.2178	0.00001	0.0001	0.0831	0.0691	0.0140
	$h(t_{01}, t_{02})$	0.0051	0.1666	5.2811e-08	0.0016	0.0055	0.0046	0.0009
100	$R(t_{01}, t_{02})$	0.0759	0.2160	0.0000	0.0001	0.0826	0.0688	0.0138
	$h(t_{01}, t_{02})$	0.0050	0.1654	5.0551e-08	0.0015	0.0055	0.0046	0.0009

Table 4
ML averages, relative absolute biases, variance, estimated risks and 95% confidence intervals of the reliability and hazard rate functions
 ($N = 10000, a = 0.6, b = 1.2, \alpha_1 = 0.75, \alpha_2 = 1.1, \theta_1 = 0.95, \theta_2 = 1.55, t_{01} = 2, t_{02} = 4$)

n	rf and hrf	Averages	RAB	Var	ER	UL	LL	Length
30	$R(t_{01}, t_{02})$	0.0524	0.8323	1.2496e-08	0.0226	0.0527	0.0522	0.0004
	$h(t_{01}, t_{02})$	0.0014	0.1493	5.0258e-10	0.0009	0.0016	0.0013	0.0003
50	$R(t_{01}, t_{02})$	0.0522	0.8321	1.8122e-09	0.0224	0.0525	0.0523	0.0002
	$h(t_{01}, t_{02})$	0.0013	0.1386	1.8525e-10	0.0006	0.0015	0.0012	0.0003
100	$R(t_{01}, t_{02})$	0.0520	0.8320	1.7635e-09	0.0221	0.0521	0.0519	0.0002
	$h(t_{01}, t_{02})$	0.0011	0.1382	2.5761e-12	0.0002	0.0012	0.0010	0.0002

Table 5
ML estimates and standard errors of the parameters for the real data set

Parameters	Estimates	Se
a	0.9952	0.0109
b	1.8246	0.0307
α_1	1.5123	0.0002
α_2	2.2275	0.0008
θ_1	1.9156	0.0002
θ_2	3.1387	0.0015

Table 6
ML estimates and standard errors of the reliability and hazard rate functions for the real data set

rf and hrf	Estimates	Se
$R(t_{01}, t_{02})$	0.0727	0.0001
$h(t_{01}, t_{02})$	0.0052	0.0044

Table 7
ML estimates and standard errors of the reliability and hazard rate functions for the real data set

rf and hrf	Estimates	Se
$R(t_{01}, t_{02})$	0.0654	0.0001
$h(t_{01}, t_{02})$	0.0034	0.0038

Table 8
ML predictors and bounds of the future observation
under two-sample prediction
(n = 30, a = 0.3, b = 0.5, $\alpha_1 = 0.7, \alpha_2 = 2, \theta_1 = 1.9, \theta_2 = 3.1$)

s	$\hat{Y}_{(s)}$	Estimates	UL	LL	Length
7	$\hat{Y}_{1(s)}$	0.0068	0.0448	0.0000	0.0448
	$\hat{Y}_{2(s)}$	0.1141	0.3374	0.0147	0.3227
15	$\hat{Y}_{1(s)}$	0.0295	0.0452	0.0001	0.0451
	$\hat{Y}_{2(s)}$	0.2062	0.3563	0.0113	0.3450
18	$\hat{Y}_{1(s)}$	0.4845	2.1960	0.0222	2.1738
	$\hat{Y}_{2(s)}$	0.6831	1.3169	0.2638	1.0532

4.3 Concluding remarks

1. It is noticed, from Tables 1 and 2 that the ML averages are very close to the population parameter values as the sample size increases. Also, ER is decreasing when the sample size is increasing. This is indicative of the fact that the estimates are consistent and approaches the true parameter values as the sample size increases.
2. The lengths of the CIs of the parameters become narrower as the sample size increases.
3. The ML averages for the rf and hrf perform better as the sample size increases. Also, ER is decreasing when the sample size is increasing.
4. The length of the CI for the first future order statistic is smaller than the length of the CI for the last future order statistic [Tables 8 and 9].
5. The ML interval includes the estimates (between the LL and UL).

5. Bayesian Method

In this section Bayesian estimation and prediction for the vector of parameters $\omega = (a, b, \alpha_1, \alpha_2, \theta_1, \theta_2)$ for the BCESFLO distribution will be studied.
of

5.1 Bayesian estimation

AL-Hussaini and Ateya (2005) estimated the parameters under a *squared error loss* (SEL) function using Tierney- Kadane’s (1986) approximation form. Iliopoulos *et al.* (2005) considered bivariate gamma distribution for estimating the unknown parameters based on SEL function. Chadi *et al.* (2013) estimated the parameters and the mean time between failures of a bivariate exponential model under various loss functions, namely SEL, absolute error, DeGroot, LINEX

Throughout this section, highlight the sign of all the parameters

and Entropy loss functions. Lin *et al.* (2013) obtained the estimators for the parameters of Moran-Downton bivariate exponential distribution based on complete and Type-II censoring. Independent gamma priors were assumed for scale parameters and beta distribution for correlation parameter. Pradhan and Kundu (2015) derived the estimators for the parameters of the Block and Basu bivariate Weibull distribution. **Highlight which criterion you will apply for parameter estimates.**

If (a, b) , (α_1, θ_1) and (α_2, θ_2) are independent, a prior density function of **why do you say that are independent, if are parameter vectors?** $\underline{\omega} = (a, b, \alpha_1, \theta_1, \alpha_2, \theta_2)$ is given by

$$\pi(\underline{\omega}) \propto \pi_1(a, b)\pi_2(\alpha_1, \theta_1)\pi_3(\alpha_2, \theta_2), \tag{31}$$

where the first prior is **Why does it state that the expressions are proportional?**

$$\pi_1(a, b) = \pi_{11}(a|b)\pi_{12}(b), \quad \pi_{11} \sim \text{Gamma}(c_1, b) \text{ and } \pi_{12} \sim \text{Gamma}(c_2, c_3),$$

Explain why the densities are Gamma

the second prior is

$$\pi_2(\alpha_1, \theta_1) = \pi_{21}(\alpha_1|\theta_1)\pi_{22}(\theta_1), \quad \pi_{21} \sim \text{Gamma}(c_4, \theta_1) \text{ and } \pi_{22} \sim \text{Gamma}(c_5, c_6),$$

Explain why the densities are Gamma

and the third prior

$$\pi_3(\alpha_2, \theta_2) = \pi_{31}(\alpha_2|\theta_2)\pi_{32}(\theta_2), \quad \pi_{31} \sim \text{Gamma}(c_7, \theta_2) \text{ and } \pi_{32} \sim \text{Gamma}(c_8, c_9).$$

The three priors can be written as

$$\pi_1(a, b) \propto b^{c_1+c_2-1} a^{c_1-1} e^{-b(c_3+a)}, \tag{32}$$

$$\pi_2(\alpha_1, \theta_1) \propto \theta_1^{c_4+c_5-1} \alpha_1^{c_4-1} e^{-\theta_1(c_6+\alpha_1)}, \tag{33}$$

and

$$\pi_3(\alpha_2, \theta_2) \propto \theta_2^{c_7+c_8-1} \alpha_2^{c_7-1} e^{-\theta_2(c_9+\alpha_2)}. \tag{34}$$

Substituting from (32) and (33) in (31) and using the likelihood function in (21), then the posterior density function will separate into three posteriors, which are

$$\pi_1^*(a, b|t_1, t_2, \beta) \propto \frac{b^{na+c_1+c_2-1}}{(\Gamma(a))^n} \left(\prod_{j=1}^n \beta_j \right)^{a+1} a^{c_1-1} \exp \left[-b \left(\sum_{j=1}^n \beta_j + c_3 + a \right) \right], \tag{35}$$

$$\pi_2^*(\alpha_1, \theta_1|t_1, t_2, \beta) \propto \theta_1^{c_4+c_5+n-1} \alpha_1^{c_4+n-1} \prod_{j=1}^n (1+t_{1j})^{-1} \exp \left[-\theta_1 \sum_{j=1}^n \alpha_1 \beta_j \ln(1+t_{1j}) + c_6 + \alpha_1 \right], \tag{36}$$

and

Bayes' Theorem for probability distributions is often stated as:

$$\text{Posterior} \propto \text{Likelihood} \times \text{Prior}.$$

Now, substituting from (32), (33) and (34) in (31) and using the likelihood function in (21), we have that the posterior density function will separate into three posteriors, which are

Update the numbering of the formulas.

$$h(y_{1(s)}, y_{2(s)} | \underline{\omega}) = \int_{\underline{\omega}} I_1 I_2 I_3 I_4 I_5 d\underline{\omega}, \tag{43}$$

where

$$I_1 = \frac{(a+1)a^{c_1}}{(\Gamma(a))^n} b^{na+c_1+c_2-3} \theta_1^{c_4+c_5+n} \alpha_1^{c_4+n} \theta_2^{c_7+c_8+n} \alpha_2^{c_7+n},$$

$$I_2 = e^{-[b(\sum_{j=1}^n \beta_j + c_3 + a) + \theta_1(c_6 + \alpha_1) + \theta_2(c_9 + \alpha_2)]} (1 + y_{1(s)})^{-1} (1 + y_{2(s)})^{-1},$$

$$I_3 = \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1 + y_{1(s)}) + \frac{\theta_2 \alpha_2}{b} \ln(1 + y_{2(s)}) \right\} \right]^{-(a+2)},$$

$$I_4 = \left(\prod_{j=1}^n (1 + y_{1j}) \right)^{-1} \left(\prod_{j=1}^n (1 + y_{2j}) \right)^{-1} \left(\prod_{j=1}^n \beta_j \right)^{a+1} e^{-[\theta_1 \alpha_1 \sum_{j=1}^n \beta_j y_{1j}^{-\alpha_1} + \theta_2 \alpha_2 \sum_{j=1}^n \beta_j y_{2j}^{\alpha_2}]}$$

and

$$I_5 = \sum_{j=0}^{m-s} C_{m,s,j} \left\{ 1 + \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1 + y_{1(s)}) + \frac{\theta_2 \alpha_2}{b} \ln(1 + y_{2(s)}) \right\} \right]^{-a} - \left[1 + \left\{ \frac{\theta_1 \alpha_1}{b} \ln(1 + y_{1(s)}) \right\} \right]^{-a} - \left[1 + \left\{ \frac{\theta_2 \alpha_2}{b} \ln(1 + y_{2(s)}) \right\} \right]^{-a} \right\}^{s+j-1}. \tag{44}$$

identify the sign of the parameters

The point predictors of the future ordered observation and their concomitants $(Y_{1(s)}, Y_{2(s)})$,

$s = 1, 2, \dots, m$, under SEL function can be obtained as follows:

$$Y_1^* = E(y_{1(s)} | \underline{\omega}) = \int_{0}^{\infty} \int_{0}^{\infty} y_{1(s)} f(y_{1(s)}, y_{2(s)} | \underline{\omega}) dy_{2(s)} dy_{1(s)}, \tag{45}$$

and

$$Y_2^* = E(y_{2(s)} | \underline{\omega}) = \int_{0}^{\infty} \int_{0}^{\infty} y_{2(s)} f(y_{1(s)}, y_{2(s)} | \underline{\omega}) dy_{1(s)} dy_{2(s)}, \tag{46}$$

identify the sign of the parameters

From (45) and (46), the point predictors Y_1^* and Y_2^* cannot be obtained in closed form. The joint Bayes points predictors of future ordered observation is

$$Y_1^*, Y_2^* = E(y_{1(s)}, y_{2(s)} | \underline{\omega}) = \int_0^\infty \int_0^\infty y_{1(s)} y_{2(s)} f(y_{1(s)}, y_{2(s)} | \underline{\omega}) dy_{1(s)} dy_{2(s)}. \tag{47}$$

$$Y^*_1, Y^*_2 = ?$$

6. Numerical Illustration

This section aims to investigate the precision of the theoretical results of Bayesian estimation and prediction based on the simulated and real data set.

6.1 Simulation study

In this subsection, a simulation study is conducted to illustrate the performance of the presented Bayes estimates on the basis of generated data from the BCESFLO distribution. Bayes averages of the estimates for the parameters, rf and hrf are computed. Moreover, credible intervals of the parameters, rf and hrf are calculated, Bayes point predictors for a future observation from the BCESFLO distribution are computed for the two-sample case. All simulation studies are performed using R programming language.

Simulation algorithm

A. In similar manner to the steps used in Subsection 4.1, different samples can be generated.

B. The Bayes estimates of $a, b, \alpha_1, \alpha_2, \theta_1$ and θ_2 are obtained by following the steps:

1. Assuming the population parameters and the sample size n .
2. Generate random samples with different sizes (30, 50, and 100) from the population distribution under study.
3. Repeat Step 2, N times, where $N = 10000$.
4. If ω_j^* is an estimate of ω , based on sample $j, j = 1, 2, \dots, N$, then the average estimate over the

samples is given by $\bar{\omega}^* = \frac{1}{N} \sum_{j=1}^N \omega_j^*$.

5. The ER of ω^* , over the N samples is given by

$$ER(\omega^*) = \frac{1}{N} \sum_{j=1}^m (\omega_j^* - \omega)^2.$$

Using Steps 4 and 5, compute $\bar{a}^*, \bar{b}^*, \bar{\alpha}_1^*, \bar{\alpha}_2^*, \bar{\theta}_1^*, \bar{\theta}_2^*$,

$ER(a^*), ER(b^*), ER(\alpha_1^*), ER(\alpha_2^*), ER(\theta_1^*)$ and $ER(\theta_2^*)$.

In the case of two-sample Bayesian prediction

1. Assuming the population parameters and the sample size n .
2. Generate a bivariate random sample of size n , say $(T_1, Y_1), (T_2, Y_2)$ as shown in the beginning of this algorithm.

3. Follow steps in Subsection 5.2.

- The underlying population in Tables 10 and 11 displays the averages estimates, ERs and variances of the Bayes case based on sample of different sizes n and $N=10000$ repetitions with informative prior. The generated population parameters are

$$(a = 0.6, b = 0.8, \alpha_1 = 1.1, \alpha_2 = 1.5, \theta_1 = 1.2, \theta_2 = 1.7)$$

$$\text{and } (a = 2.5, b = 0.67, \alpha_1 = 3, \alpha_2 = 2.5, \theta_1 = 7.4, \theta_2 = 5.1),$$

the given vector of hyper parameters is

$$(c_1 = 0.1, c_2 = 0.2, c_3 = 0.3, c_4 = 0.4, c_5 = 0.5, c_6 = 0.6, c_7 = 0.7, c_8 = 0.8, c_9 = 0.9).$$

Tables 12 and 13 present the Bayes averages, ERs and credible intervals of rf and hrf for different values of the time t_{01}, t_{02} based on informative priors.

- The Bayes two-sample predictors under informative priors are presented in Tables 18 and 19.
- Considering the two-sample prediction and using informative prior, in Tables 18 and 19 the hyper parameters are

$$(c_1 = 0.1, c_2 = 0.2, c_3 = 0.3, c_4 = 0.4, c_5 = 0.5, c_6 = 0.6, c_7 = 0.7, c_8 = 0.8, c_9 = 0.9),$$

$$\text{the population parameters are } (a = 0.6, b = 0.8, \alpha_1 = 1.1, \alpha_2 = 1.5, \theta_1 = 1.2, \theta_2 = 1.7)$$

$$\text{and } (a = 1.5, b = 0.55, \alpha_1 = 5.8, \alpha_2 = 3.5, \theta_1 = 3.5, \theta_2 = 2.5).$$

6.2 Example data set

The data set is given in Subsection 4.2 and analyzed to illustrate the theoretical results of Bayesian estimation and prediction. Tables 14- 17 present the Bayes averages and ERs, of the estimates of the parameters, rf and hrf, for the real data set under informative prior. Bayes predictors and standard errors of the future observation are given in Tables 20.

6.3 Concluding remarks

In our study we observe the following

1. The variance of the estimates is inversely proportional to the sample size and that the variance of an estimate tends to zero as the sample size tends to infinity.
2. The lengths of the CIs of the parameters become narrower as the sample size increases.
3. The Bayes averages for the rf and hrf performs better as the sample size increases. Also, ER is decreasing when the sample size is increasing.

4. It is interesting to notice that if the variables of the prior density are independent and if the likelihood function factors out with respect to these variables, then the variables of the posterior given data are also independent.

- That if $\pi(\omega_1, \dots, \omega_k) = \prod_{i=1}^k \pi(\omega_i)$ and if $L(\omega_1, \dots, \omega_k | \underline{t}) = \prod_{i=1}^k L(\omega_i | \underline{t})$, then $\pi^*(\omega_1, \dots, \omega_k | \underline{t}) \propto \pi(\omega_1, \dots, \omega_k) L(\omega_1, \dots, \omega_k | \underline{t}) = \prod_{i=1}^k \pi(\omega_i) L(\omega_i | \underline{t})$
 $= \prod_{i=1}^k \pi^*(\omega_i | \underline{t}) \Rightarrow (\omega_1 | \underline{t}, \dots, \omega_k | \underline{t}),$

are independent, the analysis will be easier. identify the sign of the parameters, (t?)

5. The likelihood function of the BCESFLO distribution can be derived using the pdf in (16) directly but compounding of $\prod_{i=1}^k f(t_i | \underline{\omega})$ and $\eta(\beta)$ can be applied to make the ML estimation easier. The results become better as the informative sample size gets larger. In all cases, the simulated percentage coverage is at least 95%.

Table 10
Bayes averages, relative absolute biases, estimated risks and
95% credible intervals for the parameters of BCESFLO
(N = 10000, a = 0.6, b = 0.8, $\alpha_1 = 1.1$, $\alpha_2 = 1.5$, $\theta_1 = 1.2$, $\theta_2 = 1.7$)

n	Parameters	Averages	RAB	ER	UL	LL	Length
30	<i>a</i>	0.60059	9.9478e-04	4.6360e-07	0.6012	0.5986	0.0026
	<i>b</i>	0.8007	0.0008	8.9479e-07	0.8019	0.7992	0.0028
	α_1	1.0985	1.3765e-03	2.8541e-06	1.1002	1.0973	0.0029
	α_2	1.4993	0.0015	3.9345e-06	1.2030	1.1998	0.0032
	θ_1	1.2018	0.0005	8.7362e-07	1.5004	1.4982	0.0021
	θ_2	1.7015	0.0009	3.7897e-06	1.7037	1.6998	0.0039
50	<i>a</i>	0.5998	0.0004	4.4439e-07	0.6006	0.5982	0.0024
	<i>b</i>	0.7996	4.5745e-4	6.5481e-07	0.8008	0.7982	0.0026
	α_1	1.0997	0.0003	3.6181e-07	1.1001	1.0978	0.0023
	α_2	1.5002	9.6223	1.7531e-06	1.2023	1.1996	0.0027
	θ_1	1.2011	1.1497e-04	3.7496e-07	1.5012	1.4989	0.0024
	θ_2	1.7006	3.6611e-04	8.0349e-07	1.7016	1.6994	0.0021
100	<i>a</i>	0.5997	5.3088e-04	2.3494e-07	0.6004	0.5988	0.0017
	<i>b</i>	0.7999	7.8312e-05	3.9512e-07	0.8009	0.7986	0.0024
	α_1	1.0999	4.1095e-05	2.9580e-07	1.1007	1.0986	0.0022
	α_2	1.5001	6.3095e-04	8.3359e-07	1.2002	1.1982	0.0019
	θ_1	1.1992	3.4961e-05	3.2374e-07	1.5007	1.4988	0.0019
	θ_2	1.6996	2.1334e-04	4.4509e-07	1.7006	1.6985	0.0021

Table 11
Bayes averages, relative absolute biases, estimated risks
and 95% credible intervals for the parameters of BCESFLO
(N = 10000, a = 2.5, b = 0.67, $\alpha_1 = 3$, $\alpha_2 = 2.5$, $\theta_1 = 7.4$, $\theta_2 = 5.1$)

n	Parameters	Averages	RAB	ER	UL	LL	Length
30	<i>a</i>	2.5014	0.0005	2.6580e-06	2.5024	2.4994	0.0029
	<i>b</i>	0.6710	0.0015	2.0137e-06	0.6724	0.6694	0.0029
	α_1	3.0017	0.0005	4.5668e-06	3.0044	2.9998	0.0045
	α_2	2.5017	0.0007	3.9767e-06	2.5029	2.4998	0.0030
	θ_1	7.3985	0.0006	3.2725e-06	7.4001	7.3970	0.0020
	θ_2	5.1023	0.0004	7.0838e-06	5.1039	5.0998	0.0041
50	<i>a</i>	2.4991	0.0003	1.0481e-06	2.4999	2.4978	0.0020
	<i>b</i>	0.6709	0.0013	1.8134e-06	0.6720	0.6695	0.0025
	α_1	3.0007	0.0002	1.1669e-06	3.0025	2.9982	0.0043
	α_2	2.4984	0.0001	3.0778e-06	2.4999	2.4971	0.0028
	θ_1	7.4008	0.0001	9.0744e-07	7.4016	7.3995	0.0020
	θ_2	5.1013	2.6363e-04	2.5136e-06	5.1029	5.0996	0.0033
100	<i>a</i>	2.4994	2.1057e-04	6.5731e-07	2.5002	2.4984	0.0018
	<i>b</i>	0.6698	2.5083e-04	4.2225e-07	0.6710	0.6686	0.0023
	α_1	3.0001	6.5163e-05	9.1350e-07	3.0015	2.9974	0.0041
	α_2	2.5003	5.7028e-05	1.3608e-06	2.5010	2.4993	0.0016
	θ_1	7.4004	1.4222e-04	3.3561e-07	7.4023	7.4005	0.0018
	θ_2	5.0988	0.0002	1.5726e-06	5.0997	5.0975	0.0021

Table 12
Bayes averages, relative absolute biases, estimated risks
and 95% credible intervals for the reliability and
hazard rate functions of BCESFLO

(N = 10000, a = 0.6, b = 0.8, $\alpha_1 = 1.1$, $\alpha_2 = 1.5$, $\theta_1 = 1.2$, $\theta_2 = 1.7$, $t_{01} = 2$, $t_{02} = 4$)

n	rf and hrf	Averages	RAB	ER	UL	LL	Length
30	$R(t_{01}, t_{02})$	0.2886	0.0006	5.4372e-07	0.2897	0.2874	0.0024
	$h(t_{01}, t_{02})$	0.0032	0.4062	6.6236e-06	0.0055	0.0012	0.0042
50	$R(t_{01}, t_{02})$	0.2881	0.0011	3.3358e-07	0.2888	0.2868	0.0019
	$h(t_{01}, t_{02})$	0.0055	0.0387	2.8049e-07	0.0063	0.0042	0.0022
100	$R(t_{01}, t_{02})$	0.2884	6.3252e-05	1.2286e-07	0.2890	0.2877	0.0014
	$h(t_{01}, t_{02})$	0.0055	2.6847e-02	2.7532e-07	0.0062	0.0043	0.0019

Table 13
Bayes averages, relative absolute biases, estimated risks and
95% credible intervals for the reliability and
hazard rate functions of BCESFLO

(N = 10000, a = 1.5, b = 0.55, $\alpha_1 = 5.8$, $\alpha_2 = 3.5$, $\theta_1 = 3.5$, $\theta_2 = 2.5$, $t_{01} = 2$, $t_{02} = 3$)

n	rf and hrf	Averages	RAB	ER	UL	LL	Length
30	$R(t_{01}, t_{02})$	0.0035	0.7896	5.5620e-06	0.0061	0.0009	0.0052
	$h(t_{01}, t_{02})$	0.0472	0.0402	4.2526e-06	0.04841	0.0449	0.0034
50	$R(t_{01}, t_{02})$	0.0025	0.2684	4.1521e-07	0.0031	0.0014	0.0017
	$h(t_{01}, t_{02})$	0.0441	0.0268	2.5180e-06	0.0453	0.0429	0.0024
100	$R(t_{01}, t_{02})$	0.0019	0.0051	1.4099e-07	0.0026	0.0009	0.0016
	$h(t_{01}, t_{02})$	0.0456	0.0059	2.8211e-07	0.0464	0.0447	0.0017

Table 14
Bayes estimates and standard errors
for the parameter of BCESFLO

Parameters	Estimate	Se
a	0.6013	0.0009
b	0.8001	0.0008
α_1	1.1025	0.0009
α_2	1.5003	0.0008
θ_1	1.2003	0.0007
θ_2	1.6999	0.0004

Table 15
Bayes estimates and standard errors
for the parameters of BCESFLO

Parameters	Estimates	Se
a	1.5018	0.0014
b	0.5510	0.0005
α_1	5.8005	0.0007
α_2	3.5027	0.0008
θ_1	3.5027	0.0014
θ_2	2.4978	0.0013

Table 16
Bayes estimates and standard errors for the
reliability and hazard rate functions of BCESFLO

rf and hrf	Estimates	Se
$R(t_{01}, t_{02})$	0.2897	0.0004
$h(t_{01}, t_{02})$	0.0035	0.0008

Table 17
Bayes estimate and standard errors for
the reliability and hazard rate functions

n	rf and hrf	Estimate	Se
30	$R(t_{01}, t_{02})$	0.0029	0.0009
	$h(t_{01}, t_{02})$	0.0461	0.0005

Table 18
Bayes predictors, relative absolute biases, estimated risks
and 95% credible interval of the future observation
(N = 10000, a = 0.6, b = 0.8, $\alpha_1 = 1.1$, $\alpha_2 = 1.5$, $\theta_1 = 1.2$, $\theta_2 = 1.7$)

n	s	$\hat{Y}_{1(s)}$	Averages	RAB	ER	UL	LL	Length
30	1	$\hat{Y}_{1(s)}$	3.9999	2.7731e-05	5.0017e-07	4.0009	3.9982	0.0026
		$\hat{Y}_{2(s)}$	7.0003	4.0064e-05	3.5349e-07	7.0011	6.9989	0.0021
	12	$\hat{Y}_{1(s)}$	4.0003	6.6825e-05	8.0271e-07	4.0015	3.9986	0.0029
		$\hat{Y}_{2(s)}$	6.9982	2.5996e-04	3.7367e06	6.9994	6.9968	0.0026
	18	$\hat{Y}_{1(s)}$	4.0009	0.0002	1.6289e-06	4.0022	3.9992	0.0031
		$\hat{Y}_{2(s)}$	6.9981	0.0003	5.1302e-06	7.0001	6.9959	0.0042
50	1	$\hat{Y}_{1(s)}$	4.0005	1.1660e-04	4.6721e-07	4.0013	3.9993	0.0019
		$\hat{Y}_{2(s)}$	7.0000	2.0907e-06	1.6751e-07	7.0006	6.9989	0.0017
	12	$\hat{Y}_{1(s)}$	4.0007	1.6923e-04	1.0526e-06	4.0016	3.9989	0.0027
		$\hat{Y}_{2(s)}$	6.9994	8.2851e-05	5.9514e-07	7.0004	6.9983	0.0020
	18	$\hat{Y}_{1(s)}$	4.0014	0.0004	3.2526e-06	4.0028	3.9998	0.0030
		$\hat{Y}_{2(s)}$	7.0008	0.0001	1.5965e-06	7.0024	6.0024	0.0033
100	1	$\hat{Y}_{1(s)}$	3.9997	6.9537e-05	2.9892e-07	4.0006	3.9989	0.0017
		$\hat{Y}_{2(s)}$	7.0004	51594e-05	2.9367e-07	7.0009	6.9994	0.0016
	12	$\hat{Y}_{1(s)}$	3.9997	8.0968e-05	3.7275e-07	4.0005	3.9983	0.0022
		$\hat{Y}_{2(s)}$	7.0014	2.0482e-04	2.4549e-06	7.0023	6.9996	0.0027
	18	$\hat{Y}_{1(s)}$	4.0002	5.7244e-05	7.7567e-07	4.0016	3.9983	0.0032
		$\hat{Y}_{2(s)}$	6.9985	2.1164e-04	3.8607e-06	7.0005	6.9964	0.0041

Table 19
Bayes predictors, relative absolute biases, estimated risks and
95% credible intervals of the future observation
(N = 10000, a = 1.5, b = 0.55, $\alpha_1 = 5.8$, $\alpha_2 = 3.5$, $\theta_1 = 3.5$, $\theta_2 = 2.5$)

n	s	$\hat{Y}_{(s)}$	Averages	RAB	ER	UL	LL	Length
30	1	$\hat{Y}_{1(s)}$	3.9986	0.0003	3.1005e-06	4.0007	3.9982	0.0025
		$\hat{Y}_{2(s)}$	7.0008	1.1791e-04	1.1019e-06	7.0016	6.9991	0.0026
	12	$\hat{Y}_{1(s)}$	3.9983	4.3381e-04	3.2831e-06	3.9993	3.9957	0.0036
		$\hat{Y}_{2(s)}$	6.9994	7.2306e-05	1.6241e-06	7.0013	6.9969	0.0043
	18	$\hat{Y}_{1(s)}$	3.9978	5.4197e-04	6.6847e-06	3.9999	3.9956	0.0043
		$\hat{Y}_{2(s)}$	7.0022	0.0003	7.1597e-06	7.0041	6.9994	0.0048
50	1	$\hat{Y}_{1(s)}$	3.9997	7.5194e-05	3.0675e-07	4.0005	3.9989	0.0017
		$\hat{Y}_{2(s)}$	7.0002	2.6742e-05	2.5213e-07	7.0011	6.9990	0.0021
	12	$\hat{Y}_{1(s)}$	4.0001	2.7094e-05	4.2356e-07	4.0010	3.9986	0.0024
		$\hat{Y}_{2(s)}$	6.9983	0.0002	3.5432e-06	6.9999	6.9971	0.0029
	18	$\hat{Y}_{1(s)}$	4.0008	0.0002	1.0349e-06	4.0018	3.9993	0.0025
		$\hat{Y}_{2(s)}$	7.0023	3.3174e-04	7.4926e-06	7.0049	6.9995	0.0054
100	1	$\hat{Y}_{1(s)}$	3.9997	8.631e-05	3.6679e-07	4.0004	3.9989	0.0015
		$\hat{Y}_{2(s)}$	6.9999	1.5412e-05	2.2928e-07	7.0006	6.9988	0.0019
	12	$\hat{Y}_{1(s)}$	4.0007	0.0001	1.0153e-06	4.0019	3.9995	0.0024
		$\hat{Y}_{2(s)}$	6.9989	0.0001	1.8428e-06	7.0003	6.9975	0.0028
	18	$\hat{Y}_{1(s)}$	3.9996	9.3514e-05	1.1006e-06	4.0010	3.9979	0.0031
		$\hat{Y}_{2(s)}$	7.0027	3.9138e-04	9.4936e-06	7.0048	6.9999	0.0049

Table 20
Bayes predictors and standard errors
of the future observation

s	$\hat{y}_{(s)}$	Estimate	Se
1	$\hat{y}_{1(s)}$	3.9999	0.0005
	$\hat{y}_{2(s)}$	7.0009	0.0006
12	$\hat{y}_{1(s)}$	3.9993	0.0007
	$\hat{y}_{2(s)}$	7.0021	0.0007
18	$\hat{y}_{1(s)}$	3.9987	0.0008
	$\hat{y}_{2(s)}$	7.00248	0.0019

REFERENCES

- AL-Hussaini, E. K. and Ateya, S. F. (2005).** Parametric estimation under a class of multivariate distributions. *Statistical Papers*, 46, 321-338.
- AL-Hussaini, E. K. (2011).** A class of compound exponentiated survival function and multivariate extension. *Journal of Statistical Theory and Applications*, 10(1), 63-84.
- Arnold, B. C. (1983).** *Pareto distributions*. Silver Spring, Maryland: International Cooperative Publishing House, Fairland, MD.
- Arnold, B. C., Castillo, E. and Sarbia, J. M. (1999).** *Conditional specification of statistical models*. Springer Series in Statistics, Springer Verlag, New York.
- Arnold, B. C., Castillo, E. and Sarbia, J. M. (2001).** Conditionally specified distributions; an introduction with discussion. *Statistical Science*, 16, 249-274.
- Balakrishnan, N. and Lai, C. D. (2009).** *Continuous bivariate distributions*, Second Edition Springer Science.
- Chadi, A., Talhi, H. and Fellag, H. (2013).** Comparison of the maximum likelihood and Bayes estimators for symmetric bivariate exponential distribution under different loss functions. *Afrika Statistika*, 8(1), 499-514.
- Iliopoulos, G., Karlis, D. and Ntzoufras, I. (2005).** Bayes estimation in Kibbles bivariate gamma distribution. *The Canadian journal of Statistics*, 33(4), 571-589.
- Kotz, S., Balakrishnan, N. and Johnson, N. L. (2000).** *Continuous multivariate distributions: models and Applications*. 1, Second Edition, Wiley, New York.

Lin, Y. J., Lio, Y. L. and Ng, H. K. T. (2013). Bayes estimation of Moran- Downton bivariate exponential distribution based on censored samples. *Journal of Statistical Computation and Simulation*, 83, 837-852.

Mardia, K. V. (1962). Multivariate Pareto distribution. *Annals of Institute of Statistical Mathematics*, 33, 1008-1015.

Muliere, P. and Scarini. M. (1987). Characterization of the Marshall-Olkin type class of distributions. *Annals of the Institute of Statistical Mathematics*. 39, 429-441.

Padamadan, V. and Nair, K. R. M. (1994). Characterization of a bivariate Pareto distribution. *Journal of the Indian Statistical Association*, 32, 15-20.

Pradhan, B. and Kundu, D. (2015). Bayes estimation for the Block and Basu bivariate and multivariate Weibull distributions. *Journal of Statistical Computation and Simulation*, 86(1), 170-182.

Sankaran, P. G. and Kundu, D. (2014). A bivariate Pareto model. *Journal of Theoretical and Applied Statistics*, 48(2), 241-255.

Sarabia Alegria, J. M. and Gómez-Déniz, E. (2008). Construction of Multivariate distributions: a review of some recent results. *Statistics and Operations Research Transactions*, 32(1), 3-36.

Tierney, L. and Kadane, J. B. (1986). Accurate approximations for posterior moments and marginal densities. *Journal of the American Statistical Association*, 81(393), 82-86.