

## Case study

### Aicardi-Goutières syndrome: a case report

#### Abstract :

The Aicardi-Goutières syndrome is an encephalopathy characterized by an arrest of psychomotor development in the first months of life, spastic tetraplegia spastic tetraplegia, abnormal movements and acquired microcephaly. Neuroimaging reveals calcifications of the basal ganglia and cerebrospinal fluid (CSF) study shows lymphocytosis. Our patient is a 7 month old child, from a 1st degree consanguineous marriage. She presents a psychomotor delay with hypotonia, nystagmus and convulsive myoclonic seizures. Cerebral CT shows an active tri ventricular hydrocephalus with multiple parenchymal calcifications with multiple cerebral parenchymal calcifications and basal ganglia of infectious appearance. The intercritical sleep EEG was unremarkable. Ophthalmological examination, including an EPI, revealed a probable retinal dystrophy predominantly on the scopic system. Brain MRI shows a mesencephalic atrophy of the annular protuberance and increased volume of the volume of the cisterns of the posterior cerebral fossa. The dosage of interferon alpha in the CSF is increased to 113 pg /ml.

#### Introduction:

Aicardi-Goutières syndrome is a progressive encephalopathy characterized by arrested psychomotor development in the first months of life, spastic tetraplegia, abnormal movements, and acquired microcephaly. Neuroimaging shows calcifications of the basal ganglia and cerebrospinal fluid (CSF) study shows a lymphocytosis.

**Comment [m1]:** This part is a repetition of a paragraph of the abstract

#### Observation:

Our patient is a 7 month old child, from a consanguineous marriage of the 1st degree and a 29 years old mother, G1P1, grouping B+, pregnancy well followed, the delivery took place by caesarean section for exceeding the term, birth weight= 2kg600 with immediate cry, put under exclusive breastfeeding. She presented a psychomotor delay with hypotonia, nystagmus and myoclonic convulsive seizures. Cerebral CT shows an active tri ventricular

hydrocephalus with multiple cerebral parenchymal calcifications and basal ganglia of infectious appearance.

The intercritical sleep EEG was unremarkable. Ophthalmological examination including an EPI revealed a probable retinal dystrophy predominating on the scopic system. Brain MRI shows a mesencephalic atrophy with atrophy of the annular protuberance and increased volume of the cisterns of the posterior cerebral fossa. The interferon alpha level in the CSF is increased to 113 pg /ml. The genetic study is in progress.

On these clinical, neuroradiological, ophthalmological signs and biochemical results, we concluded to an Aicardi-Goutières Syndrome.

**Discussion:**

Aicardi-Goutières syndrome (AGS) was first described in 1984 by Jean Aicardi and Françoise Goutières two eminent pediatric neurologists working at Necker Hospital in Paris, France [1]. They reported eight children from five families with an early onset of central nervous system disorder with bilateral spasticity and dystonia and acquired microcephaly. The course was rapid to death.

This syndrome was defined as the triad of infantile spasms, agenesis of the corpus callosum and chorioretinal lacunae. [2]. The syndrome is genetically heterogeneous with mutations in 7 genes, including TREX1, RNASEH2A, RNASEH2A, RNASEH2B, RNASEH2C, SAMHD1, ADAR1 and IFIH1 (interferon-induced helicase c domain-containing protein 1) [3].

Aicardi-Goutières syndrome (AGS) is an inflammatory mendelian leukoencephalopathy. Aicardi-Goutières syndrome (AGS) is an inflammatory mendelian leukoencephalopathy, part of the type I interferonopathies - a group of neuroinflammatory phenotypes associated with increased type I interferon (IFN) signaling, characterized by basal ganglia calcification, white matter abnormalities, and cerebrospinal fluid (CSF) lymphocytosis. It is a rare autosomal recessive neuro-pediatric syndrome with an encephalopathy beginning before the age of one year and of progressive aggravation, leading to a loss of acquisitions then a severe intellectual and psycho-motor deterioration.

The symptoms evolve over a few months with the appearance of microcephaly and a pyramidal syndrome before stabilizing. However, more moderate forms have been

described with onset after 1 year of age, relative preservation of language and cognitive functions, and normal head circumference. Inter- and intra-familial phenotypic variability is usually observed.

The diagnosis of Aicardi-Goutières syndrome is still often misunderstood. Epileptic spasms are often early, as in our patient's case, and occur before the age of 3 months, and as early as the neonatal period in 18% of cases [4]. On EEG, true hypsarrhythmia remains rare, the most typical aspect being asynchronous pseudoperiodic discharges on both hemispheres [5]. Chorioretinal lacunae are found and are yellowish white, rounded with a clear unpigmented border. They are most often multiple and bilateral and are pathognomonic of the syndrome. Other ocular abnormalities may be associated with chorioretinal lacunae such as unilateral colobomas of the optic nerve, iris and peripapillary choroid [6], microphthalmia often unilateral, microcornea, calcified cataracts, iris synechiae and pupillary membrane remnants, optic nerve hypoplasia, optic atrophy, retinal detachment, or sclerectasia [7].

CT scan shows calcifications of the basal ganglia and MRI shows suggestive hypodensities in the white matter. The increase of interferon alpha in the CSF is almost constant. Cerebral histological specimens may show microinfarcts and calcifying vasculitis of the brain and systemic vessels.

TORCH complex embryo-fetopathies, neonatal enteroviruses, Cockayne syndrome, hereditary progressive CNS degeneration with generalized intracranial calcifications are the main differential diagnoses to be ruled out. [8].

The prognosis of these patients is guarded; the evolution is usually towards profound mental retardation, blindness and the absence of any motor or language acquisition.

The treatment remains symptomatic, consisting in the management of eating disorders, psychomotor deficits, and possible epilepsy. The evolution towards a profound deterioration and death is generally severe and rapid.

### **Conclusion**

Aicardi-Goutières syndrome is a rare encephalopathy. This observation illustrates the interest of making the diagnosis and the importance of genetic counseling because there is a

risk of recurrence in a subsequent pregnancy. Prenatal diagnosis can be made by molecular biology on amniotic fluid or trophoblast.

#### References:

- [1] Yanick J. Crow1-3 and Nicolas Aicardi-Goutières syndrome and the type I interferonopathies *Nature Reviews Immunology* | AOP 2015
- [2] Aicardi J, Chevrie JJ, Rousselle F (1969) Le syndrome spasmes en flexion agenesie calleuse, anomalies chorio-retiniennes. *Arch Fr Pediatr* 26:1103-1120
- [3] Ahmed Gilani1 , Laura A Adang2 , Adeline Vanderver2 , Abigail Collins *Pediatric and Developmental Pathology* 1-5 2019, Society for Pediatric Pathology  
Neuropathological Findings in a Case of IFIH1-Related Aicardi-Goutières Syndrome
- [4] Jeavons PM, Livet M. West's syndrome. In: Roger J, editor. *Epileptic syndromes of the child and adolescent*. Paris: John Libbey; 1992. p. 53-66
- [5] Bour F, Chiron C, Dulac O, Plouin P. Electroclinical features of seizures in Aicardi syndrome. *Rev EEG Neurophysiol Clin* 1986;16:341-53.
- [6] Béby F, Zech C, Touraine R, Guibaud P, Masset H, Trepsat CH, et al. Persistent and hyperplastic primary vitreous syndrome in a girl with Aicardi syndrome. *J Fr Ophthalmol* 2000;23: 703-7
- [7] Aicardi J. Aicardi syndrome. *Brain Dev* 2005;27:164-71.
- [8]. Menezes AV, Lewis TL, Buncic JR. Role of ocular involvement in the prediction of visual development and clinical prognosis in Aicardi syndrome. *Br J Ophthalmology* 1996;80:805-11.

**Comment [m2]:** Since there is no immediate solution, it would not be better to educate the couples not to marry at all. Not to warn them about the risks to their offspring afterwards.

**Comment [m3]:** It is necessary to homogenize the writing of the bibliography (date) according to the required format