

Evaluation of Hepatic Enzyme levels and Haematological Profiles in Chikungunya patients- An institutional study

Abstract

Background: Chikungunya is a viral infection caused by the Chikungunya virus (CHIKV) belonging to the Togaviridae family. The transmission of the virus is taking place through the bite of infected daytime biting female mosquitoes – primarily *Aedes aegypti* and *Aedes albopictus*. There were many reports which highlighted the number of reported cases of Chikungunya has been increasing in India in recent years, especially since 2016-2017.

Materials and Methods: Blood samples were collected from patients diagnosed with having Chikungunya. Routine laboratory investigations were carried out for analysing haematological and serum enzyme profiles.

Result: Our results highlighted that the alterations of haematological and hepatic dysfunction are prevalent in Chikungunya infection, with SGPT rising significantly more than SGOT. Hepatic Enzymes levels appear to have a directly proportional correlation with severity of infections.

Conclusion: Our study demonstrated that haematological and hepatic enzymes levels should be explored as routine laboratory markers for assessing the severity of Chikungunya infection, as they will help in employing an appropriate patient therapy, and thus optimise the use of available resources.

Keywords: Chikungunya, Dengue, Hepatic, Haematological

Introduction

Chikungunya is a viral illness transmitted by mosquitoes affecting all age groups, but severe complications are more often observed in children. Over the past two decades, there has been global increase in the frequency of Chikungunya and its epidemics. Symptoms generally start 4-7 days after the mosquito bite. The acute phase is characterised by joint and muscle pain, high fever, extreme weakness, headache, vomiting, and rashes. In the chronic phase, various neurological syndromes and non-neurological manifestations can occur [1]. In 2006, the CHIK virus (CHIKV) re-emerged in India after 32 years, causing the epidemic affecting more than 1.4 million people across the 13 States, and post epidemic, a declining trend was seen till 2011 [1-3].

The WHO South-East Asia Regional Office has reported a significant number of Chikungunya cases in almost 151 districts in 8 states of India suffered from many cases of chikungunya fever. The most affected states are Andhra Pradesh, Andaman & Nicobar Islands, Tamil Nadu, Karnataka, Maharashtra, Gujarat, Madhya Pradesh, Kerala and Delhi [1,2]. As usual, the outbreaks of Chikungunya infections are most likely to occur in the post-monsoon period when the vector density significantly increases. There are 3 epidemiological factors: the host (man and mosquito), agent (virus), environment (abiotic and biotic factors) agent for Chikungunya viruses. Human beings are considered as the reservoir for the chikungunya virus during the epidemic time.

The rapid spread of the disease in Southern part of India from 2004 has afflicted millions of people and left many of them with crippling disabilities. Chikungunya disease continues to cause epidemics in many countries in the region. The patient in this context acts as the reservoir of infection for others in the household and the community. Therefore, public health measures to minimise the transmission of infection become imperative to prevent and control the outbreak from spreading [3-5].

Comment [VFS1]: References on the epidemiology of chikungunya in India are old, it should be updated with reference to the study period. Furthermore, the impact of the COVID-19 epidemic during that period could be indicated.

Fever with or without arthralgia is a prevalent manifestation of several other diseases. CHIK fever may not have typical manifestations, or it may co-exist with other infectious diseases like dengue fever or non-infectious diseases like rheumatoid arthritis [6].

The analysis and diagnosis of the case in the community need to be communicated immediately to the nearest public health official for identification of clusters by person, place and time and expansion of the control measures in the community and district levels.

The most significant features seen in CHIKV-positive patients are lymphopenia, leukopenia, hypocalcemia, and elevated transaminases; but these features were not consistently reported in all studies and vary depending the clinical phase of infection.

The general laboratory diagnosis of Chikungunya often reveals lymphopenia, neutrophils and thrombocytopenia, an elevated creatinine, and elevated hepatic transaminases. In this study we tried to establish the relationship between the severity of Chikungunya infection and its impact on levels hepatic enzymes and haematological profiles of individuals.

Comment [VFS2]: What are the bibliographic references in this paragraph?

Comment [VFS3]: What are the bibliographic references in this paragraph?

UNDER PEER REVIEW

Materials and Methods

Recruitments of Patients and Healthy Individuals: This study was carried out in the Department of Biochemistry and General medicine at mamata Medical College and general Hospital, Khammam, Telangana from the period of July 2021 to November 2021.

Study Design: It was a prospective cohort study over two years through sample and sampling techniques. A total of 50 patients admitted to the hospital with a history of fever of more than 38.5°C and CHICKV positive were selected using purposive sampling techniques. Clinical signs and symptoms, severity, and outcomes with relevant laboratory parameters were compared in detail. They are followed from the onset of fever to time of recovery or discharge according to WHO discharge criteria whichever is earlier. Same numbers of normal healthy individuals were also recruited for this study. Duly signed Informed consent forms were collected from all the participants during the study.

Routine Laboratory Investigations: The 5-7ml of blood samples were collected from all these participants' investigations were done- blood counts, clinically patients are monitored and for platelets, hematocrit count, haemoglobin count, total leukocyte count.

Liver Function Test: SGPT, SGOT were estimated by using protocol prescribed by the International Federation of Clinical Chemistry. ALP, Total Protein, Albumin, Total Bilirubin were estimated by colourimetric assay.

Statistical analysis: The collected data analysis was done using SPSS software. The results were presented as mean \pm standard deviation (SD) and percentages. Mean, and SD was computed for all continuous variables and comparisons were done using Student's t-test.

Comment [VFS4]: were the patients followed for two years? nothing is mentioned about the results of the follow-up period

Comment [VFS5]: it remains to describe the laboratory techniques used for diagnostic confirmation. serological or molecular tests or both?

Comment [VFS6]: information on the ethical procedures of the study is lacking. What was the ethics committee that authorized the study? Approval number, Inclusion and exclusion criteria, etc.

Comment [VFS7]: Taking into account that it is a prospective cohort, were control hematological and biochemical analyzes performed? It would be great to be able to include and discuss those results

Comment [VFS8]: It is indicated that the Student's t test was used for the bivariate analysis of all the quantitative variables, all these variables had a normal distribution?

Results

The total numbers of 50 cases of acute febrile illness were recruited based on their clinical report as seropositive for CHIKV infection, an equal number of healthy individuals also participated in this study. There were altogether 67 (63.6%) men and 33 (36.4%) women participants. Most of the participants were in the 20–30-year age group. A seasonal peak with 42 (34.7%) cases was seen in monsoon season. The patient characteristics and salient clinical features of acute chikungunya infection, and biochemical parameters are compared in Tables 1 and 2. The rural residency was prominent in all groups. Fever was seen in all cases (100%). The symptomology of chikungunya included headaches, joint pain, weakness, and restricted joint movement. No complications were reported. Deranged liver function was seen in all patients (100%).

Table 1: Comparison of characteristics and clinical features in between normal healthy individuals and patients groups.

Characteristics	Control Group	Patient Group
Age Group (Years)	<20	4 (8)
	20-30	21 (42)
	31-40	11 (22)
	41-50	3 (6)
	51-60	2 (4)
	>60	0
Gender	Male	37 (74)
	Female	23 (46)
Residence	Urban	22 (44)
	Rural	28 (56)
Symptoms	Fever	50 (100)
	Joint Pain	31 (62)
	Backache	15 (30)
	Vomiting	4 (8)

Comment [VFS9]: Different studies have shown that chikungunya-dengue coinfection is very common in endemic countries (as has already been demonstrated in previous studies in India). Is it possible that this coinfection is frequent in the study area? And were the alterations in the hematological parameters and hepatic enzymes produced due to this coinfection? [as in the study carried out by Kularatne et al (2009) in Sri Lanka (reference 14)]

Out of 50 patients, 37 were male and 23 were female and in control group of 50 healthy individuals 40 were male and 10 were female. Most of the participants were in the age group of 20 – 40 and none of them were below 20 years and above 70 years of age. All the patients had fever as a presenting complaint. Most of them had myalgia at the time of presentation.

Table 2: Comparison of biochemical parameters in between normal healthy individuals and patients groups.

Parameters	Control Group	Patient Group	p Value
Hb	13.15±1.82	11.55±2.36	0.5012
TLC	8560±2634	4781±2219	0.0082
PC	155,700±23,523	67,622±44,254	0.0015
HCT	36.67±4.52	38.66±6.07	0.523
SGPT	28.87±4.23	239.68±589.25	0.5215
SGOT	25.92±6.22	345.52±1356.52	0.633
ALP	34.21±14.21	144.21±110.26	0.2231
Total Bilirubin	0.98±0.29	1.84±0.28	0.001
Direct Bilirubin	0.51±0.18	0.98±0.33	0.001
Total Protein	7.4±0.59	6.29±0.89	0.0747
Albumin	4.11±0.47	3.15±0.69	0.0130

Hb: Hemoglobin, TLC: Total leukocyte count, PC: Platelets count, HCT: Hematocrit, RBS: Random blood sugar, ALP: Serum alkaline phosphatase, SGPT: Serum glutamate pyruvate transaminase, SGOT: Serum glutamic-oxaloacetic transaminase.

Table 3: Comparison of deranged liver enzymes in between normal healthy individuals and patients groups.

Parameters	Control Group	Patient Group
SGPT (U/L)	60-180 (%)	1 (2)
	181-500 (%)	0
	501-1000 (%)	5 (10)
	>1000 (%)	4 (8)
SGOT (U/L)	38-120 (%)	0
	121-500 (%)	1 (2)
	500-1000 (%)	9 (18)
	>1000 (%)	3 (6)

Elevated liver enzymes were found in almost all patients. Of the liver enzymes, SGPT levels were significantly higher than SGOT levels.

Table 4: Comparison of thrombocytopenia in between normal healthy individuals and patients groups.

Platelets Count (μL)	Control Group	Patient Group
>150,000	26 (52)	2 (4)
150,000-100,000	23 (46)	5 (10)
100,000-50,000	1(2)	20 (40)
50,000-20,000	0	13 (26)
<20,000	0	6 (12)
<10,000	0	3 (6)
Platelets transfusion	0	1 (2)

Thrombocytopenia was noted in 43 patients, of whom 12 had bleeding, and 31 had no bleeding. Statistically, the Chi-square test was used to test the significance of thrombocytopenia and bleeding tendencies. There was no significant difference ($p = 0.945$) found between the variations in the thrombocytopenia and the bleeding tendencies. This implies that bleeding tendencies does not depend upon thrombocytopenia.

Discussion

Currently, Chikungunya is causing significant public health concern throughout the world, particularly in South-East Asian countries. Recently dengue outbreaks caused significant morbidity and mortality in certain parts of India mainly in Andhra Pradesh, Uttar Pradesh and Tamil Nadu [7]. Hepatic dysfunctions in dengue are common. It is due to either direct effect of the virus on hepatocytes or due to reactive hepatitis. Hepatic involvement in dengue fever is in the form of elevated serum aminotransferase [8]. Those patients with elevated liver enzymes are more likely to have increased risk of bleeding tendencies, shock, ARDS, renal failure. In addition to decreased platelet count, hepatic dysfunction plays a significant role in

bleeding. Hence, it is mandatory to evaluate serum aminotransferases in all patients with dengue fever [9]. It is observed that, out of 50 patients, most of the patients had elevated liver enzymes, SGPT levels were significantly higher than SGOT levels, and these patients had severe complications.

Chikungunya viral infection is not generally fatal but can cause neurological and optical manifestations. Severe joint pain is the prominent clinical manifestation and can persist for months to a year [10]. As Dengue Fever (DF) has a high incidence and mortality rate, symptomatic patients are tested only for DENV and only in rare cases for chikungunya viral infection. This scenario is an important reason why chikungunya cases go undiagnosed in dengue-endemic regions, and the actual burden of the chikungunya viral infection has been missed. Thus, investigation for both viruses should be done, especially in endemic regions. Accurate and early diagnosis of coinfections would help inappropriate management [11, 12].

The number of cases increases during and after the monsoon months because higher humidity lengthens the life span of mosquitoes and increased temperatures shorten the extrinsic incubation period [13]. Rising of the cases during the monsoon period with highest cases in different months had been reported in previous studies from India [11-14]. November was observed as a peak among all the months of the year in this study.

Earlier studies have reported increased incidences of atypical hemorrhagic manifestations [12-14] and hepatic dysfunction [15]. Hepatic dysfunction was the most common complication in our dengue and coinfection groups. SGPT levels were significantly higher than SGOT levels.

Conclusion

Haematological profile alteration and Hepatic involvement are common in Chikungunya. It is characterized by elevated liver enzymes, i.e. SGPT more than SGOT levels. Elevated liver enzymes are associated with complications like bleeding, shock and organ impairment. In

Comment [VFS10]: Of the references cited in the last paragraph of the discussion, with the exception of reference 14, they do not mention analysis of hematological and biochemical parameters in chikungunya. In addition, reference 14 mentions findings of alteration but in people with DENV-CHIKV co-infection

addition to thrombocytopenia, hepatic involvement plays a significant role in bleeding. Elevated liver enzymes have got prognostic value in this study. Hence, liver enzymes are mandatory in dengue fever to look for complications, and it is of prognostic value. Those patients with elevated liver enzymes should be monitored carefully than those patients with normal liver enzymes.

Comment [VFS11]: The conclusions of the study would be subject to resolving the previous observations

UNDER PEER REVIEW

References

1. World Health Organisation. Strengthening implementation of the global strategy for dengue fever and dengue haemorrhagic fever, prevention and control. Report of the informal consultation. 18 – 20 October 1999. Geneva: WHO; 1999.
2. World Health Organisation. Scientific working group on dengue. Meeting report, Geneva, Switzerland, 3-5 April 2000. Geneva:WHO;2000.
3. Singhi S, Kissoon N, Bansal A. Dengue and dengue hemorrhagic fever: Management issues in an intensive care unit. *J Peiatr (Rio J)*. 2007;83(2 suppl): S22-35.
4. Satarsinghe RL, Arulnithy K, Amerasena NL, Bulugahapitiya U, Sahayam DV. Asymptomatic myocardial involvement in acute dengue virus infection in a cohort of adult Sri Lankans admitted to a tertiary referral centre. *Br J Cardiol* 2007; 14:171-3.
5. Kamath SR, Ranjit S. Clinical features, complications and atypical manifestations of children with severe forms of dengue hemorrhagic fever in South India. *Indian J Pediatr* 2006; 73:889-5.
6. Singh M. *Medical emergencies in children*. 4th Ed. New Delhi: Sagar Publications; 2004.
7. Deeba F, Afreen N, Islam A, Naqvi IH, Broor S, Ahmed A, Parveen S. Co-infection with Dengue and Chikungunya Viruses. In: RodriguezMorales A, editor. *Current Topics in Chikungunya*. InTech; 2016 (London-United Kingdom) DOI:10.5772/64308. Available from: [http:// www.intechopen.com/books/current-topics-in-chikungunya/co-infectionwith-dengue-and-chikungunya-viruses](http://www.intechopen.com/books/current-topics-in-chikungunya/co-infectionwith-dengue-and-chikungunya-viruses). [Last accessed on 2018 Jan 30].
8. Taraphdar D, Sarkar A, Mukhopadhyay BB, Chatterjee S. A comparative study of clinical features between monotypic and dual infection cases with chikungunya virus and dengue virus in West Bengal, India. *Am J Trop Med Hyg* 2012;86:720-3.

9. Kalawat U, Sharma KK, Reddy SG. Prevalence of dengue and chikungunya fever and their co-infection. *Indian J Pathol Microbiol* 2011;54:844-6.
10. Carey DE, Myers RM, DeRanitz CM, Jadhav M, Reuben R. The 1964 chikungunya epidemic at Vellore, South India, including observations on concurrent dengue. *Trans R Soc Trop Med Hyg* 1969;63:434-45.
11. Afreen N, Deeba F, Khan WH, Haider SH, Kazim SN, Ishrat R, et al. Molecular characterization of dengue and chikungunya virus strains circulating in New Delhi, India. *Microbiol Immunol* 2014;58:688-96.
12. Mohanty I, Dash M, Sahu S, Narasimham MV, Panda P, Padhi S, et al. Seroprevalence of chikungunya in Southern Odisha. *J Family Med Prim Care* 2013;2:33-6.
13. Saswat T, Kumar A, Kumar S, Mamidi P, Muduli S, Debata NK, et al. High rates of co-infection of dengue and chikungunya virus in Odisha and Maharashtra, India during 2013. *Infect Genet Evol* 2015;35:134-41.
14. Kularatne SA, Gihan MC, Weerasinghe SC, Gunasena S. Concurrent outbreaks of chikungunya and dengue fever in Kandy, Sri Lanka, 2006-07: A comparative analysis of clinical and laboratory features. *Postgrad Med J* 2009;85:342-6
15. Krishna A.P, Chandrika, Suchetha, Manasa A and Shrikant L. Patil. Variation in Common lipid parameters in malaria infected Patients. *Indian Journal of Physiology and Pharmacology*, 2009; 53 (3): 271-274.