

# Original Research Article

## Papilionoideae Species Abundance and Diversity in Forest and Outside a Forest Reserve in Taraba State, Nigeria

---

### ABSTRACT

**Aims:** To assess the abundance and diversity of Papilionoideae species in Ngel-Nyaki forest reserve and its surrounding landscape.

**Study design:** The study area was divided into two habitat types; Protected (Ngel-Nyaki forest reserve) and unprotected habitat (surrounding landscape). We identified all species of Papilionoideae in the study area, and we used general linear models to compare the abundance and diversity of Papilionoideae species between the protected habitat and unprotected habitat.

**Place and Duration of Study:** The study was conducted in Ngel-Nyaki forest reserve and surrounding landscape, in Taraba State, South-eastern Nigeria from April to August 2019.

**Methodology:** Thirteen line transects, each measuring 1000m was laid across the study sites, and on each of the 13 line transects, sixteen 10 x 10 m quadrats were established and spaced at an interval of 50m. All Papilionoideae species in our established 10 x 10 m quadrats were identified and recorded. All unidentified Papilionoideae species were collected and transferred to Ngel-Nyaki forest reserve herbarium for identification.

**Results:** We recorded 560 individual species, consisting of 42 species from 6 genera. The protected habitat had the highest abundance and diversity of Papilionoideae species.

**Conclusion:** Overall, we have shown that the Ngel-Nyaki forest reserve and surrounding landscape host many individual Papilionoideae species, suggesting the need for conservation to ensure that these habitats continue to support these species.

*Keywords:* Papilionoideae species, forest, abundance, diversity, surrounding landscape

## 1. INTRODUCTION

Increasing human pressures and corresponding land-use changes are among the greatest threats to tropical biodiversity [1, 2]. It is therefore important to preserve biodiversity in habitats that are impacted by human activities, because these habitats make up an increasingly large portion of the earth [3].

Over the years, government across the globe have invested in establishing protected areas as a fundamental strategy towards biodiversity conservation [4]. The protected areas create habitats for threatened species, support healthy ecosystems and provides ecosystem services to humans (e.g., climate regulation, food security, and clean water) [5]. However, the ability to solely conserve biodiversity within a network of strictly protected areas has become limited by lack of funding, political will, and conservation opportunity [6]. As a result, the protected areas have not totally slowed rates of biodiversity loss. There are numerous gaps in the protected areas networks, especially because these areas do not cover the distribution of many threatened species [5]. Therefore, to develop effective conservation strategies, it may be important to assess biodiversity values both within and outside the protected areas [6, 7]. Moreover, Primack [8] have shown that unique ecosystems and several endangered species are comparatively or solely found within unprotected areas.

In Nigeria, protected areas were originally established around the country in recognition of the significance of many flora and fauna species [9]. However, many of these species are currently threatened or endangered due to lack of proper management and missing data on the status of biodiversity e.g., *Eugenia gilgii* [9-11]. Importantly, some plant species recorded in lowland rainforest outside protected areas in Nigeria are vulnerable, facing a high risk of extinction in the wild, e.g., timber trees *Entandrophragma angolense* and *Khaya grandifoliola* [11]. Therefore, to conserve biodiversity in/for a longtime, it is important to understand the distribution of plant species in and outside protected areas boundaries [12], particularly in Nigeria where there is increasing biodiversity loss [13].

Our study tries to improve the current knowledge on the distribution of a plant species inside and outside Ngel-Nyaki forest reserve, in Taraba State, Nigeria. To this end, we focused on the Fabaceae: Papilionoideae species; a good proxy for understanding plant species distribution patterns in many habitats [14]. Papilionoideae species exhibit wonderful variation in their distribution patterns; from trees, shrubs, lianas, vines [15, 16], and are key stone species in many tropical rain forests because of their ability to provide food and shelter for other animals [17]. However, some species of Papilionoideae are endangered, for example, *Dalbergia melanoxylum* [18]. Therefore, understanding the distribution pattern of these species is a crucial step in management and conservation of plant communities as well as ecosystems.

We compared the abundance and diversity of Papilionoideae species between habitats (protected and unprotected). We predict that Papilionoideae species abundance and diversity will be higher in the Ngel-Nyaki forest reserve as compared to outside the reserve since the reserve is protected against anthropogenic disturbance (e.g., livestock grazing, farming, and cutting of trees for fuelwood).

## 2. MATERIAL AND METHODS

### 2.1 Study area and location of sampling sites

Our study took place in forest and outside the Ngel Nyaki Forest Reserve in Taraba State Nigeria (7.0876°N, 11.0534°E), South Eastern Nigeria [11]. There is a distinct wet and dry season with a mean annual rainfall of around 1700mm. The minimum average monthly temperature is between 15.5-18.5 °C and the maximum ranges from 27.5-30.5 °C [19]. The forest, with an average elevation of ~1650 m asl is characterized by gently undulating hills covered in overgrazed *Sporobolus* grassland. This forest is well known for its rich flora and

fauna, containing several Nigeria's endangered plant species, notable among which are the large forest tree species such as *Etandrophragma angolense*, *Lovoa trichilioides*, *Millettia conraui* and *Pouteria altissima*.

The Ngel Nyaki forest reserve is a vulnerable site of conservation concern because it has been negatively affected by annual bushfires that are usually ignited by Fulani pastoralists [20]. These fires have led to the creation of wide-open grasslands principally consisting of *Sporobolus pyramidalis* and *Hyperhnenia rufa* within the reserve [21]. Outside the reserve, habitat degradation is evident from anthropogenic activities (e.g., livestock grazing, farming and fuelwood), giving way to degraded grazing land and farmlands.

## 2.2 Sampling plots

We sampled Papilionoideae species between April and August 2019 when species can be easily identified. Two hundred and eight study plots, each measuring 10 x 10 m quadrats and spaced at an interval of at least 50 m were established in the forest reserve and those of the surrounding landscapes. We selected areas sampled outside the reserve using the methodology of Abiem, et al [9], with modification. All unidentified Papilionoideae species were collected and transferred to Ngel-Nyaki forest reserve herbarium for identification.

## 2.3 Data analyses

All statistical analyses were carried out using the R statistical package version 3.5.6 (R Core Team 2018). We used general linear models (GLMs) to compare the abundance of Papilionoideae species between the reserve and its surrounding areas. Plant species diversity was calculated using Shannon Wiener's diversity index and GLMs was used to compare species diversity between the reserve and surrounding area. In all cases, statistical test was performed at the significance level of  $p < 0.05$ .

## 3. RESULTS

A total of 560 individual plants of Papilionoideae species consisting of 42 species from 6 genera were recorded during the survey: forest reserve 316 (56%), and surrounding area 244 (44%). See Appendix 1 for checklist of species.

As shown in Table 1, the abundance of Papilionoideae species was higher inside the protected area. General linear model with a Poisson error structure (count data) showed a significant difference between Papilionoideae species abundance ( $p < 0.001$ ) of the protected area and the unprotected area.

The results from the general linear model in Table 2, showed the diversity of Papilionoideae species differed significantly between the protected and unprotected areas, with the protected area having the highest diversity of Papilionoideae species ( $p < 0.001$ ).

**Table 1:** Abundance of Papilionoideae species between forest reserve (protected) and surrounding areas (unprotected)

Parameters	Estimate	SE	T value	P value
Protected habitat	0.75	0.056	13.25	<b>&lt;0.001</b>
Unprotected habitat	0.11	0.085	1.26	0.206

**Note:** Protected habitat was set as the intercept based on alphabetical order. P values in bold show significant difference. Model; abundance = habitat, family = Poisson. Null deviance: 492.17 on 253 degrees of freedom.

**Table 2:** Diversity of Papilionoideae species between protected and unprotected habitats.

Parameters	Estimate	SE	T value	P value
------------	----------	----	---------	---------

Protected habitat	1.63	0.032	50.69	<b>&lt;0.001</b>
Unprotected habitat	0.11	0.050	2.28	<b>0.024</b>

**Note:** Protected habitat was set as the intercept based on alphabetical order. P values in bold show significant difference. Model; diversity = habitat. Null deviance: 40.090 on 253 degrees of freedom.

#### 4. DISCUSSION

The Ngel-Nyaki forest reserve and its surrounding landscape host many individual Papilionoideae species, including *Dalbergia melanoxyllum*, an endangered species in the IUCN red list [18]. This suggests the need to ensure that these habitats continue to support the diversity of these species.

The protection effect was evidenced by high abundance and diversity of Papilionoideae species in the protected habitat compared to the unprotected habitat, indicating that protected area plays important role in guaranteeing the conservation of biodiversity. On the other hand, the unprotected areas may have experienced anthropogenic interference from livestock grazing, farming, cutting of trees for firewood, and access road, making the soil unsuitable for the growth of the plants. Another possible explanation for the lower diversity in the surrounding landscape could be the extensive collection of seeds, fruits, and roots by the locals from some of the Papilionoideae species may have possibly affect the distribution, and natural regeneration of the plant species in the surrounding areas, especially for species that could not adapt to the change in the environmental condition [22]. Therefore, there is need for a broad spectrum conservation [9].

The soil composition within protected areas is rich in humus because of the minimal anthropogenic activities, when compared to surrounding areas [23]. This may support the growth of vigorous plants species, an indication of ecological healthy ecosystem. Perhaps, this is true because we recorded a good number of the Papilionoideae family in the protected areas only. E.g., *Abrus cicimanga*, *Adenocarpus celerata*, *Alysicarpus rugosus*, *Calopogonium muconoides*, *Crotalaria retusa*, *Dalbergia afzeliana*, *Dalbergia melanoxyllum*, *Dalbergia nitoduna*, and *Dalbergia vogelii*. However, it is important to note that, at the time of this study, even within the protected habitat, anthropogenic activities e.g., illegal forest fire still occurs, although at a minimal level compared to the surrounding areas. If this persists, it could lead to habitat degradation of the protected habitat, which may result to loss of the Papilionoideae species as well as other plant species. Hence, the need for humans, policy makers and the management of the forest reserve to ensure the protection of this habitat.

#### CONCLUSION

Overall, the results of this study have direct implications for conservation. The high abundance and diversity of Papilionoideae species in the reserve when compared to the surrounding landscape can be attributed to the differences in environmental conditions and the ability of the species to adapt to these conditions [24]. However, conservation of the surrounding landscape is as important as the protected reserve because we also recorded some Papilionoideae species outside the reserve. This surrounding landscape could act as either population sinks for some fauna and/or valuable corridors increasing connectivity [25-27].

#### ETHICAL APPROVAL

Not applicable. This research does not require an ethics approval.

## REFERENCES

1. Laurance WF, Camargo JL, Fearnside PM, Lovejoy TE, Williamson GB, Mesquita RC, et al. An Amazonian rainforest and its fragments as a laboratory of global change. *Biological Reviews*. 2018;93(1):223-47.
2. Gibson L, Lee TM, Koh LP, Brook BW, Gardner TA, Barlow J, et al. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature*. 2011;478(7369):378-81.
3. Buechley ER, Şekercioğlu ÇH, Atickem A, Gebremichael G, Ndungu JK, Mahamued BA, et al. Importance of Ethiopian shade coffee farms for forest bird conservation. *Biological Conservation*. 2015;188:50-60.
4. Lopoukhine N, Crawhall N, Dudley N, Figgis P, Karibuhoye C, Laffoley D, et al. Protected areas: providing natural solutions to 21st Century challenges. *SAPI EN S Surveys and Perspectives Integrating Environment and Society*. 2012;(5.2).
5. Sritharan S, Burgess ND. Protected area gap analysis of important bird areas in Tanzania. *African Journal of Ecology*. 2012;50(1):66-76.
6. Gardner TA, Caro T, Fitzherbert EB, Banda T, Lalbhai P. Conservation value of multiple-use areas in East Africa. *Conservation Biology*. 2007;21(6):1516-25.
7. Vandermeer J, Perfecto I. The agricultural matrix and a future paradigm for conservation. *Conservation biology*. 2007;21(1):274-7.
8. Primack R. Conservation outside protected areas. *A Primer of Conservation Biology*. 2012;255-81.
9. Abiem I, Saha S, Manu SA, Elisha EB. Woody vegetation composition and diversity in woodlands inside and outside a Forest Reserve in Jos, Nigeria. *African Journal of Ecology*. 2018;56(2):202-7.
10. Usman B, Adefalu L. Nigerian forestry, wildlife and protected areas: Status report. *Biodiversity*. 2010;11(3-4):54-62.
11. Chapman J, Chapman H. The forests of Taraba and Adamawa States, Nigeria. An ecological account and plant species checklist University of Canterbury, Christchurch, New Zealand. 2001.
12. Zhang J-T, Xu B, Li M. Vegetation patterns and species diversity along elevational and disturbance gradients in the Baihua Mountain Reserve, Beijing, China. *Mountain Research and Development*. 2013;33(2):170-8.
13. Tela M. The value of forest fragments in maintaining ecosystems for food security in Sub-Saharan Africa. . PhD Thesis University of Canterbury, New Zealand. 2019.
14. Yahara T, Javadi F, Onoda Y, de Queiroz LP, Faith DP, Prado DE, et al. Global legume diversity assessment: concepts, key indicators, and strategies. *Taxon*. 2013;62(2):249-66.
15. Tapia-Pastrana F, Delgado-Salinas A, Caballero J. Patterns of chromosomal variation in Mexican species of *Aeschynomene* (Fabaceae, Papilionoideae) and their evolutionary and taxonomic implications. *Comparative cytogenetics*. 2020;14(1):157.
16. Arbonnier M. Trees, shrubs and lianas of West African dry zones: Quae; 2004.
17. Schulze E-D, Mooney HA. Biodiversity and ecosystem function: Springer Science & Business Media; 2012.

18. Yin X, Huang A, Zhang S, Liu R, Ma F. Identification of three *Dalbergia* species based on differences in extractive components. *Molecules*. 2018;23(9):2163.
19. Matthesius A, Chapman H, Kelly D. Testing for Janzen–Connell effects in a west African montane forest. *Biotropica*. 2011;43(1):77-83.
20. Yadok BG. Aspects of the behaviour of African giant pouched rats (*Cricetomys* sp. nov) which impacts seed dispersal of large-seeded tree species in a West African montane forest landscape. 2018.
21. Barnes AD, Chapman HM. Dispersal traits determine passive restoration trajectory of a Nigerian montane forest. *Acta Oecologica*. 2014;56:32-40.
22. Bandyopadhyay S. A Systematic Census on the Sedges of Koch Bihar district, West Bengal. *Indian Journal of Applied and Pure Biology*. 2017;32:181-8.
23. Fenu G, Cogoni D, Ulian T, Bacchetta G. The impact of human trampling on a threatened coastal Mediterranean plant: The case of *Anchusa littorea* Moris (Boraginaceae). *Flora-Morphology, Distribution, Functional Ecology of Plants*. 2013;208(2):104-10.
24. Bibby C, Burgess N. Hill, DA & Mustoe, SH 2000. Bird census techniques. London: Academic Series; 2000.
25. Neuschulz EL, Botzat A, Farwig N. Effects of forest modification on bird community composition and seed removal in a heterogeneous landscape in South Africa. *Oikos*. 2011;120(9):1371-9.
26. MacGregor- Fors I, Schondube JE. Use of tropical dry forests and agricultural areas by Neotropical bird communities. *Biotropica*. 2011;43(3):365-70.
27. Tela M. The value of forest fragments in maintaining ecosystems for food security in Sub-Saharan Africa: University of Canterbury; 2019.

## APPENDIX

Appendix 1: Checklist of Papilionoideae species identified in forest and outside Ngel-Nyaki Forest Reserve.

S/N	Common name	Species name
1	Rosary pea	<i>Abrus precatorius</i> L.
2	Cosedo	<i>Adenocarpus celerata</i> ** Recheck the specific epithet
3	American jointvetch	<i>Aeschynomene americana</i> L.
4	Budda pea	<i>Aeschynomene indica</i> L.
5	Red moneywort	<i>Alysicarpus rugosus</i> (Willd.) DC.
6	Pigeon pea	<i>Cajanus cajan</i> (L.) Huth.
7	Calopo	<i>Calopogonium muconoides</i> Desv.
8	Showy pea	<i>Campylotropis speciosa</i> (Royle ex Schindl.) Schindl.
9	Devil bean	<i>Crotalaria goreensis</i> Guill. & Perr.
10	Rattle weed	<i>Crotalaria retusa</i> L.
11	Smooth rattlebox	<i>Crotalaria striata</i> DC. This is a synonym of <i>Crotalaria pallida</i> var. <i>pallida</i> Or <i>Crotalaria striata</i> A. Braun This is a synonym of <i>Crotalaria impressa</i> Nees ex Walp.

---

		***which one is your plant. Confirm it.
12	Rose wood	<i>Dalbergia afzeliana</i> G. Don
13	African blackwood	<i>Dalbergia melanoxydon</i> Guill. & Perr.
14	indian rosewood	<i>Dalbergia nitoduna</i> / <i>Dalbergia nitidula</i> Welw. ex Baker
		**Check the spelling
15	Diola flup	<i>Dalbergia rufa</i> G. Don
16	North Indian rosewood	<i>Dalbergia sissoo</i> Roxb. ex DC.
17	Rose wood	<i>Dalbergia</i> spp
18	Sarivan	<i>Desmodium cordifolia</i> is a synonym of <i>Grona cordifolia</i> (Harms) H. Ohashi & K. Ohashi
19	Salparni	<i>Desmodium gangeticum</i> is a synonym of <i>Pleurolobus gangeticus</i> (L.) J. St.-Hil. ex H. Ohashi & K. Ohashi
20	Roundhead tickclover	<i>Desmodium rotundifolium</i> DC.
21	Scorpion tick trefoil	<i>Desmodium scorpiurus</i> (Sw.) Desv. ex DC.
22	Creeping tick trefoil	<i>Desmodium triflorum</i> (L.) DC. This is a synonym of <i>Grona triflora</i> (L.) H. Ohashi & K. Ohashi
23	Silverleaf desmodium	<i>Desmodium uncinatum</i> (Jacq.) DC.
24	Tick clover	<i>Desmodium velutinum</i> (Willd.) DC. This is a synonym of <i>Polhillides velutina</i> (Willd.) H. Ohashi & K. Ohashi
25	Gallito	<i>Erythrina vogelii</i> Hook.f.
26	Silky indigo	<i>Indigofera astragalina</i> DC.
27	Hairy indigo	<i>Indigofera hirta</i> Bojer This is a synonym of <i>Indigofera hirsuta</i> L.
28	Indigo	<i>Indigofera microphylla</i> Lam. This is a synonym of <i>Indigofera sarmentosa</i> L.f.
29	Western indigo	<i>Indigofera miniata</i> Ortega
30	True indigo	<i>Indigofera pseudotinctoria</i> Matsum. This is a synonym of <i>Indigofera bungeana</i> Walp.
31	Three leaf indigo	<i>Indigofera trifoliata</i> L.
32	Velvet bean	<i>Mucuna puggie</i> / <i>Mucuna poggei</i> Taub.
34	Caribbean stylo	<i>Stylosanthes hamata</i> (L.) Taub.
35	Hoary pea	<i>Tephrosia cajanus</i> *** Confirm the species
36	Poison pea	<i>Tephrosia vicioides</i> A. Rich. This is a synonym of <i>Tephrosia uniflora</i> subsp. <i>uniflora</i> / <i>Tephrosia vicioides</i> Schldl.
37	Fishpoison bean	<i>Tephrosia vogelii</i> Hook.f.
38	Hairy pod cowpea	<i>Vigna luteola</i> (Jacq.) Benth.
39	Zombi pea	<i>Vigna vexillata</i> (L.) A. Rich.
40	Banmoong	<i>Vigna wightii</i> Benth. ex Bedd. This is a synonym of <i>Vigna vexillata</i> var. <i>wightii</i>

---

---

		(Benth. ex Bedd.) Babu & S.K. Sharma
41	Two-leaf zornia	<i>Zornia latifolia</i> Sm.
42	Dogon	<i>Zornia</i> spp

---

UNDER PEER REVIEW