

## Original Research Article

# A Comparative Study of Hair Morphology of three Carnivore families (Canidae, Ursidae and Herpestidae)

Comment [S1]: Descriptive Study

### ABSTRACT

**Aim:** To comparatively study the morphological characteristics of dorsal guard hair of four carnivoran species native to Tamil Nadu for the creation of a reference repository for wildlife forensics.

**Study design:** The experiment was designed in a way to observe the physical characters of each hair strand and measure widely used morphological parameters from native species of the state. The results were compared with previous findings to validate the differences and similarities with our study.

**Place and Duration of Study:** Advanced Institute for Wildlife Conservation (AIWC), Tamil Nadu Forest Department, Vandalur, Chennai, Tamil Nadu. between August 2020 and September 2021.

**Methodology:** Hair samples were collected for Golden Jackal (*Canis aureus*), Indian Fox (*Vulpes bengalensis*), Dhole (*Cuon alpinus*), Sloth bear (*Melursus ursinus*) and Indian grey mongoose (*Herpestes edwardsii*). Hair characters such as scale pattern, scale distance and scale margin, medulla structure, cortex thickness, medulla thickness, medullary index and cross-sectional structure were observed.

**Results:** The hair length of golden jackal, Indian fox, dhole, sloth bear and Indian grey mongoose were as follows  $44.94 \pm 2.24$  mm,  $29.70 \pm 1.51$  mm,  $29.60 \pm 0.97$  mm,  $83.83 \pm 2.93$  mm,  $54.60 \pm 2.03$ . The cuticular characters showed variation but overall, the characters were in line with previous studies. The medullary structure was wide medulla with vacuoles for all three canid species while sloth bear has a narrow simple unbroken medulla and Indian grey mongoose had wide medulla with lattice. The medullary index for golden jackal, Indian fox, dhole, sloth bear and Indian grey mongoose were  $0.73 \pm 0.05$  (S.D.),  $0.77 \pm 0.071$  (S.D.),  $0.66 \pm 0.06$  (S.D.),  $0.18 \pm 0.08$  (S.D.) and  $0.63 \pm 0.05$  (S.D.) respectively.

**Conclusion:** The morphometric characters of hair showed variation in color, scale pattern, scale margin, scale distance and medullary index when compared with previous studies which may be attributed to environmental factors.

**Keywords:** Tricho-taxonomy, hair morphometry, carnivores, species identification, reference repository

## 1. INTRODUCTION

India is home to 427 species of terrestrial and marine mammals (1) that face survival threat from humans in the form of widespread illegal hunting and poaching (2). Mammals constitute a major share in international illegal wildlife trade mainly for their skin, body parts and meat, and live animals (3) and in most cases hair ends up as only physical evidence for examination (4). Species identification from hair characteristics has become an integral part of wildlife forensics and study of diet ecology of carnivores because the properties of the hair do not get affected by the digestive process and takes more time to decay (5).

The mammalian skin is covered by guard hair or protective hair and fur hair or under-hair (6). For species identification, dorsal guard hair is generally used because hair from other regions of the body lack clear morphological properties (7). Microscopic and macroscopic characteristics of wild mammalian hair was pioneered by Hausman (6) and later refined (8). In India, tricho-taxonomical research on various families of mammals was expansively carried out by Chakraborty and De (9, 10, 11), De et al (12, 13, 14 & 15), Sahajpal et al (4 & 16), Bahuguna (17) and in recent past decade by Kamalakannan et al (18, 19, 20 & 21).

The present work is intended to compare morphological characteristics of hair with previous findings of five species of carnivores, Golden jackal (*Canis aureus*), Indian Fox (*Vulpes bengalensis*), Dhole (*Cuon alpinus*) (Family: Canidae), Sloth bear (*Melursus ursinus*) (Family: Ursinidae) and Indian Grey Mongoose (*Herpestes edwardsii*) (Family: Herpestidae), which are overlooked species in conservation point of view. The golden jackal is hunted in many parts of India driven by superstitious beliefs (22). The Indian fox is threatened by loss of habitat throughout its native range (23) and in some parts of south India they are captured and ceremonially released during the Sankranti festival (24). The sloth bear is poached for their meat, skin, teeth, claws, bile gland and derivatives used as traditional medicine in local and international markets (2). The Indian grey mongoose and other species of mongooses had been under constant threat due to poaching for their fine hair quality, this characteristic feature has high demand in the making of painting and shaving brushes (4).

Among the five, the three canid species and Indian grey mongoose belong to schedule II and sloth bear belong to schedule I of Wildlife (Protection) Act (1972). These species are often seized by forest department officials in the state of Tamil Nadu and this work was exclusively designed with the aim of creating a standard reference morphometric data for identification of unknown hair samples from the field.

**The study ignored the individual variations within species.**

## 2. MATERIAL AND METHODS

### 2.1 Sample Collection

Naturally fallen hair from captive Golden Jackal (*Canis aureus*), Indian Fox (*Vulpes bengalensis*), Dhole (*Cuon alpinus*), Sloth bear (*Melursus ursinus*) were collected from their enclosures at Aringar Anna Zoological Park, Vandalur and Kurumbapatti Zoological Park, Salem, between 2019 and 2021. For Indian grey mongoose (*Herpestes edwardsii*), hair sample was obtained from a dead specimen from Vandalur reserve forest.

### 2.2 Sample Preparation

Pre-treatment included washing hair samples with tap water followed by acetone treatment to remove exogenic impurities from the hair. The cleansed hair was then dried and used for further observation (20). A total of thirty (n=30) hair samples from each species (150 hair for all five species) was used for morphometric analysis.

### 2.3 Morphometric analysis

Physical characteristics such as colour of the hair, banding, length of each strand, root length was observed. Cuticular characters such as cuticular width, scale pattern, scale margin, scale distance was observed for each hair strand at proximal, medial and distal ends of the hair (7).

For observation of cuticular characters, cuticle imprints of the hair were made on a 1% - 2% gelatine smear on a clear glass slide (16). Medullary characters such as type of medulla, medullary width, medullary index and percentage along with cross-section were observed for each hair strand by soaking the hair pieces in xylene (14).

All the images were taken at 400x magnification except for the root of sloth bear which was taken at 100x as they have the longest root observed among the five species. Hair length was

**Comment [S2]:** describe

**Comment [S3]:** please, write the replication ( number of animals) within each species hair used in the test

**Comment [S4]:** The samples from one animal?

measured from digital photograph of hair using ImageJ® software (26), while other microscopic measurements were carried out via Magnus CX23 microscope with Magvision® software. The hair characteristics is described using the Nomenclature (6 & 8).

There are many environmental factors besides genetics affect and also genetic-environmental interaction effects on hair status and structure, the environmental factors such as : nutrition, stress, age, pregnancy, season and weather change, so the author must describe the these factors as a fixed model not effected on the results

### 3. RESULTS

#### 3.1 Physical characters

The hair colour and texture of Golden Jackal (GJ) had brown or umber colour hair with dark tips at distal end with a smooth texture. The Indian Fox (IF) had soft hair with grey to buff colour depending on the part of the body. The Dhole (D) had a distinct brown to orange colour with dark distal ends and a rough texture to its hair. The Sloth Bear (SB) had long black human-like hair with kinks and a very rough texture. The Indian Grey Mongoose (IGM) had brown tips followed by alternating light and dark bands with smooth texture to its hair. The macroscopic appearance of hair of all the species is giving in the figure (Figure 1).



Figure 1. Macroscopic appearance of hair

The structure and length of hair root (Table 1) were observed for providing additional information of hair samples. The root of GJ and F had similar appearing root but the later had comparatively shorter root. The D had much longer root than the other two canid species. The SB had the longest root among the five species that tapers to the tip and the IGM had an oblong root (Figure 2). The morphometric data and physical characteristics of hair are given in Table 1.




Figure 2. Root structure of different species

Table 1. Physical characteristics of hair of the species under study

Features	Golden Jackal	Indian Fox	Dhole	Sloth Bear	Indian grey mongoose
Colour	Umber dark tips with	Grey to buff	Orange brown to	Black	Brown tips with alternate white and brown bands
Shape	Straight	Straight	Short Straight	Long with kinks	Straight
Texture	Rough	Smooth	Rough	Very rough	Smooth
Hair Length	44.94 ± 2.24 mm (S.E.)	29.70 ± 1.51 mm (S.E.)	29.60 ± 0.97 mm (S.E.)	83.83 ± 2.93 mm (S.E.)	54.60 ± 2.03 mm (S.E.)
Root length	585.21 ± 11.96 μm (S.E.)*	319.54 ± 10.38 μm (S.E.)*	710.31 ± 19.10 μm (S.E.)*	1235.58 ± 36.63 μm (S.E.)*	419.22 ± 15.68 μm (S.E.)*

\*S.E. - Standard Error

### 3.2 Cuticular characters

The cuticular characters varied along the length of each hair. Hence, it was documented in proximal, medial and distal parts of the hair. The cuticular scales are transversal in nature at distal and medial parts of the hair samples. The scale margin at proximal end was observed to be smooth for all the species. In the medial region the scale margin of GJ, D and SB were crenate type, while IF had a mixture of smooth and crenate type and the IGM had a combination of crenate and rippled margin type. The distal part of the hair of all species had crenate type margin.


The scale distance was observed to be distant at the proximal end for GJ, IF, D and SB while the Indian grey mongoose had nearer scales. At medial region, scales were nearer in GJ, D, SB and IGM but the IF has distant scales. Similarly, the scales at distal region were nearer in GJ, IF and D while SB and IGM had both nearer and cuticular scales.

The scale pattern of GJ had mixture of diamond petal and regular wave in the proximal end of their hair. The IF had broad diamond petal in proximal end. The D hair had regular wave pattern in most of the hair with broad petal in a few samples. The scale pattern of SB and IGM had regular wave pattern in their proximal ends. The scale pattern at medial and distal end of GJ, D, SB and IGM had irregular wave type pattern with few outliers. The IF showed diamond petal scale pattern in medial and distal end of its hair (refer Table 2 and Figure 3).

**Table 2. Cuticular characteristics comprising scale margin, scale distance and scale pattern**

Cuticular characteristics	Golden Jackal	Indian Fox	Dhole	Sloth Bear	Indian grey mongoose
Scale margin	Basal (proximal)	Smooth	Smooth	Smooth	Smooth
	Medial	Crenate	Smooth to Crenate	Crenate	Crenate
	Apical (distal)	Crenate	Crenate	Crenate	Crenate
Scale distance	Basal (proximal)	Distant	Distant	Distant	Distant
	Medial	Near (Intermediate)	Distant	Near (Intermediate)	Near (Intermediate)
	Apical (distal)	Near (Intermediate)	Near (Intermediate)	Near (Intermediate)	Close

<b>Scale pattern</b>	<b>Basal (proximal)</b>	Broad Diamond Petal to Regular wave	Broad diamond petal	Broad Diamond Petal to Regular wave	Regular wave	Regular wave
	<b>Medial</b>	Irregular wave	Broad Diamond Petal to regular wave	Irregular wave	Irregular wave	Irregular wave
	<b>Apical (distal)</b>	Irregular wave	Irregular wave	Irregular wave	Irregular wave	Irregular wave



**Figure 3. Cuticular characteristics of hair at distal, medial and proximal regions**


### 3.3 Medullary characters

In our study it was observed that GJ and IF had wide medulla lattice with vacuoles ( $> 0.70$  medullary index). A similar structure was observed in D but the thickness of medulla was much less with vacuoles ( $< 0.70$  medullary index). The SB hair had a simple narrow continuous medulla ( $< 0.20$  medullary index) and the IGM hair had a wide medulla ( $> 0.60$  medullary index) with intrusion of cortex. Cross section of GJ, IF and D exhibited circular to oval shape with pigments around the medulla. SB hair exhibited oval shape with heavy pigmentation in the cortex. The IGM exhibited oval to oblong shaped hair (Figure 4). The medullary index values and medulla percentage for all the five species is given in the table (Table 3).

**Table 3. Medullary characteristics of hair**

Medullary characteristics	Golden Jackal	Indian Fox	Dhole	Sloth Bear	Indian grey mongoose
Medulla type	Wide medulla with vacuoles	Wide medulla with vacuoles	Wide medulla with vacuoles	Narrow unbroken medulla	Wide medulla lattice
Medullary index	0.73 ± 0.05 (S.D.)	0.77 ± 0.071 (S.D.)	0.66 ± 0.06 (S.D.)	0.18 ± 0.08 (S.D.)	0.63 ± 0.05 (S.D.)
Medullary percentage	73.44%	76.91 %	66.66%	18.52%	63.99%
Cross-section type	Circular	Circular	Circular	Oval to oblong	Oval to oblong

S.D. Standard Deviation



**Figure 4. Medullary structure and cross-section of hair**

A graph (Figure 5) was plotted with average medullary index against the cortical thickness of five species of carnivores. This simple method is useful to understand the relationship between hair thickness (width) and medulla thickness (width) and can be applied to unknown hair samples to find their relatedness to standard values.




Figure 5. Cuticle thickness and medullary index of hair of carnivores under study

#### 4. DISCUSSION

A cross reference with previously published literature for all five species provided the similarities and differences in hair characteristics present in our current research.

##### 4.1 Golden jackal

The hair length of GJ from our observation  $44.94 \pm 2.24$  (S.E.) mm was similar to the one observed by Chakraborty and De (2001)  $46.76 \pm 13.14$  (S.E.) mm but Arpacik (2021) reported an average length of  $6.66 \pm 1.22$  (S.E.) cm from the GJ of Turkey. There was a difference in the cuticular pattern of Golden Jackal hair, but the scale margin is same as observed by (7, 10 & 17). Bahuguna reported broad diamond petal scale pattern at proximal end of the hair (17) while our observation showed both regular wave and broad diamond petal pattern. The medullary index had variation between  $0.67 \pm 0.02$  S.D. (10),  $0.72 \pm 0.005$  S.D. (7) and our observation  $0.73 \pm 0.05$  S.D.

##### 4.2 Indian Fox

The IF hair characteristics observed were mostly in line with the findings in (10). The hair length was reported to be  $26.6 \pm 2.9$  (S.E.) mm with smooth scale margin in the proximal end and irregular wave and diamond petal scale pattern along the hair length. The medulla is vacuolated with a medullary index of  $0.81 \pm 0.012$  (S.D.) and is circular in shape.

##### 4.3 Dhole

The hair length of dhole was reported to have  $44.2 \pm 3.6$  (S.E.) mm (10) while we observed only  $29.60 \pm 0.97$  (S.E.) mm. The cuticle pattern is described to be of regular wave type at the proximal end (17) and irregular wave type of hair (7) but our observation had regular wave with some broad diamond petal pattern at proximal end and irregular wave pattern along other parts of the hair. A much lower medullary index  $0.59 \pm 0.02$  S.D. was observed in (7) compared to our result.

##### 4.4 Sloth bear

The average hair length,  $88.83 \pm 2.93$  (S.E.) mm was comparatively greater than the previous observation of  $69.44 \pm 3.02$  (S.E.) mm (14). The scale pattern at proximal end was reported to have regular mosaic wave (14) and broad petal (17) though we observed only regular wave pattern. The medullary index was  $0.91 \pm 0.008$  S.D. (14) while our observation was  $0.18 \pm 0.08$  S.D. The cross-section type was oval to oblong (14) but an oval cross-section is reported in (17). Our results were inclined to (14).

#### 4.5 Indian grey mongoose

The average hair length of *Herpestes edwardsii* was observed to be  $25.61 \pm 3.34$  (S.E.) mm (13). Medullary index was at  $0.82 \pm 0.003$  (S.D.) (13) and  $0.792 \pm 0.005$  (S.D.) (4) while our observation had an average medullary index of  $0.63 \pm 0.05$  (S.D.). Our cross-section shape was concurrent to previous record (4).

#### 5. CONCLUSION

The hair characteristics of same species exhibit variation among different individuals. Such variations may be attributed to geographical differences, environmental factors, sex, and age of the species (7 & 25). The species belonging to the same family share some similarities in hair morphology in canids (7, 10), in ursids (14, 16) and in herpestids (4, 13) of India. In our case, we found similarity in cuticle patterns, medulla structure and cross-section shapes of Golden Jackal, Dhole and Indian Fox.

This research reiterates the established statement about tricho-taxonomy that it takes a combination of different cuticular and medullary characteristics to identify a species from hair. The inclusion of the root length and medullary index graph to ascertain unknown hair samples can provide valuable information when dealing with unknown hair samples. A robust method could be achieved by the application of statistical methods for studying variations within and between species. Nevertheless, the present method has been efficient in species identification alongside creating a reference standard for mammalian hair is essential for wildlife forensic investigatory groups respective to their geographical ranges.

**Comment [S5]:** The study ignored these factors and not describe the experimental units according to these factors

#### ETHICAL APPROVAL

The present work does not involve any kind of experimentation on live animals. As the study involved non-invasive biological sampling of mammalian hair, which are generally considered as animal waste, permissions from ethical committees on animal experimentation and conservation are not required.

#### REFERENCES

1. Sharma G, Kamalakannan M, Venkataraman K. A checklist of mammals of India with their distribution and conservation status. *Zool Surv India*. 2015;111.
2. Gomez L, Wright B, Shepherd CR, Joseph T. An analysis of the illegal bear trade in India. *Global Ecology and Conservation*. 2021;27, e01552.
3. Nijman V. An overview of international wildlife trade from Southeast Asia. *Biodiversity and conservation*. 2010;19(4):1101-1114.
4. Sahajpal V, Goyal SP, Raza R, Jayapal R. Identification of mongoose (genus: *Herpestes*) species from hair through band pattern studies using discriminate functional analysis (DFA) and microscopic examination. *Science & Justice*. 2009;49(3):205-209.
5. Joshi HR, Gaikwad SA, Tomar MPS, Shrivastava K. Comparative trichology of common wild herbivores of India. *Advances in Applied Science Research*. 2012;36:3455-8.
6. Hausman LA. Structural characteristics of the hair of mammals. *The American Naturalist*. 1924;54(635):496-523.
7. Arpacık A. Microanatomical Observations of Hair Characteristics of Red Fox (*Vulpes vulpes*), Golden Jackal (*Canis aureus*), and Gray Wolf (*Canis lupus*): A Comparative Study. 2021.
8. Brunner H, Coman B. The identification of mammalian hair Shanghai Printing Press Ltd. Hong Kong. 1974.
9. Chakraborty R, De JK. Structure and Pattern of Cuticular Scales on Mid-Dorsal Guard Hairs of Marbled Cat, *Felis marmorata charltoni* Gray (Mammalia: Carnivoro: Felidae). *Records of the Zoological Survey of India*. 1996;95(1-2):65-70.
10. Chakraborty R, De JK. Identification of dorsal guard hairs of five Indian species of the family Canidae (Carnivora: Mammalia). 2001.
11. Chakraborty R, De JK. Identification of dorsal guard hairs of nine Indian species of the family Viverridae (Carnivora: Mammalia). *Records of the Zoological Survey of India*. 2005;104(3-4):13-21.
12. De JK. Study of surface structure of hair of some primates of Indian sub-continent. *Records of the Zoological Survey of India*. 1993;93(1-2):31-34.

13. De JK, Chakraborty S, Chakraborty R. Identification of dorsal guard hairs of five Indian species of mongoose, *Herpestes Illiger* (Mammalia: Carnivora). 1998.
14. De JK, Chakraborty R. Identification of dorsal guard hairs of four Indian species of Bear (Mammalia: Carnivora: Ursidae). *Records of Zoological Survey India*. 2006;106(3):19-26.
15. De JK, Chakraborty R. Identification of dorsal guard hairs of nine species of the family Bovidae (Artiodactyla: Mammalia). *Records of the Zoological Survey of India-A Journal of Indian Zoology*. 2012;112(2):39-52.
16. Sahajpal V, Goyal SP, Jayapal R, Yoganand K, Thakar MK. Hair characteristics of four Indian bear species. *Science & Justice*. 2008;48(1):8-15.
17. Bahuguna A. Species identification from guard hair of selected Indian mammals: a reference guide. *Wildlife Institute of India*. 2010.
18. Kamalakannan M, De JK, Manna CK. Identification of dorsal guard hairs surface structure of Indian chevrotain *Moschiola Indica* Gray, 1852 (Tragulidae: Artiodactyla: Mammalia). 2013.
19. Kamalakannan M, De JK, Manna CK. Tricho-taxonomic studies of dorsal guard hairs of Indian Cervids (Artiodactyla: Mammalia) for species identification. 2015.
20. Kamalakannan M, De JK. Hair Morphology of Striped Hyena *Hyaena hyaena* (Linnaeus, 1758). *Int. J. Curr. Microbiol. App. Sci*. 2017;6(5):1438-1441.
21. Kamalakannan M, Chandra K, De JK, Venkatraman C. The identification of pika and hare through tricho-taxonomy (Mammalia: Lagomorpha). *Journal of Threatened Taxa*. 2019;11(10):14301-14308.
22. Chawla MM, Srivathsa A, Singh P, Majgaonkar I, Sharma S, Punjabi G, Banerjee A. Do wildlife crimes against less charismatic species go unnoticed? A case study of Golden Jackal *Canis aureus* Linnaeus, 1758 poaching and trade in India. *Journal of Threatened Taxa*. 2020;12:15407-15413.
23. Sharma G. A review on the Studies on Faunal diversity, status, threats and conservation of Thar Desert or Great Indian Desert Ecosystem. In *Biological Forum—An International Journal*. 2013;5(2):81-90.
24. Gommer ME, Vanak AT. *Vulpes bengalensis*. *Mammalian Species*, 2006;(795):1-5.
25. Sarma KK, Bhattacharjee PC, Dey S. Microscopical analysis of guard hair of Tiger, *Panthera tigris*, with reference to wildlife forensic applications. *Journal of Advanced microscopy research*. 2014;9(3):199-205.
26. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*. 2012;9(7):671–675. doi:10.1038/nmeth.2089

**APPENDIX**

**GOLDEN JACKAL (*Canis aureus*)**

S. No.	Hair length (mm)	Root length (µm)	Cortical thickness(µm)	Medullary thickness (µm)	Medulla Percentage in hair (%)	Medullary index
1	59.16	583.45	98.18	72.44	73.783	0.738
2	65.48	640.7	118.77	91.71	77.216	0.772
3	58.98	578.61	86.06	59.93	69.637	0.696
4	73.13	703.03	95.77	64.189	67.024	0.670
5	44.32	526	94.57	56.59	59.839	0.598
6	49.92	698.05	158.97	118.66	74.643	0.746
7	29.87	614.02	98.44	71.79	72.928	0.729
8	46.36	461.47	73.34	49.93	68.080	0.681
9	56.72	696.08	189.96	136.89	72.063	0.721
10	48.69	601.95	139.2	104.49	75.065	0.751
11	49.95	572.72	145.68	103.15	70.806	0.708
12	51.8	542.63	142.86	96.61	67.626	0.676
13	52.81	509.93	117.51	84.2	71.653	0.717
14	32.33	573	159.48	110.24	69.125	0.691
15	41.81	451.69	97.21	74.82	76.967	0.770
16	37.51	534.23	163.32	127.98	78.361	0.784
17	27.9	524.43	148.12	116.23	78.470	0.785
18	29.85	562.97	146.8	111.51	75.960	0.760
19	29.56	632.11	148.07	118.87	80.280	0.803
20	38.58	590.47	167.13	130.26	77.939	0.779
21	37.76	518.56	136.57	106.79	78.194	0.782
22	56.25	555.53	122.52	80.91	66.038	0.660
23	22.76	580.42	147.08	97.08	66.005	0.660
24	35.29	591.59	166.02	133.83	80.611	0.806
25	56.72	705.27	148.84	104.53	70.230	0.702
26	41.42	624.89	170.79	134.61	78.816	0.788
27	34.78	610.61	130.63	96.64	73.980	0.740
28	53	651.32	189.19	141.6	74.845	0.748
29	50.7	573.6	163.25	124.66	76.361	0.764
30	35.05	546.99	138.56	111.64	80.572	0.806
<b>Mean</b>	<b>44.949 ± 2.247 (Standard error)</b>	<b>585.211 ± 11.96 (Standard error)</b>	<b>136.763 ± 30.613 (Standard deviation)</b>	<b>101.093 ± 25.814 (Standard deviation)</b>	<b>73.437 ± 5.162 (Standard deviation)</b>	<b>0.734 ± 0.052 (Standard deviation)</b>

**INDIAN FOX (*Vulpes bengalensis*)**

S. No.	Hair length (mm)	Root length (µm)	Cortical thickness(µm)	Medullary thickness (µm)	Medulla Percentage in hair (%)	Medullary index
1	23.06	399.6	80.04	62.01	77.474	0.775
2	19.71	403.45	66.13	44.4	67.140	0.671
3	25.61	298.35	49.43	33.28	67.328	0.673
4	46.07	352.37	53.62	34.16	63.708	0.637
5	22.73	377.82	95.72	77.08	80.527	0.805
6	44.04	363.43	135.81	120.19	88.499	0.885
7	31.19	346.17	89.63	64.91	72.420	0.724
8	49.08	392.79	138.48	113.18	81.730	0.817
9	28.77	374.77	108.68	86.71	79.785	0.798
10	26.9	337.28	132.47	104.39	78.803	0.788
11	27.59	270.86	81.47	62	76.102	0.761
12	25.36	233.58	147.05	127.66	86.814	0.868
13	29.47	315.86	74.31	58	78.051	0.781
14	29.54	314.57	103.86	84.48	81.340	0.813
15	25.76	388.57	82.43	67.45	81.827	0.818
16	24.87	304.78	75.39	55.09	73.073	0.731
17	31.67	382.13	54.96	43.8	79.694	0.797
18	26.59	327.28	148.47	124.490	83.849	0.838
19	24.57	285.92	78.24	56.3	71.958	0.720
20	30.29	275.26	135.01	107.04	79.283	0.793
21	36.21	299.4	71.46	54.82	76.714	0.767
22	31.3	206.86	118.89	97.23	81.781	0.818
23	34.04	252.12	103.52	82.56	79.753	0.798
24	30.09	339.1	119.17	100.86	84.635	0.846
25	19.79	221.18	57.37	36.05	62.838	0.628
26	20.45	296.34	83.3	67.3	80.792	0.808
27	31.46	381.52	60	38.36	63.933	0.639
28	26.24	261.04	118.42	98.97	83.575	0.836
29	18.14	240.66	67.01	43.36	64.707	0.647
30	50.48	343.21	77.21	61.01	79.018	0.790
<b>Mean</b>	<b>29.702 ± 1.516 (Standard error)</b>	<b>319.542 ± 10.38 (Standard error)</b>	<b>93.585 ± 30.204 (Standard deviation)</b>	<b>73.571 ± 28.738 (Standard deviation)</b>	<b>76.905 ± 7.144 (Standard deviation)</b>	<b>0.769 ± 0.071 (Standard deviation)</b>

**DHOLE (*Cuon alpinus*)**

S. No.	Hair length (mm)	Root length (μm)	Cortical thickness (μm)	Medullary thickness (μm)	Medulla Percentage in hair (%)	Medullary index
1	23.18	927.42	122.88	70.61	57.463	0.575
2	26.37	931.83	160.42	121.69	75.857	0.759
3	34.25	811.31	142.68	86.99	60.969	0.610
4	29.67	768.72	182.08	147.08	80.778	0.808
5	28.69	746.06	156.79	88.65	56.541	0.565
6	39.22	744.24	135.05	87.31	64.650	0.647
7	26.12	702.97	138.85	78.31	56.399	0.564
8	23.72	823.06	162.57	117.52	72.289	0.723
9	37.6	965.94	148.29	107.24	72.318	0.723
10	40.25	694.56	136.16	75.91	55.751	0.558
11	30.24	611.93	156.86	109.6	69.871	0.699
12	29.57	728.16	152.24	88.93	58.414	0.584
13	37.76	587.56	164.08	124.75	76.030	0.760
14	31.14	766.7	159.92	106.75	66.752	0.668
15	25.96	687.84	161.58	101.64	62.904	0.629
16	26.85	718.77	153.46	89.54	58.347	0.583
17	27.49	783.53	158.52	107.38	67.739	0.677
18	29.52	658.29	159.81	115.23	72.104	0.721
19	28.72	659.31	162.55	120.38	74.057	0.741
20	35.86	727.76	178.53	133.27	74.649	0.746
21	23.9	634.27	171.52	130.29	75.962	0.760
22	31.64	605.37	112.12	70.39	62.781	0.628
23	29.91	619.71	153.92	104.62	67.970	0.680
24	26.32	572.33	140.87	97.69	69.348	0.693
25	15.47	635.76	93.88	59.64	63.528	0.635
26	29.75	674.89	128.51	87.22	67.870	0.679
27	26.07	702.23	121.51	80.57	66.307	0.663
28	28.84	559.62	126.52	83.22	65.776	0.658
29	35.96	647.86	101.9	62.34	61.178	0.612
30	28.07	611.43	137.86	89.76	65.110	0.651
<b>Mean</b>	<b>29.604 ± 0.976 (Standard error)</b>	<b>710.310 ± 19.109 (Standard error)</b>	<b>146.064 ± 21.396 (Standard deviation)</b>	<b>98.151 ± 21.898 (Standard deviation)</b>	<b>66.657 ± 6.821 (Standard deviation)</b>	<b>0.667 ± 0.068 (Standard deviation)</b>

**SLOTH BEAR (*Melursus ursinus*)**

S. No.	Hair length (mm)	Root length (µm)	Cortical thickness(µm)	Medullary thickness (µm)	Medulla Percentage in hair (%)	Medullary index
1	71.6	1128	94.1	10.57	11.233	0.112
2	77.76	961.53	151.35	32.75	21.639	0.216
3	68.73	1017.42	100.21	29.64	29.578	0.296
4	83.93	1488.28	106.33	26.01	24.462	0.245
5	75.22	1325.52	134.66	16.02	11.897	0.119
6	77.49	1439.72	152.07	25.29	16.630	0.166
7	98.77	1200.48	184.53	19.04	10.318	0.103
8	91.59	1056.17	158.1	16.1	10.183	0.102
9	79.25	1087.35	147.9	45.63	30.852	0.309
10	114.38	1574.53	138.97	34.02	24.480	0.245
11	76.27	1094.82	121.95	32.42	26.585	0.266
12	87.08	1251.19	107.09	32.9	30.722	0.307
13	93.39	1021.46	106.08	29.2	27.526	0.275
14	93.63	1528.21	137.58	33.3	24.204	0.242
15	105.19	1208.82	141.16	32.29	22.875	0.229
16	100.73	1112.51	145.48	31.95	21.962	0.220
17	107.02	1514.19	131.82	31.29	23.737	0.237
18	98.74	1103.38	126.93	30.72	24.202	0.242
19	88.48	1313.43	148.29	50.33	33.940	0.339
20	89.85	1144.92	106.4	20.15	18.938	0.189
21	92.85	1158.68	102.9	7.34	7.13	0.071
22	43.84	1500.17	100.9	6.72	6.66	0.067
23	79.91	1434.25	136.03	13.41	9.86	0.099
24	64.82	1279.2	136.03	20.02	14.72	0.147
25	70.7	1455.99	102.69	12.38	12.06	0.121
26	68.66	1095.53	122.3	12.44	10.17	0.102
27	64.54	1349.53	101.02	11.4	11.28	0.113
28	93.31	939.72	87.28	7.52	8.62	0.086
29	100.01	850.53	88.68	15.03	16.95	0.169
30	57.28	1431.75	115.82	20.62	17.80	0.178
<b>Mean</b>	<b>83.834 ± 2.930 (Standard error)</b>	<b>1235.57 ± 36.63 (Standard error)</b>	<b>124.488 ± 23.834 (Standard deviation)</b>	<b>23.550 ± 11.302 (Standard deviation)</b>	<b>18.707 ± 8.033 (Standard deviation)</b>	<b>0.187 ± 0.080 (Standard deviation)</b>

**INDIAN GREY MONGOOSE (*Herpestes edwardsii*)**

S. No.	Hair length (mm)	Root length (µm)	Cortical thickness(µm)	Medullary thickness (µm)	Medulla Percentage in hair (%)	Medullary index
1	48.96	112.5	153.43	102.37	66.72	0.67
2	39.88	423.18	134.78	78.67	58.37	0.58
3	49.44	494.79	139.88	89.61	64.06	0.64
4	48.3	444.15	130.12	86.53	66.50	0.67
5	61.94	388.95	85.59	56.24	65.71	0.66
6	47.61	456.18	123.35	76.5	62.02	0.62
7	50.57	538.36	129.45	83.56	64.55	0.65
8	53.22	508.83	128.74	69.49	53.98	0.54
9	52.9	341.25	118.48	86.69	73.17	0.73
10	49.02	436.14	131.13	91.62	69.87	0.70
11	35.52	276.93	120.62	84.78	70.29	0.70
12	55.14	409.46	130.13	79.85	61.36	0.61
13	46.84	452.6	136.57	92.93	68.05	0.68
14	53.13	387.63	114.15	84.23	73.79	0.74
15	43.91	443.02	148.21	95.93	64.73	0.65
16	49.69	347.08	115.04	63.74	55.41	0.55
17	44.45	395.29	138.72	79.59	57.37	0.57
18	41.68	362.54	92.14	64.44	69.94	0.70
19	44	444.74	142.26	85.16	59.86	0.60
20	45.49	462.7	129.39	73.25	56.61	0.57
21	73.64	482.28	122.73	68.72	55.99	0.56
22	69.77	475.55	139.88	88.27	63.10	0.63
23	59.12	461.29	148.88	89.11	59.85	0.60
24	61.13	348.26	148.58	91.63	61.67	0.62
25	71.77	464.36	145.53	91.34	62.76	0.63
26	74.99	391.35	137.67	92.68	67.32	0.67
27	58.62	317.98	98.2	69.06	70.33	0.70
28	76.23	498.13	122.21	83.9	68.65	0.69
29	66.57	514.96	151.57	93.97	62.00	0.62
30	64.63	496.06	150.56	98.9	65.69	0.66
<b>Mean</b>	<b>54.605 ± 2.033 (Standard error)</b>	<b>419.21 ± 15.68 (Standard error)</b>	<b>130.266 ± 17.150 (Standard deviation)</b>	<b>83.092 ± 11.236 (Standard deviation)</b>	<b>63.991 ± 5.330 (Standard deviation)</b>	<b>0.640 ± 0.053 (Standard deviation)</b>