

Original Research Article

PERFORMANCE OF TRANSPLANT AUS RICE VARIETIES UNDER DIFFERENT NITROGEN MANAGEMENT PRACTICES

Abstract

The experiment was carried out during April to July 2015 at the Agronomy Field Laboratory of Patuakhali Science and Technology University, Dumki, Patuakhali ~~with a view~~ to find out the influence of different nitrogen management and variety on the yield performance of transplant Ausrice. The experiment consisted of four levels of nitrogen viz. Control (without N), 30 kg N ha⁻¹, 60 kg N ha⁻¹ and urea super granule @ 52 kg N ha⁻¹ and four T. Ausrice varieties viz. *Kali Hitta*, *ChaitaBoro*, *Abdul Hai* and *Gota IRRI*, and The experiment was laid out in a split-plot design with three replications. The levels of nitrogen were assigned in the main plot and varieties were allocated in the sub-plots. Nitrogen management, variety and their interactions exerted significant influence on plant characters, yield contributing characters and yield of transplanted Ausrice. In the case of nitrogen management, the tallest plant (161.60 cm), maximum leaf area index (2.97), the highest number of effective tillers hill⁻¹ (15.45), longest panicle (24.30 cm) with the maximum number of filled grains (94.73), 1000-grain weight (29.97 g) and grain yield (2.48 t ha⁻¹) were obtained from USG @ 52 kg N ha⁻¹ and the shortest plant height (136.90 cm), lowest leaf area index (1.78), lowest number of effective tillers hill⁻¹ (8.43), shortest panicle (18.84 cm) with the lowest numbers of filled grains panicle⁻¹ (53.18), 1000-grain weight (24.33 g) and grain yield (1.40 t ha⁻¹) were obtained in control (N₁₌₀, kg N ha⁻¹). Among the varieties, *ChaitaBoro* gave the tallest plant height (151.60 cm) and maximum leaf area index (2.54). While the highest number of effective tillers hill⁻¹ (12.20), longest panicle (22.42 cm) with the maximum number of filled grains panicle⁻¹ (73.50), highest 1000-grain weight (27.41 g) and highest grain yield (2.39 t ha⁻¹) were recorded from *Gota IRRI* than other varieties. In case of interaction, *Gota IRRI* Fertilized with USG at 52 kg N ha⁻¹ produced the highest number of effective tillers hill⁻¹ (16.87), panicle length (25.13 cm), number of grains panicle⁻¹ (105.70) and grain yield (3.13 t ha⁻¹). The lowest number of effective tillers hill⁻¹ (8.13), lowest panicle length (17.47 cm) with minimum numbers of filled grains 47.67) and grain yield (1.12 t ha⁻¹) were produced by the interaction of control (N₁₌₀, kg N ha⁻¹) and *Kali Hitta*. So, cultivation of transplant Ausrice (*Gota IRRI*) appeared to be the best performance with USG @ 52 kg N ha⁻¹. and recommended to the end users.

Comment [H1]: Non-standard abbreviations should not be used, if necessary they should be clearly defined at first use.

Keywords: Ausrice , nitrogen management , variety, yield.

Introduction

Rice (*Oryza sativa* L.) plays an important role in the food security of the world and it is the staple food of more than fifty percent people of the world's population (IRRI, 2015). At present a hectare of

rice field can feed 27 people, it must feed 43 people by 2050 (Chen S, 2012). In Bangladesh rice is the staple food of about 17 million people which covers about 75% of the total-cropped area (Nasim et al., 2017). For increasing yield in rice high yielding varieties along with appropriate management practices are needed. Among management practices, nitrogen fertilizer management is a key nutrient element ~~which that~~ plays a vital role in vegetative growth, development of yield components and yield of rice (BRRI, 2000). Efficient fertilizer management gives a higher yield of ~~the~~ crop and reduces fertilizer cost (Hossain and Islam, 2006). Like other rice-growing countries, Bangladesh also needs to increase production to feed ~~her-its~~ overloaded population since rice is the staple food, though the country occupies 4th position in the world ~~with respect~~ to rice area and production (FAO, 2020). Aman season is rainfed but Boro is ~~totally~~ an exhaustive season where a considerable amount of electric power is used for ~~irrigation~~. Again, sub-soil water is mining for irrigation which exerts a negative impact on our natural resources. But the main drawback is that ~~the~~ average yield of aus rice (2.16 t ha⁻¹) is lower than aman and boro ~~season~~. Besides that, in the southern region of Bangladesh, most of the farmers cultivate aus rice with local varieties whose yield potential is very low. Out of ~~the~~ total Aus area (1.05 mha), local varieties covers about 0.26 mha with an average yield of only 1.27 t ha⁻¹ (DAE, 2014). This lower yield in aus rice is characterized by local low yielding ~~varieties, varieties~~; though most of the farmers prefer it and poor management especially nitrogen fertilizer management has not been optimized in local aus rice varieties. ~~The~~ optimum dose of N is one of the most effective means for maximum yield of rice which could increase about 70-80% yield in ~~rice~~. Nitrogen fertilizer increases tillering, ~~and~~ vegetative growth, ~~increases~~ plant height, grain and straw yield and number of heads usually are proportional~~ly~~ to the amount of nitrogen added to ~~the~~ soil. Application of total N fertilizer in several splits matching the demand of the crop for N at critical stages of growth, deep placement of N in the reduced zone of the soil or thorough incorporation in the soil, use of coated/modified (e.g. USG) fertilizers are some of the useful techniques which improve N-use efficiency in rice. As N fertilizer ~~is the~~ the main promoter of crop growth and yield, it is important to improve management ~~practices that minimize~~ N losses and increase the recovery of applied N by the crop. This will increase productive efficiency and reduce negative impact of N use on ~~the environment~~ ~~the environment~~. In the light of ~~the~~ above points, the present study was undertaken to, evaluate the effect of Aus rice varieties on growth and yield performance, observe the effect of nitrogen fertilizer on Aus rice, and the interaction effect of Aus rice varieties and nitrogen management practices.

Comment [H2]: Follow citation style of the journal in all section of the manuscript

Comment [H3]: Support with references/data

Comment [H4]: Support with reference/es

Comment [H5]: Reference (support with references

Comment [H6]: Rephrase this part using your own words rather than direct use of other sources

Formatted: Pattern: Clear (Custom Color(RGB(217,149,148)))

Materials and Methods

Location and site

The experiment was carried out at the Agronomy experimental field of Patuakhali Science and Technology University, Patuakhali, Bangladesh during the period from April, 2015 to July, 2015. The area is located at 22.26°N latitude and 90.22°E longitude at an elevation of 1.5 m above sea level.

Location and site

~~The experiment was conducted at experimental field of Patuakhali Science and Technology University, Dumki, Patuakhali with geographical location of 22.26°N latitude and 90.22°E longitude at an elevation of 1.5 m above the sea level.~~

Soil type

The experimental field was medium high land with silty clay textured soil with a pH value of 6.5 belonging to the Ganges Tidal Flood Plain (AEZ 13) in the coastal non-saline zone of Bangladesh. The experimental field was more or less neutral in reaction, low in organic matter content and its general fertility level was also low.

Experimental material

Four popular T. aus rice varieties viz. Kali Hitta, ~~Chaita Boro, Chaita Boro~~, Abdul Hai and Gota IRRI were used as experimental materials. The seeds were collected from local trusted farmers.

Experimental treatments

The experiment was a two factor factorial experiment consisted of four T. aus varieties and four doses of Nitrogen fertilizers. The details of treatments are

Factor A: Nitrogen management -4 (Main plot) Control (No Nitrogen) N₁, Nitrogen @ 30 Kg ha⁻¹ (N₂), Nitrogen @ 60 Kg ha⁻¹ (N₃), Urea super granule @ 52 Kg ha⁻¹ (N₄)

Factor B: T. Aus Variety -4 (subplot) Kali Hitta (V₁), Chita Boro (V₂), Abdui Hai (V₃), Gota IRRI (V₄).

Experimental design and layout

The experiment was laid out in a split-plot design with three replications. Where N-fertilizer levels were assigned in the main plot and varieties were in the sub-plot. The treatments were randomly assigned in each replication. There were 48 unit plots in the experiment. The size of each plot was 4 m × 2.5 m. Each unit plot and block was separated from each other by 50 cm and 1m, respectively. A High border was maintained to control the movement of fertilizers to the adjacent plots.

Raising seedlings

Comment [H7]: Dear author, include the information on temperature, rainfall (amount and its distribution) of the area, and also major crops grown in the area. Support with references

Comment [H8]: ????

Comment [H9]: State shortly the description of this variety with its agro-ecological requirements (RF, altitude, Tem, its potential yield on farmers and research station...)

Formatted: Highlight

Formatted: Highlight

Healthy seeds were selected by a specific gravity method and then sprouted by immersing in water in the bucket for 24 hours. Then the seeds were taken out of the water and kept thickly in a gunny bag. After 48 hours the seeds started sprouting and sown after 72 hours in the nursery bed. A piece of high land was selected for raising seedlings. The nursery plot was irrigated 12 days before of sowing seeds to hasten germination of *Kharif* annual weeds. After that nursery plot was ploughed first by soil turning plough followed by cross ploughing with a cultivator. The sprouted seeds were sown gently and uniformly in the wet nursery beds on 1 April 2015. Proper care was taken to raise the seeding in the nursery. Weeds were removed and irrigation was given in the nursery beds as and when necessary.

Formatted: Pattern: Clear (Custom Color(RGB(229,184,183)))

The experimental land was prepared by a tractor drawn disc plough on 27 April, 2015. Then the land was puddled thoroughly by repeated ploughing and cross ploughing with a country plough and leveled by laddering. Weeds and stubble of the previous crop were removed from individual plots and finally plots were leveled so properly by wooden plank that no water pocket could remain in the puddled field.

Formatted: Pattern: Clear (Custom Color(RGB(229,184,183)))

Application of fertilizers

Fertilizer with a uniform dose except for nitrogen of 80, 60, 21 and 8 kg per hectare P, K, S and Zn through TSP, MOP, gypsum and Zinc Sulphate was applied in all the plots. Full doses of phosphorus, potassium, gypsum and zinc sulphate were applied as basal applications just before transplanting. Nitrogen was applied as per treatment in the form of prilled urea and urea super granule (USG). Prilled urea were urea was applied in three equal splits at 15, 30 and 45 days after transplanting (DAT) and USG were applied at a time at 10 DAT.

Transplanting of seedlings

The seedlings were uprooted from the nursery bed on the day of transplanting. Thirty days old seedlings were transplanted on 5 May 2015 in the main field with 4 seedlings hill⁻¹ spaced at 20 cm × 20 cm.

To ensure and maintain the normal growth and development of the crop intercultural operations were done at proper time. The following intercultural operations were done.

Irrigation and drainage

A thin film of water was maintained at the time of transplanting. The plots were kept in saturated condition for a week after transplanting. Two irrigation were applied during the first month of transplanting. At the latter stage, frequent rainfall occurred, so there was no need for further irrigation. Excess water was drained out from the plots before 15 days of harvest to enhance maturity of the crop.

Plant protection measures

Crops were mildly attacked by rice stem borer at tillering stage of crop and rice bug at the dough stage. Furadan @ 10 kg ha⁻¹ were applied to control the stem borer and rice bug, respectively.

Sampling, Harvesting and Processing

Harvesting was done depending upon the maturity of the crop. Maturity of the crop was determined when 90% of the grain became golden yellow color. Five hills from each plot excluding border hills were selected at random prior to harvesting and taken out for studying yield attributes data. The crop of individual plots was harvested on 28 July 2015. These plants were taken out with respective tag levels. An area of central 5 m² in each plot excluding the crop sampling zone was harvested for measurement of grain and straw yields. The harvested crop of each plot was separately bundled, properly tagged and then brought to the threshing floor. The crop was threshed by pedal thresher. Grain was sun dried and cleaned. Straw was also sun dried properly. Finally, grain and straw yields were adjusted to 14% moisture level and converted to ton per hectare.

Formatted: Pattern: Clear (Custom Color(RGB(229,184,183)))

Data Collection

Data was collected on pPlant height (cm), lLeaf area index, Ddays to 50% flowering-, nNumber of total tillers per hill-, nNumber of effective tillers per hill, nNumber of non-effective tillers per hill-, pPanicle length (cm)-, nNumber of grains per panicle-, 1000-grain weight (g), Grain yield (t ha⁻¹)-, sStraw yield (t ha⁻¹) and Harvest index (%).

Statistical Analysis

~~Data were compiled and tabulated in proper form for statistical analysis.~~ The ~~recorded~~ collected data on various plant characters were statistically analyzed using the 'Analysis of variance technique' with the help of a 'R' and the mean differences s were adjusted by Duncan's Multiple Range Test (Gomez and Gomez, 1984).

RESULT AND DISCUSSIONS

Effect of nitrogen management, variety and interaction (nitrogen and variety)

Plant height at maturity

The plant height of transplant Ausrice varied significantly due to different nitrogen management (Table 1). The plant height ranged from 136.90 cm to 161.60 cm. The tallest plant (161.60 cm) was recorded at USG 52 kg N ha⁻¹. Increased plant with the treatment of USG might be the availability of sufficient nitrogen ~~in sufficient~~ amount for the rice plant throughout the life cycle,

which might have favored increased cell division and cell enlargement. Faraji (1998) recorded different plant heights due to different nitrogen rates.

The plant height at maturity of transplant Ausrice was significantly influenced by variety (Table 2). The tallest plant (151.60 cm) was recorded in *ChaitaBoro* followed by *Kali Hitta* (149.37 cm) and *Abdul Hai* (147.45 cm). These differences are mostly due to the genetic variation among the varieties. These results were consistent with those of Om *et al.* (1998), Khisha, (2002) and Rahman, (2003) who recorded variable plant height among the varieties.

Interaction of nitrogen management and variety, the highest plant height (165.30 cm) was found from the interaction between USG @ 52 kg N ha⁻¹ × *ChaitaBoro* and the lowest plant height (135.60 cm) was found from *Gota IRRI* (V₄) with control (N₁) application, which was similar to 30 kg prilled urea × *Abdul Hai* interaction (135.80 cm) (Table 3)

Leaf area index of transplant Ausrice

Leaf area of transplant Ausrice was varied significantly due to nitrogen management practices. Application of USG (52 kg N ha⁻¹) produced maximum LAI (2.97), while the lowest LAI (1.78) was recorded from control treatment. No significant variation was observed in the case of prilled urea at both N₃₀ and N₆₀ levels (Table 1).

Varieties showed significant variation as the LAI ranged from 2.13 to 2.54 (Table 2). The maximum LAI (2.54) was recorded in *ChaitaBoro* and the lowest LAI (2.13) was recorded from *Gota IRRI*. This variation in LAI may be due to the genetic make of the varieties.

In interaction, variety *ChaitaBoro* and *Kali Hitta* showed statistically similar (significantly at 5% level) to LAI at USG @ 52 kg N ha⁻¹, which were 3.26 and 3.07, respectively. On the other hand, *Gota IRRI* at zero N level gave the lowest (1.43) LAI (Table 3).

Days to 50% flowering and maturity

Response to flowering showed dependencies on the level of application of nitrogen fertilizer. A significant variation of days to 50% flowering was found highest (104.80 days) in USG @ 52 Kg N ha⁻¹ and without N application evoked early flowering than any other treatment, which took only 62.83 days (Table 1).

ChaitaBoro took a longer time (96.42 days) to reach 50% flowering followed by *Gota IRRI* (93.67 days), and variety *Kali Hitta* took shortest time (83.67 days) to reach 50% flowering. Time required to set flowers showed significant variation due to nitrogen management and variety interaction at a

5% level of significance (Table 3). *ChaitaBorow* with USG @ 52 Kg N ha⁻¹ took a longer time (111.30 days) to reach 50% flowering and *Kali Hitta* without N fertilizer reported earlier (57.00 days) in 50% flower production.

Days to maturity

Response to maturity also showed significant variation due to the level of application methods of nitrogen fertilizer. A significant variation of days to maturity was found highest (128.33 days) in USG @ 52 Kg N ha⁻¹ and the lowest days to maturity (104.42 days) was recorded from without N application (Table 1). Among *T. aus* varieties *ChaitaBoro* took longer time (125.67 days) to reach maturity followed by *Gota IRRI* (122.33 days), and variety *Kali Hitta* took shortest time (111.00 days) to attain maturity (Table 2). In interaction, *ChaitaBoro* with USG @ 52 Kg N ha⁻¹ took a longer time (137.00 days) to get matured and *Kali Hitta* without N fertilizer reported earlier in maturity (98.67 days) (Table 3).

Number of effective tillers hill⁻¹

Results showed that USG 52 kg N ha⁻¹ produced the highest number of effective tillers hill⁻¹ (15.45) and the lowest one (8.43) was obtained in control (N₀ kg N ha⁻¹) (Table 1). Adequacy of nitrogen as USG probably favored the cellular activities during panicle initiation and development which led to increased number of productive tillers hill⁻¹. Ahmed *et al.* (2005) also reported that the number of effective tillers hill⁻¹ increased with the better management of nitrogen.

The highest number of effective tillers hill⁻¹ was produced by *Gota IRRI* (12.20) and the lowest one (10.50) was produced by *Kali Hitta* (Table 2). The probable reason ~~of for~~ the differences in producing the effective tillers hill⁻¹ is the genetic make-up of the variety which is primarily influenced by heredity. This finding corroborates with those reported by BINA, (1998) Om *et al.* (1998) and Bhowmick and Nayak, (2000) who stated that the effective tillers hill⁻¹ was varied with variety.

The effect of interaction between nitrogen management and variety was found to be significant in respect of number of effective tillers hill⁻¹ (Table 3). The highest number of effective tillers hill⁻¹ (16.87) was counted in the treatment combination of USG 52 kg N ha⁻¹ with V₄.

Number of non-effective tillers hill⁻¹

A significant difference was found due to the level of nitrogen as a source of nitrogen in respect of the number of non-effective tillers hill⁻¹ (Table 1). The total non-effective tillers hill⁻¹ ranged from

Comment [H10]: I didn't see the importance of this section, better to removing it.

0.52 to 2.86. The results show that the highest number (2.86) of non-effective tillers hill⁻¹ was produced by the control (N₁ kg N ha⁻¹) treatment. The lowest number (0.52) of non-effective tillers was found at USG 52 kg N ha⁻¹. The observation in the formation of non-effective tillers hill⁻¹ was decreased with an appropriate nitrogen level. The highest number of non-effective tillers hill⁻¹ was produced by *Kali Hitta* (1.85) and the lowest one (1.24) was produced by *Gota IRRI*. The probable reason of for the differences in producing the non-effective tillers hill⁻¹ is the genetic make-up of the variety which is primarily influenced by heredity. Devarajauet *al.* (1998) and Rahman, (2006) also reported that the number of non-effective tillers hill⁻¹ was significantly influenced by varieties.

The N number of non-effective tillers hill⁻¹ was significantly affected by the interaction. The M maximum number of non-effective tillers hill⁻¹ (3.47) was produced by the combination of *Kali Hitta* and control (N₁kgN ha⁻¹), which was statistically similar with *ChaitaBoroat* control (3.20) and the minimum number of non-effective tillers hill⁻¹ (0.27) was produced by the interaction of *Gota IRRI* and USG 52 kg N ha⁻¹, which was statistically similar to *Abdul Haiat* USG 52 kg N ha⁻¹ (0.47) (Table 3).

Length of panicle (cm)

Table 1 shows that the longest panicle length (24.30 cm) was found in treatment with USG 52 kg N ha⁻¹ and the shortest one (18.84 cm) was found from control i.e. without N application (N₁).

Gota IRRI had the taller panicle length (22.42 cm) followed by *Abdul Hai* (21.88 cm), *ChaitaBoro* (21.32 cm) and *Kali Hitta*(20.93 cm), respectivelyrespectively. Variation in length of panicle due to variety as obtained in the study might be because of their genetic potentiality. This finding is also supportive to the findings of Bhuiyanet *al.* (2002).

From the table 3 it is obvious that the longest panicle length (25.13 cm) was found in the interaction between *Gota IRRI* × USG @ 52 kg N ha⁻¹ and the shortest panicle length (17.47 cm) was found in the interaction between *Kali Hitta*at without N. The variation in panicle length due to interaction of variety and integrated nitrogen management was also reported by Parvin (2012).

Number of filled grains panicle⁻¹

The highest number of grains panicle⁻¹ (94.73) was obtained from the treatment USG @ 52 kg N ha⁻¹ and the lowest number of grain panicle⁻¹ (53.18) was found in treatment without N, which was again similar to the treatment N₃₀ (59.06). This finding is in contrast with the research findings of Rajarathinam and Balasubramanivan,(1999) where they found the application of N increased the number of filled grains per panicle.

Gota IRRI produced the highest number of grains panicle⁻¹ (73.50) and *Kali Hitta*produced the lowest number of grains panicle⁻¹ (63.12)(Table 2). This might be due to the differences in the

genetic makeup of the varieties. The variable number of grain panicle⁻¹ among the varieties was reported by Kamal *et al.* (1999).

The highest number of grains panicle⁻¹ (105.70) was observed in the interaction between *Gota IRRI* × USG @ 52 kg N ha⁻¹ and the lowest (47.76) was found in the interaction between *Kali Hitta* with N₁. This finding of the study is also supportive with the research findings of Surendra *et al.* (1995).

Number of unfilled grains panicle⁻¹

Different application methods of nitrogenous fertilizers exerted a significant effect on the number of unfilled grains panicle⁻¹. The maximum number of unfilled grains panicle⁻¹ (24.88) was found in treatment without N application and the lowest (7.08) was found in USG @ 52 kg N ha⁻¹ (Table 1)

For variety-, *Kali Hitta* produced the higher number of unfilled grains panicle⁻¹ (18.92) and the lowest (15.07) was recorded from *Gota IRRI*. The results are in agreement with the findings of Chowdhury, (1993) who reported that varietal differences in the number of unfilled grains panicle⁻¹ might be due to genotypic variation.

The highest number of unfilled grains panicle⁻¹ (27.47) was observed in the interaction between *Kali Hitta* having no N, which was statistically identical with *Chaita Boro* at the same level of N (N₁) (26.07) (Table 3).

1000-grain weight

The highest 1000-grain weight (29.97 g) was obtained from the treatment USG @ 52 kg N ha⁻¹. Baligar, (2001) reported that the weight of 1000-grain increased significantly with increasing nitrogen levels. Variety *Gota IRRI* produced the highest weight (27.41 g) of 1000 grains followed by *Abdul Hai* (27.03 g), *Chaita Boro* (26.61 g) and *Kali Hitta* (25.94 g). This might be due to the differences in the genetic makeup of the varieties. These results are in agreement with Hasan, (2007) and Thakur, (1999) who reported differences in 1000-grain weight among the varieties. Significantly the highest weight of 1000 grains (31.13 g) was observed in combination between USG @ 52 kg N ha⁻¹ × *Gota IRRI* and the lowest weight of 1000-grains (23.52 g) was observed in *Kali Hitta* at N₁ (Table 3).

Grain yield

Different levels of application methods of nitrogenous fertilizers exerted a significant effect on the grain yield of T. Ausrice at a 1% level of probability (Table 1). The highest grain yield (2.48 t ha⁻¹) was found in USG @ 52 kg N ha⁻¹. On the other hand, the lowest grain yield (1.40 t ha⁻¹) was recorded in treatment without N. The results also indicate that the application of USG performed better in terms of grain yield compared to the application of prilled urea. The highest grain yield was obtained in the study from the application of USG @ 52 kg N ha⁻¹ was significantly higher than that of any other application levels of nitrogenous fertilizer application. The highest grain yield might be due to the resultant effect of the highest number of effective tillers hill⁻¹ and the highest number of grains panicle⁻¹ as obtained in the treatment. Budhar and Palaniappan, (1993) reported that nitrogen use efficiency was highest in USG than any other levels of urea application.

The effect of variety on grain yield was significant at 1% level of probability (Table 2). The higher grain yield (2.39 t ha⁻¹) was recorded in the variety *Gota IRRI* and the lowest in *Kali Hitta* (1.52 t ha⁻¹). Varietal differences regarding grain yield were reported elsewhere (Patel, 2000; Khisha, 2002).

The interaction effect of different levels of nitrogenous fertilizers and variety was significant on grain yield at a 1% level of probability (Table 3). The highest grain yield (3.13 t ha⁻¹) was observed in combination between USG @ 52 kg N ha⁻¹ × *Gota IRRI*. On the other hand, the lowest grain yield (1.12 t ha⁻¹) was observed in *Kali Hitta* at without N, which was again statistically similar to N₁ with V₂(1.23 t ha⁻¹).

Straw yield

The Straw yield was significantly influenced by the application of different levels of nitrogenous fertilizers (Table 1). The highest straw yield (3.67 t ha⁻¹) was found in USG @ 52 kg N ha⁻¹ and the lowest straw yield (2.39 t ha⁻¹) was achieved from without application of N. The result showed that the higher dose of nitrogen especially USG influenced vegetative growth in terms of plant height and total tillers hill⁻¹ which resulted in differences of straw yield. These findings corroborated with those of Pasha and Reddy, (2007).

The effect of variety on straw yield was significant at a 1% level of probability (Table 2). The higher straw yield (3.32 t ha⁻¹) was recorded in the variety *Gota IRRI* and the lowest (2.70 t ha⁻¹) was in *Kali Hitta*. Hossain *et al.* (2003) reported variable straw yields among the varieties.

Interaction of different level of nitrogenous fertilizers and variety exerted a significant influence on the straw yield of T. Ausrice varieties (Table 3). The highest straw yield (4.17 t ha⁻¹) was obtained

from the interaction between USG @ 52 kg N ha⁻¹ × *Gota IRRI*, and the lowest straw yield (2.28 t ha⁻¹) was observed in *Kali Hitta* with no nitrogen.

Harvest index

The highest value of harvest index (40.02%) was obtained from USG @ 52 kg N ha⁻¹ and the lowest harvest index (36.69%) was achieved from control (N₁) nitrogen application. A similar result was reported by Bhuiyan, (2007) was found increasing harvest index with increasing N fertilizer.

Harvest index was significantly influenced by varieties at 1% level of probability (Table 2). The higher harvest index (41.61%) was obtained from *Gota IRRI* followed by *AbdulHai* (40.74%) and the lowest of 35.73% and 35.95%, respectively was obtained from *Kali Hitta* and *ChaitaBoro*, respectively. Kabiret *al.* (2004) reported variable harvest index among varieties.

The effect of interaction between levels of nitrogenous fertilizers and variety on harvest index was significant at 5% level of significance (Table 3). The highest value of harvest index (42.92%) was found in the interaction between USG (N₄) @ 52 kg N ha⁻¹ × *Gota IRRI* (V₄) and the lowest value (32.97%) was found in interaction between control (N₁) treatment × *Kali Hitta* (V₁).

Conclusion

According to the result of the experiment, it can be concluded that, the performance of *Gota IRRI* was better than that of *Kali Hitra*, *ChaitaBoro* and *Abdul Hai*. Among different levels of nitrogen, USG @ 52 kg N ha⁻¹ performed better than prilled urea. It can be suggested that USG @ 52 kg N ha⁻¹ and T. Ausvariety, *Gota IRRI*, could be a better combination for transplant Ausrice cultivation in terms of grain yield.

Comment [H11]: This section is short and need to enhance.

REFERENCES

Comment [H12]: Follow the reference style of the journal

- Nasim, M., Shahidullah, S. M., Saha, A., Muttaleb, M. A., Aditya, T. L., Ali, M. A., & Kabir, M. S. (2017). Distribution of crops and cropping patterns in Bangladesh. *Bangladesh Rice Journal*, 21(2), 1-55.
- Ahmed, M.H., Islam, M.A., Kader, M.A. and Anwar, M.P. (2005). Evaluation of urea super granules as source of nitrogen in transplant *Amanrice*. *Pakistan J. Bio1.Sci.* 3(5): 735-737.
- Baligar, A.S. and Ganin, B.A. (2001). Response of rice genotypes to nitrogen level under transplanted condition in Brazil. *Newsl.* 9 (5) : 29-31.
- Bhowmick, N. and Nayak, R.L. (2000). Response of hybrid rice (*Oryza sativa*) varieties to nitrogen, phosphorus and potassium fertilizers during dry season in West Bengal. *Indian J. Agron.* 45(2): 323- 326.
- Bhuiyan, F.I. (2007). Effect of nitrogen and spacing on the yield and yield attributing characteristics of *Boro* rice (cv. BRRI dhan28). M. S. Thesis, Dept. Agron., Bangladesh Agril. Univ., Mymensingh. pp. 42-44.
- Bhuiyan, N.I., Paul, D.N.R. and Jabber, M.A. (2002). Feeding the extra millions by 2025: Challenges for rice research and extension in Bangladesh. Proc. Workshop on Modern Rice Cultivation in Bangladesh, BRRI, Joydebpur, Gazipur, January 29-31.
- BINA (Bangladesh Institute of Nuclear Agriculture) 1996. Annual Report (1998-99). Bangladesh Inst. Nucl. Agric., Mymensingh. P. 175.
- BRRI (Bangladesh Rice Research Institute) 2000: "*Adhunik Dhaner Chash*" Bangladesh Rice Res. Inst. Joydebpur, Gazipur, Bangladesh, pp. 7-17.
- Budhar, M.N. and Palaniappan, S.P. (1993). Effect of source and levels of nitrogen on grain yield, yield attributes, N uptake, recovery and response by rice in deep water conditions *Crop Res.* 6(2): 213-216.
- Chen, S. (2012). Bid to lift hybrid rice yield to 15 tonnes a hectare slammed. South China Morning Post, 04 December 2012.
- Chowdhury, M.J.U. (1993). Effect of variety and number of seedlings hill⁻¹ on the yield and its components on late transplanted *amanrice*. *Bangladesh J. Aril. Sci.* 20 (2): 311-316.
- DAE 2014: Department of agricultural extension. <http://www.dae.gov.bd>.
- Devaraju, K.M., Gowda, H. and Raju, B.M. (1998). Nitrogen response of Karnataka Rice Hybrid 2. *Intl. Rice Res. Notes.* 23 (2): 43.

- FAO (Food and Agricultural Organization) (2010). Production Year Book. Food and Agricultural Organization of the United Nations, Rome: 76-77.
- Faraji, M.A. (1998). Effect of splitting and rate of nitrogen application on yield and yield components of rice (*Oryza sativa* L.) Agril. Res. J. 2(3): 25-34.
- Gomez, K.A. and Gomez, A.A. (1984). Statistical Procedure for Agricultural Research. 2nd Edn. AWileyInterscience Publication, John Wiley and Sons, New York. pp. 207-215.
- Hasan, S.M. (2007). Effect of level of urea supergranules on the performance of transplanted *Aman* rice. M. Sc. Ag. Thesis Dept. of Agronomy, BAU, Mymensingh.
- Hossain, S.M.A. and Islam, M.S. (2006). Fertilizer Management. in Bangladesh. Adv. Agron. Res. in Joydebpur, Gazipur. pp. 48-54.
- Hussain, M.A., Mani, H. and Tanimu, B. (2003). Effect of nitrogen, phosphorus and planting pattern on growth and yield of rice/sorghum intercrop. *Journal of Sustainable Agricultural Environment* 7 19-25.
- IRRI (International Rice Research Institute) 2013: Annual Report Int. Rice Res. Inst. Los Banos, Philippines. p. 179.
- Kabir, M.E., Kabir, M.R., Jahan, M.S. and Das, G.G. (2004). Yield performance of three aromatic fine rice in a coastal -medium -high land. *Asian. J. Plant. Sci.* 3 (5): 531- 563.
- Kamal, A.M.A, Islam, M.R. and Chowdhury, B.L.D. (1999). Growth performance and nutrient uptake by modern varieties of rice under irrigation condition in Bangladesh. *Thai J. Agril. Sci.* 32(1): 105-110.
- Khisha, K. (2002). An Evaluating of Madagascar System of Rice Production in *Aman* season with three high potential rice varieties. M.S. Thesis, Dept. Agron., Bangladesh Agril. Univ., Mymensingh. pp. 36-98.
- Om, H., Dhiman, S.D., Nandal, D.P. and Verma, S.L. (1998). Effect of method of nursery raising and nitrogen on growth and yield of hybrid rice (*Oryza sativa*). *Indian J. Agron.* 43(1): 68-70.
- Parvin, S. (2012). Improving yield of *T. aman* rice (cv. BRRI dhan52) through integrated use of nitrogen. MS Thesis Dept of Agron, Bangladesh Agril. Univ., Mymensingh. pp. 23-33.
- Pasha, M.L. and Reddy, M.G. (2007). Performance of dry seeded irrigated rice under different seed densities and nitrogen levels. *Crop Res. Hisar*, 33(1/3): 25-29.
- Patel, J.R. (2000). Effect of water regime, variety and blue green algae on rice (*Oryza sativa*). *Indian J. Agron.* 45 (1): 103-106.

- Rahman, M.A.(2003). Effect of levels of urea super granules and depth of placement on the growth and yield of transplant *Amanrice*. M. S. Thesis, Dept. Agron., Bangladesh Agril, Univ., Mymensingh. 100 p.
- Rahman, M.M.(2006). Effect of cultivar, depth of transplanting and depth of placement of urea super granules on growth and yield of *Bororice*. M. S. Thesis, Dept. Agron., Bangladesh Agril. Univ., Mymensingh. 94.
- Rajarathinam, P. and Balasubramaniyan, P.(1999). Effect of plant population and nitrogen on yield attributes and yield of hybrid rice (*Oryza sativa* L.). *Indian Journal of Agronomy* 44(4):717-721.
- Surendra, S., Prasad, R. and Sharma, S.N.(1995). Effects of blue green algae, nitrogen levels and modified urea materials on yield attributes, and yield of wet land rice (*Oryza sativa*) *Indian J. Agron.* 40(4): 594-597.
- Thakur, K.M.(1999). Relative efficiency of prilled urea and modified urea fertilizer in rainfed low land rice. *Indian J. Agron.* 31(1): 87-90.

UNDER PEER REVIEW

Table 1. Effect of nitrogen fertilizer management on yield and yield contributing characters of Ausrice**Comment [H13]:** Follow the norms of the journal in using tables/figures

Nitrogen fertilizer (dose)	Plant height (cm)	Leaf Area Index	50% flowering (days)	Maturity (days)	Effective tillers per hill(no.)	Non-effective tillers per hill (no.)	Panicle length (cm)	Filled grains per panicle (no.)	Un-filled grains per panicle (no.)	1000-seed weight (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Harvest index (%)
Control (N ₁)	136.90 d	1.78 c	62.83 d	104.42 d	8.43 c	2.86 a	18.84 d	53.18 c	24.88 a	24.33 d	1.40 d	2.39 d	36.69 d
N ₃₀	142.4 c	2.16 b	92.00 c	115.83 c	9.70 c	1.58 b	20.95 c	59.06 c	19.93 b	25.63 c	1.65 c	2.67 c	37.98 c
N ₆₀	153.60 b	2.46 b	103.00 b	126.25 b	11.43 b	1.28 b	22.47 b	67.73 b	15.65 c	27.06 b	2.19 b	3.34 b	39.34 b
USG ₅₂	161.60 a	2.97 a	104.80 a	128.33 a	15.45 a	0.52 c	24.30 a	94.73 a	7.08 d	29.97 a	2.48 a	3.67 a	40.02 a
LSD _{0.05}	4.79	0.31	1.12	1.14	1.72	0.70	1.31	7.69	1.99	1.07	0.01	0.02	0.11
Significance level	**	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	1.33	5.23	1.21	0.82	5.68	11.82	1.63	4.87	5.37	1.01	0.73	0.57	0.28

In a column figures with same letter or without letter do not differ significantly whereas figures with dissimilar letter differs significantly (as per DMRT)

Table 2. Effect of variety on yield and yield contributing characters of Ausrice

Variety	Plant height (cm)	Leaf Area Index	50% flowering (days)	Maturity (days)	Effective tillers per hill (no.)	Non-effective tillers per hill (no.)	Panicle length (cm)	Filled grains per panicle (no.)	Un-filled grains per panicle (no.)	1000-seed weight (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Harvest index (%)
Kali Hitta	149.37 b	2.42 b	83.67 d	111.00 d	10.50 c	1.85 a	20.93 d	63.12 c	18.92 a	25.94 d	1.52 d	2.70 d	35.73 c
ChaitaBoro	151.60 a	2.54 a	96.42 a	125.67 a	10.93 bc	1.67 b	21.32 c	68.06 b	17.50 b	26.61 c	1.71 c	3.01 c	35.95 c
Abdul Hai	146.07 c	2.28 c	88.83 c	115.83 c	11.38 b	1.48 c	21.88 b	70.03 b	16.06 c	27.03 b	2.10 b	3.04 b	40.74 b
Gota IRRI	147.45 c	2.13 d	93.67 b	122.33 b	12.20 a	1.24 d	22.42 a	73.50 a	15.07 d	27.41 a	2.39 a	3.32 a	41.61 a
LSD _{0.05}	1.41	0.10	0.92	0.82	0.54	0.16	0.30	2.06	0.76	0.23	0.03	0.03	0.14
Significance level	**	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	3.22	13.43	1.23	0.94	15.27	44.69	6.05	11.20	11.05	4.00	1.64	1.05	0.43

In a column figures with same letter or without letter do not differ significantly whereas figures with dissimilar letter differs significantly (as per DMRT)

Table 3. Combined effect of nitrogen management and variety on yield and yield contributing characters of Ausrice

Interaction (Nitrogen X Variety)	Plant height (cm)	Leaf Area Index	50% flowering (days)	Maturity (days)	Effective tillers per hill (no.)	Effective tillers per hill (no.)	Panicle length (cm)	Filled grains per panicle (no.)	Filled grains per panicle (no.)	1000- seed weight (g)	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Harvest index (%)
N ₁ V ₁	137.20 jk	1.93 jk	57.00 l	98.67 l	8.13 i	3.47 a	17.47 j	47.67 j	27.47 a	23.52 k	1.12 n	2.28 l	32.97 l
N ₁ V ₂	138.90 ij	2.02 ij	68.67 i	110.70 i	8.20 i	3.20 a	18.43 i	54.13 i	26.07 a	24.40 j	1.23 m	2.44 j	33.61 k
N ₁ V ₃	135.80 k	1.75 k	60.33 k	101.30 k	8.67 hi	2.73 b	19.47 h	54.93 i	23.60 b	24.66 j	1.57 j	2.37 k	39.76 e
N ₁ V ₄	135.60 k	1.43 l	65.33 j	107.00 j	8.73 hi	2.03 c	20.00 gh	56.00 hi	22.40 bc	24.73 ij	1.67 i	2.46 j	40.42 d
N ₂ V ₁	143.30 gh	2.20 ghi	85.67 h	108.30 j	9.20 hi	1.80 cd	20.53 fg	57.13 hi	21.80 cd	25.18 hi	1.33 l	2.48 j	34.91 j
N ₂ V ₂	145.20 g	2.26 fgh	96.67 f	121.30 e	9.47 gh	1.60 de	20.73 f	57.77 ghi	20.33 de	25.36 h	1.47 k	2.68 i	35.42 i
N ₂ V ₃	141.70 hi	2.11 g-j	90.33 g	114.00 h	9.73 fgh	1.47 ef	21.13 ef	59.60 gh	18.93 ef	25.87 g	1.82 g	2.71 i	40.21 d
N ₂ V ₄	139.40 ij	2.06 hij	95.33 f	119.70 f	10.40 efg	1.47 def	21.40 e	61.73 fg	18.67 f	26.10 g	1.99 f	2.81 h	41.39 c
N ₃ V ₁	154.60 de	2.50 cde	95.33 f	117.70 g	10.60 efg	1.40 ef	22.00 d	64.67 ef	17.93 fg	26.58 f	1.76 h	2.97 g	37.28 g
N ₃ V ₂	157.10 cd	2.61 cd	109.00 b	133.70 b	10.87 ef	1.27 ef	22.13 d	67.00 de	16.47 gh	26.97 f	1.87 g	3.20 e	36.93 h
N ₃ V ₃	152.30 ef	2.44 def	101.30 e	123.30 d	11.47 e	1.27 ef	22.60 cd	68.67 de	15.00 h	27.03 f	2.36 d	3.34 d	41.44 bc
N ₃ V ₄	150.50 f	2.31 efg	106.30 c	130.30 c	12.80 d	1.20 f	23.13 c	70.60 d	13.20 i	27.67 e	2.75 b	3.85 b	41.72 b
N ₄ V ₁	162.30 b	3.07 a	96.67 f	119.30 f	14.07 c	0.73 g	23.73 b	83.00 c	8.47 j	28.47 d	1.85 g	3.05 f	37.76 f
N ₄ V ₂	165.30 a	3.26 a	111.30 a	137.00 a	15.20 b	0.60 g	24.00 b	93.33 b	7.13 jk	29.70 c	2.27 e	3.73 c	37.85 f
N ₄ V ₃	160.00 bc	2.83 b	103.30 d	124.70 d	15.67 b	0.47 gh	24.33 b	96.93 b	6.70 k	30.57 b	2.66 c	3.75 c	41.55 bc
N ₄ V ₄	158.80 c	2.71 bc	107.70 bc	132.30 b	16.87 a	0.27 h	25.13 a	105.70 a	6.00 k	31.13 a	3.13 a	4.17 a	42.92 a
LSD _{0.05}	2.82	0.21	1.84	1.64	1.08	0.31	0.59	4.12	1.53	0.46	0.05	0.05	0.28
Sig. level	*	*	*	**	*	**	*	**	*	**	**	**	**
CV (%)	1.33	5.23	1.21	0.82	5.68	11.82	1.63	4.87	5.37	1.01	1.64	1.05	0.43

In a column figures with same letter or without letter do not differ significantly whereas figures with dissimilar letter differs significantly (as per DMRT)

Where, V₁ = Kali Hitta, V₂ = ChaitaBoro, V₃ = Abdul Hai and V₄ = Gota IRRI

F₁ = Control (0 kg N ha⁻¹), F₂ = Prilled urea @ 30 kg N ha⁻¹, F₃ = Prilled urea @ 60 kg N ha⁻¹, and F₄ = USG @ 52 kg N ha⁻¹