

Short communication

***In vitro* Evaluation of Three Conservation Methods of Probiotics from 42-day-Old Tropical Calves**

ABSTRACT

Aims: The conservation of probiotics is of great importance due to the microorganism's viability; therefore, their properties and benefits depend on it. In this work, we studied three methods of conservation; refrigeration, cryopreservation, and lyophilization of two probiotics isolated from the rumen of 42 day-old calves, previously evaluated: A1D42 and A3D42.

Place and Duration of Study: Laboratorio de Rumiología y Metabolismo Nutricional, Facultad de Estudios Superiores Cuautitlán, Universidad Nacional Autónoma de México, between August 2020 and September 2021.

Methodology: It was evaluated the viability of the probiotics after of 0, 30, 60 and 90 days of refrigeration, cryopreservation, and lyophilization as preservation methods. The effect of the lyophilized probiotic A1D42 on the digestibility *in vitro*, such as its effect on the volatile fatty acids (VFAs) production was also determined.

Results: Cryopreserved and lyophilization methods had better performance about the viability in both probiotics. A1D42 lyophilized maintained its viability (67%) until 60 days and after 90 days it was affected significantly. Despite the fact that VFA production *in vitro* did not increase with probiotic addition, A1D42 significantly increased *in vitro* digestibility by up to 66%. The results demonstrated that lyophilization is the best method of conservation of probiotic production due to it preserves vial cells, as well as the properties and effects of the probiotic.

Comment [H1]: The use of pronouns in reporting experiment or writing of manuscripts is not encouraged. Kindly recast the sentence here and in other places in the study

Comment [H2]: Out of place

Comment [H3]: What do they stand for? Abbreviations must on first mention be defined before subsequent use.

Conclusion: Therefore, lyophilized A1D42 probiotic can be used as an additive in calf diets, which could improve their digestibility, and, therefore, their weight gain, which could have a positive impact on animal production, in addition to the health benefits.

Keywords: Probiotics; calves; lyophilization; refrigeration; cryopreservation.

1. INTRODUCTION

Probiotics are live microorganisms which administered in adequate doses to confer health benefits to their host animal such as nutritional, immunological, bacteriostatic, and bactericide effects [1,2].

Comment [H4]: Reconstruct the sentence

Therefore, they can improve animal productivity [3].

The prohibition of antibiotics used in animals as growth promoters by the European Union (Regulation CEE 1831/2003), increased the use of probiotics in animals. Therefore, nowadays the need and interest to optimize probiotic production, researching and applying new strategies for production improvement [4], mainly due to the low viability of probiotics during commercial process and storage, for this reason it is necessary to study new methods of preservation to maintain their viability, properties, and effects.

Cryopreservation and refrigeration are easy and cheap preservation methods [5]. Nonetheless, keeping refrigerated is not a great preservation method for probiotics due to its effect on its viability [6].

Freezing-thawing was reported to damage bacterial cells, inhibit their growth and reduce or affect metabolic activity, which affects its viability [7].

Comment [H5]: Check these sentences. If refrigeration is not a great preservation method for probiotics, do you need to consider refrigeration in your study?

Slowly freezing causes severe cellular damage and can inactivate the cell, whereas fast freezing can decrease this cellular damage [8]. Moreover, the commercial production of frozen probiotics is complicated to manage, transport and to storage. Then, in recent days, had been studied and used other preservation methods more efficiently, which allows for easier handling of the probiotic. One of these methods is lyophilization (freeze-drying) which was reported to be a great preservation method for probiotics [6,9-11].

The use of probiotics on calves has a positive effect on weight gain, digestively, volatile fatty acids (VFA) production, then, animal production. High feed digestibility and VFA production will provide more energy for the animal, and metabolize it as a fuel for body and animal production. It was reported that some lactic acid bacteria (LAB) increase digestibility when added to the rumen fluid [12-14].

Hence, the aim of this study was to evaluate the effect of three preservation methods; refrigeration, cryopreservation and lyophilized on the viability of two calves' bacterial consortia that previously demonstrated their potential as probiotics. Also, it was studied the effect of a lyophilized probiotic on the digestibility and production of VFA *in vitro*.

Comment [H6]: Remove "was"

2. MATERIALS AND METHODS

2.1 Sample preparation

The bacteria consortiums (BC) were inoculated in MRS (Man, Rogosa y Sharpe) broth, and were incubated at 37 °C for 18 hours when the culture was near the stationary phase of growth (10^9 CFU/mL). After, they were refrigerated (4°C), cryopreserved (-70°C), and lyophilized to conserve them.

2.2 Refrigeration

The BC were conserved in their liquid state (in MRS broth) and were stored at 4 °C until their use.

2.3 Cryopreservation

The BC were centrifugated at 16,000 x g for 10 min and were washed twice with 10 mL of Ringer solution. The pellet was suspended in 20 mL of MRS broth with glycerol (20 %, v/v, ratio), it was mixed on vortex and was immediately frozen in liquid nitrogen (-196 °C) to do not affect the cells. After, it was stored at -70 °C [15,16].

Comment [H7]: Centrifuged, not centrifugated

Comment [H8]:

Comment [H9]: Reconstruct the sentences

2.4 Lyophilization

2.4.1 Cryoprotective medium

The cryoprotectant used in this study was skim milk powder (Svelty, Nestlé, México) in 24%, which showed high capacity as a cryoprotectant [9,10,15,17-18]. Moreover, it is cheap and safe to use in the food industry. The cryoprotectant medium was prepared by suspending the cryoprotectant 24 g in 100 mL of distilled water and was sterilized at 121 °C for 20 min [17].

2.4.2 Sample preparation for lyophilization

Cells were centrifuged at 16,000 x g for 5 min, the pellet was separated from the liquid. The pellet was washed twice with 40 mL of sterile distilled water and centrifuged again. The pellet was resuspended in 50 mL of skim milk (24%) in special glass containers to be subjected to lyophilization [17, 19].

Comment [H10]: As above

Comment [H11]: As above

2.4.3 Lyophilization procedure

The pellet mixed with the skim milk (24%) was frozen with dry ice (CO₂) and acetone, and it was stored at -20 °C. The samples were freeze-dried at -20 °C for 8 h in a freeze-dryer (LABCONCO) to 5 x10² mbar [17-18].

2.5 Viability of Microorganisms

The number of viable cells (CFU/mL) were determined in the BC before and after refrigeration, cryopreservation, and lyophilization. 100 µL of suitable dilutions of each BC before refrigeration, freezing, and lyophilization was seeded on MRS agar plates and was incubated at 37 °C for 24 h for subsequent counting. After 30, 60, and 90 days of refrigeration, freezing [15], and lyophilization, the CFU/mL was determined. The freeze-dried samples were resuspended in skim milk by shaking, and incubated at room temperature for 15 min, afterwards, 100 µL of appropriated dilutions were seeded in MRS agar plates at 37 °C for 24 h, and the colonies were counted.

The viability of cell suspension was calculated using the following equation [18]:

$$Viability (\%) = \frac{Viable\ cells\ after\ treatment\ \left(\frac{CFU}{mL}\right)}{Viable\ cells\ before\ treatment\ \left(\frac{CFU}{mL}\right)} \times 100$$

2.6 Digestibility of the lyophilized probiotic

The diet used for the digestibility test was grass dried in an oven for 24 h at 60°C and ground.

The diet was heat sealed in Ankom F57 bags (ANKOM Technology Corp., Fairport, NY). Ruminant fluid was collected (2 L) through a rumen cannula and transported to the laboratory in a prewarmed container.

The reagents used to produce the buffer solution were: solution A (10 g of KH₂PO₄, 0.5 g of MgSO₄·7H₂O, 0.5 g of NaCl, 0.1 g of CaCl₂·H₂O, and 0.5 g of urea in 1 L of deionized water) and solution B (15 g of Na₂CO₃ and 1 g of Na₂S₉H₂O in 100 mL of deionized water). A buffer solution

was made just before to each digestion run by warming solutions A and B to 39°C and adding 20 mL of solution B to 1 L of solution A. The pH of the buffer solution was adjusted to 6.8.

Ten bags were introduced in each digestion vessel, 1600 mL (5:1) of the buffer solution, 400 mL of ruminal fluid and the probiotic at doses 4×10^{11} , 4×10^{12} and 4×10^{13} UFC/g were added; the vessels were injected with CO₂, and were placed in the DAISY^{II} digester at 39 °C for 48 h rotating. After that, the bags were washed, dried and were weighted to calculate the digestibility.

2.7 Determination of the total production of volatile fatty acids (VFA's).

2.7.1 Quantification of VFA's

For the determination of VFA concentration, the sample was acquired from the rumen liquid of the *in vitro* digestibility of each treatment. 1.5 mL of the ruminal fluid was taken and centrifuged at 3,500 x g for 10 min at 4°C for pellet sedimentation, then, 1200 µL of supernatant was recovered in a new Eppendorf tube and 240 µL of 25% metaphosphoric acid was added to obtain a 5:1 ratio. The tubes were incubated on ice for 30 min to promote protein sedimentation and immediately centrifuged at 12,000 x g for 15 min at 4°C. The supernatants were filtered through glass fiber membranes. For measurement of VFA, a Varian 3300 gas chromatograph with flame ionization detector and a Zebron ZB-FFAP capillary column with characteristics of length 15 cm, internal diameter of 0.53 mm (Phenomenex, CA, USA) were used. The operating conditions of the gas chromatograph were as follows: column temperature 70°C for 1.5 min and increased to 130°C at a rate of 10°C per min, reaching the final temperature at 7.5 min; injector temperature 190°C; detector temperature 210°C; mobile phase: nitrogen at a flow rate of 9.5 mL/min; pressure: air 60 psi, N₂ 70 psi, and H₂ 40 psi.

2.8 Statistical analysis

Data were measured in triplicate (viability and digestibility) and expressed as means ± standard deviation (SD); differences between samples were analyzed by one-way analysis of variance (ANOVA) and Tukey's Kramer tests, and the differences with $P = .05$ were considered significant.

Data were analyzed using the JMP version 8.0 software (SAS Institute, Cary, NC).

3. RESULTS AND DISCUSSION

3.1 Viability of Microorganisms

Comment [H12]: If the study is CRD as you have stated here, how did you compare A1D42 and A3D42
One way ANOVA was only used for the days and methods of preservation

The viability of A1D42 was conserved with the lyophilization method, nevertheless, at 90 days of lyophilized was reduced significantly (Fig. 1). The viability of refrigerated was affected significantly ($P = .05$) after 30 days (least of 18%), the viability of cryopreserved also was affected after 30 days, although it was maintained until the 90 days (65%).

Comment [H13]: See my earlier comment.meanwhile, it didnt even reflect in Materials and methods




Fig. 1. Viability of BC A1D42 (a) and A3D42 (b) at 30, 60, and 90 days of different methods of preservation (% of the CFU/mL).

The data collected are the mean ± S.D. Different letters indicate differences ($P = .05$)

Unlike the A1D42, the viability of A3D42 was better cryopreserved than lyophilized and liquid (Fig. 1). Despite 95% of the liquid being maintained up to 30 days, at 60 days it was significantly ($P = .05$) affected (20%). Lyophilization was affected significantly after 30 days (39%).

The viability can be preserved over time by the lower temperature (freezing) or by the reduction of the available water (freeze-drying) [20] as can be seen in the Fig. 1. According to the results, it was evident that the refrigeration cannot be a good conservation method as was reported by Serna-Cock et al. [6], the viability of *W. confusa* (LAB) grown in MRS too decreased progressively until the six weeks of storage. Meanwhile, the viability of lyophilized strain persisted unchanged for six weeks, it was preserved throughout time.

Nevertheless, it is reported that slowly freezing produce outside ice that causes severe cellular damage, whereas fast freezing can decrease this cellular damage [8]. During the Freeze-drying process, the viability of the probiotic bacteria may decrease because the freezing step can inactivate the cell [21], although, 60-70% of cells that survived this step can survive the dehydration process [22]. On the other hand, the drying decreases the bound water from bacterial cells, producing damage to surface proteins, cell wall, and the cell membrane. The deduction of water can cause destabilization of the structural integrity of the cellular components, resulting in loss or weakening of function [8].

On the other hand, the skim milk used as a cryoprotectant in this study has demonstrated a great capacity as a cryoprotectant in *Lactobacillus*, *L. lactis* ssp. *Lactis* CECT 5180 remained 44.3% cell viable [9], *L. salivarius* had 22.4% of cell viable [10], *Lactobacillus delbrueckii* subsp. *Bulgaricus* 86.53% mixed with glycerol, sorbitol, and sucrose [18] and *L. plantarum* mixed with sucrose, and trehalose [17]. The cryoprotect effect could be due to it having the capacity to stabilize the cell membrane components and to produce a porous structure in freeze-dried products that makes rehydration easier [11]. Furthermore, skim milk contains proteins that create a protective layer for the cells during the process avoiding damage [23].

The survival of probiotic bacteria in dried powders during the storage depends on several conditions like storage temperature, relative humidity, oxygen content, moisture content of powders, exposure to light, and storage materials [20]. The viability of powder bifidobacterial probiotics during storage was greater at low temperatures (-18 °C) than at 15 °C, room temperature (20 °C), and 25 °C it was reported that the viability of probiotic storage at these temperatures decreased significantly the bacteria viable [24, 25]. The above may explain that although freezing and freeze-drying can better

preserve the probiotic over time, the viability of the probiotic decreases slightly after these processes. Moreover, the probiotic cellular oxidation of membrane fatty acids during storage is associated also with the decrease in viability [26]. Therefore, the addition of antioxidants with storage under vacuum and controlling water activity should be a better alternative and more effective to preserve the probiotic [27].

Therefore, freeze-drying can be used for large-scale production of probiotic powders, despite the little damage and loss of bacterial viability due to stresses such as cold, oxygen, and osmotic stress, during drying and storage [20].

In addition, due to better probiotic management, it is easier and feasible to manipulate dry powder than frozen, A1D42 was conserved appropriately lyophilized until 60 days maintaining its viability (67 %), therefore, this lyophilized probiotic was selected to continue with the next analysis.

3.2 Digestibility of the lyophilized probiotic

All the concentrations of probiotics had higher percentage of digestibility than the control ($P = .05$). The digestibility of A1D42 at 4×10^{13} UFC/mL concentrations had the greatest digestibility (66%), but it was no different with the other concentrations (Fig. 2).




Fig. 2. Digestibility of A1D42 lyophilized probiotic at different doses.

The data collected are the mean ± S.D. Different letters indicate differences (P = .05)

Our results were similar to those obtained by Ridwan et al. [13], where the addition of LAB strain 32 *L. plantarum* in rumen fermentation resulted in digestibility of 65-75%.

Feed digestibility is an essential parameter to measure rumen fermentation, high feed digestibility will supply more energy for the animal, and metabolize it as a fuel for body and animal production. Some studies report that some LAB increases digestibility when added directly to the rumen fluid [12-14].

Ellis et al. [14] report that *L. plantarum* increased significantly the digestibility *in vitro* in rumen compared to the control. The increasing of feed digestibility confirms that LAB can act as a probiotic by stimulating rumen bacteria activity.

The effect of the probiotic on the increase of the digestibility is because the probiotic improves cellulolytic bacteria in the rumen [28] and increases the pH ruminal, consequently, increasing the fiber degradation [29]. Some LAB break carbohydrates and produce simple carbohydrates (monosaccharides) like glucose, which is essential to produce energy, then, the animal can make better use of the food, since improved the carbohydrates bioavailability [30].

Other LAB are reported to produce enzymes that help to digest the fiber carbohydrates, all of this improve animal performance, therefore, increasing the animal productivity [30, 31].

3.3 Production of volatile fatty acids (VFA's)

The acetic, propionic, and butyric acids are produced in higher concentrations during food fermentation in the rumen by the action of rumen bacteria. Therefore, it was thought that rumen probiotics could improve VFA production. However VFA concentrations remain similar to the control and were no observed increase with the addition of the probiotic or with the increase of the doses. Despite an increase in digestibility, the probiotic inoculation did not benefit the production of VFA ($P > 0.05$, data not showed). Even decreased one VFA, acetate with 4×10^{11} and 4×10^{12} doses of probiotic.

Our results are similar to other studies that report no significant changes in VFA concentrations by adding probiotics in the rumen. Ellis et al. [14] report that there were no significant effects of LAB on VFA concentration after 72 h of incubation at 37°C in rumen compared to the control. Jal et al. [32]

Comment [H14]: Follow the referencing guideline strictly

also reported no significant increase of VFA concentration by LAB compared to the control. **Quadis et al.** [33] report that a probiotic that contains *L. plantarum* strain 220, *E. faecium* strain 26 and *Clostridium butyricum* strain Miyari did not affect ruminal VFA concentrations. **Chiquette et al.** [34] neither observed an increase in VFA production using *Prevotella bryantii* 25A as a probiotic in cows.

4. CONCLUSIONS

In summary, the viability was greater in cryopreserved and lyophilized than retriggered in both probiotics. Nevertheless, lyophilized is easier to handle due to it being a dry powder, A1D42 lyophilized had better viability than A3D42 lyophilized, and this was maintained until 60 days. Moreover, A1D42 was shown to increase significantly the digestibility, which improved with increasing probiotic dosage up to 66%. The VFAs production did not augment the addiction to the probiotic.

Based on the results, it was concluded that lyophilization is a great conservation method for the production of rumen probiotics, which proved not to affect the properties and beneficial effects of the probiotic *in vitro*. Nevertheless, its preservation could be improved with the addition of antioxidants, with storage under vacuum at low temperature and controlling water activity.

ETHICAL APPROVAL

UNAM CICUAE.DC-2019/4-2

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Comment [H15]: Check the spelling

REFERENCES

1. Food and Agriculture Organization of the United Nations/World Health Organization (FAO/WHO). Guidelines for the evaluation of probiotics in food. 2002. Accessed 20 May 2022. Available: https://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf
2. Pineiro M, Stanton C. Probiotic bacteria: legislative framework-Requirements to evidence basis. *J Nutr.* 2007;137(3 Suppl 2):850S–853S.
3. Uyeno Y, Shigemori S, Shimosato T. Effect of probiotics/ prebiotics on cattle health and productivity. *Microbes Environ.* 2015;30(2):126–132.
4. Huyghebaert G, Ducatelle R, Van Immerseel F. An Update on Alternatives to Antimicrobial Growth Promoters for Broilers. *Vet J.* 2011;187(2):182-188.
5. Coppola R, Succi M, Tremonte P, Reale A, Salzano G, Sorrentino E. Antibiotic susceptibility of *Lactobacillus rhamnosus* strains isolated from Parmigiano Reggiano cheese. *Lait.* 2005;85(3): 193-204.
6. Serna-Cock L, Enríquez-Valencia CE, Jiménez-Obando EM, Campos-Gaona R. Effects of fermentation substrates and conservation methods on the viability and antimicrobial activity of *Weissella confusa* and its metabolites. *Electron J Biotechnol.* 2012;15(3):1-6.
7. Murga ML, Cabrera GM, De Valdéz GF, Disalvo A, Seldes AM. Influence of growth temperature on cryotolerance and lipid composition of *Lactobacillus acidophilus*. *J Appl Microbiol.* 2000;88(2):342-348.
8. Fowler A, Toner M. Cryo-injury and biopreservation. *Ann N Y Acad Sci.* 2005;1066:119–135.
9. Cárcoba R, Rodríguez A. Influence of cryoprotectants on the viability and acidifying activity of frozen and freeze-dried cells of the novel starter strain *Lactococcus lactis* ssp. *Eur Food Res Technol.* 2000;211:433-437.
10. Zayed G, Roos YH. Influence of trehalose and moisture content on survival of *Lactobacillus salivarius* subjected to freeze-drying and storage. *Process Biochem.* 2004;39:1081-1086.
11. Selmer-Olsen E, Birkeland SE, Sørhaug T. Effect of protective solutes on leakage from and survival of immobilized *Lactobacillus* subjected to drying, storage and rehydration. *J Appl Microbiol.* 1999;87(3):429-437.
12. Weinberg Z, Shatz O, Chen Y, Yosef E, Nikbahat M, Ben-Ghedalia D, Miron J. Effect of lactic acid bacteria inoculants on in vitro digestibility of wheat and corn silages. *J Dairy Sci.* 2007;90(10):4754-4762.
13. Ridwan R, Ariga BW, Dwi AW, Rohmatussolihat, Fitri SN, Fidriyanto R, Jayanegara A, Wijayanti I, Widyastuti Y. The Use of Lactic Acid Bacteria as Ruminant Probiotic Candidates Based on In Vitro Rumen Fermentation Characteristics. *Buletin Peternakan.* 2018;42(1):31-36.

14. Ellis JL, Bannink A, Hindrichsen IK, Kinley RD, Pellikaan WF, Milora N, Dijkstra J. The effect of lactic acid bacteria included as a probiotic or silage inoculant on *in vitro* rumen digestibility, total gas and methane production. *Anim Feed Sci Technol.* 2016;211:61-74.
15. Succi M, Tremonte P, Reale A, Sorrentino E, Coppola R. Preservation by freezing of potentially probiotic strains of *Lactobacillus rhamnosus*. *Ann Microbiol.* 2007; 57:537-544.
16. Polo L, Mañes-Lázaro R, Olmeda I, Cruz-Pio, LE, Medina A, Ferrer S, Pardo I. Influence of freezing temperatures prior to freeze-drying on viability of yeasts and lactic acid bacteria isolated from wine. *J Appl Microbiol.* 2017;122(6):1603-1614.
17. Girme G, Arosemena EL, Perez L, Vera R, Grau E, Calvo MA. Enhancement of the Viability of *Lactobacillus plantarum* during the Preservation and Storage Process Based on the Response Surface Methodology. *Food Nutr Sci.* 2014;5:1746-1755.
18. Huang L, Lu Z, Yuan Y, Lü F, Bie X. Optimization of a Protective Medium for Enhancing the Viability of Freeze-Dried *Lactobacillus delbrueckii* subsp. *bulgaricus* Based on Response Surface Methodology. *J Ind Microbiol Biotechnol.* 2006;33(1):55-61.
19. Fernández S, Fraga M, Silveyra E, Trombert AN, Rabaza A, Pla M, Zunino P. Probiotic properties of native *Lactobacillus* spp. strains for dairy calves. *Benef Microbes.* 2018;9(4):613-624.
20. Meng XC, Stanton C, Fitzgerald GF, Daly C, Ross RP. Anhydrobiotics: The challenges of drying probiotic cultures. *Food Chem.* 2008;106(4):1406-1416.
21. Tsvetkov T, Brankova R. Viability of micrococci and lactobacilli upon freezing and freeze-drying in the presence of different cryoprotectants. *Cryobiology.* 1983;20(3):318–323.
22. Brian CST, Etzel MR. Spray-drying, freeze-drying, or freezing of three different lactic acid bacteria species. *J Food Sci.* 1997;62(3):576–585.
23. Abadias M, Benabarre A, Teixidó N, Usall J, Viñas I. Effect of freeze drying and protectants on viability of the biocontrol yeast *Candida sake*. *Int J Food Microbiol.* 2001;65(3):173-182.
24. Bruno FA, Shah NP. Viability of two freeze-dried strains of **Bifidobacterium** and commercial preparations at various temperatures during prolonged storage. *J Food Sci.* 2006;68(7):2336–2339.
25. Simpson PJ, Stanton C, Fitzgerald GF, Ross RP. Intrinsic tolerance of *Bifidobacterium* species to heat and oxygen and survival following spray-drying and storage. *J Appl Microbiol.* 2005;99:493–501.
26. Coulibaly I, Kouassi EK, N'guessan E, Destain J, Béra F, Thonart P. Lyophilization (Drying Method) Cause Serious Damages to the Cell Viability of Lactic Acid Bacteria. *Annu Res Rev Biol.* 2018;24(4):1-15.
27. Teixeira P, Castro H, Kirby R. Evidence of membrane lipid oxidation of spray-dried *Lactobacillus bulgaricus* during storage. *Lett Appl Microbiol.* 1996;22(1):34–38.

28. Wallace RJ, Newbold CJ. Rumen fermentation and its manipulation: The development of yeast culture as feed additives. In: Lyons, TP, editor. *Biotechnology in the Feed Industry*. Alltech Technical Publications, Nicholasville, KY. 1993.
29. Umberger SH, Notter DR. Evaluation of lactobacillus inoculants on feedlot lamb performance. *J Anim Sci*. 1989;8:40-45.
30. Khalid MF, Shahzad MA, Sarwar M, Rehman AU, Sharif M, Mukhtar N. Probiotics and lamb performance: A review. *Afr J Agric Res*. 2011;6(23):5198-5203.
31. Fiems LO. The use of yeast in practical diets for ruminants. In: Castanon JIR, editor. *Microorganisms and enzyme preparations in animal nutrition*. Directorate-General for Agriculture, European Communion, Brussels. 1994.
32. Jalc D, Varadyova Z, Laukova A, Homolka P, Jancik F. Effect of inoculated corn silage on rumen fermentation and lipid metabolism in an artificial rumen (RUSITEC). *Anim Feed Sci Technol*. 2009;152(3-4):256-266.
33. Quadis AQ, Goya S, Ikuta K, Yatsu M, Kimura A, Nakanishi S, Sato S. Effects of a Bacteria-Based Probiotic on Ruminal pH, Volatile Fatty Acids, and Bacterial Flora of Holstein Calves. *J Vet Med Sci*. 2014;76(6):877-85.
34. Chiquette J, Allison MJ, Rasmussen MA. *Prevotella bryantii* 25A used 346 as a probiotic in early lactation dairy cows: Effect on ruminal fermentation characteristic 347 cs, milk production, and milk composition. *J Dairy Sci*. 2008;91(9):3536-3543.