

Original Research Article

UPTAKE AND SIDE EFFECTS PROFILE OF COVID-19 VACCINE AMONGST MEDICAL DOCTORS AT A TERTIARY HOSPITAL IN SOUTH-SOUTH NIGERIA

ABSTRACT

Aim: To determine the uptake of, and the incidence and pattern of side effects to the COVID-19 vaccine amongst medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria.

Study design: This was a descriptive cross-sectional study.

Place and duration of Study: This study was carried out among medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria, from 15th June to 30th July, 2021.

Methodology: Semi-structured questionnaires were used to obtain data from 102 participants recruited by consecutive sampling. Data obtained were the socio-demographic characteristics of the participants, the presence of co-morbidities and the presence of side effects to the COVID-19 vaccine for those vaccinated.

Result: Out of 102 medical doctors who participated in the survey, only 30 (29.4%) had received the COVID-19 vaccine and 26 of them (86.7%) reported different forms of side effects. Pain at the injection site was the commonest vaccine-associated side effect (76.7%). The first day after COVID-19 vaccination was the commonest time of onset of the reported side effects. There was no statistically significant relationship between COVID-19 vaccine side effects and presence of chronic medical conditions in the participants.

Conclusion: Uptake of COVID-19 vaccine among medical doctors was low and is worrisome. Incidence of COVID-19 vaccine side effects was high; however, the symptoms were mild to moderate. In the face of this challenging pandemic, COVID-19 vaccination still remains a safe means of ensuring protection against the disease.

KEYWORDS: COVID-19 vaccine, Side effects, Medical doctors, Gender, Co-morbidities

INTRODUCTION

COVID-19 is an illness caused by a new virus called severe acute respiratory syndrome Corona virus 2 (SARS-CoV-2). It was identified first in the middle of an outbreak of respiratory illnesses in Wuhan City, Hubei province, China.¹ It was reported to WHO on the 31st of December 2019, declared a global health emergency by WHO on the 30th of January 2020 and declared a global pandemic by WHO on the 11th of March 2020.²⁻⁴

COVID-19 has serious deleterious effect globally. As of 30th June 2021, there were 181,344,224 confirmed cases of COVID-19 including 3,934,252 deaths reported to WHO globally.⁵ Of these, Nigeria had 167,543 confirmed cases and 2,120 deaths as reported by Nigeria centre for disease control.⁶ This put so much strain on health and global economy with increased unemployment rate; as more man power, infrastructure, equipment, drugs, consumables are needed to combat this pandemic. This effect is more in developing countries including Nigeria that has an already fragile economy and health system.

Comment [L1]: The title is clear, but if you read the title of this research is descriptive, it turns out that from the method and results of the study, a hypothesis test was carried out on the relationship between the side effects of the Covid-19 vaccine and chronic medical conditions.

Comment [L2]: Not yet consistent with the results of research that proves the hypothesis test that there is no relationship between the side effects of the Covid-19 vaccine and chronic medical conditions

Comment [L3]:
Write down the most side effects

Since respiratory droplets carrying infectious virus is the principal mode of transmission, the following measures are employed to prevent the spread of COVID-19: washing of hands with soap and water or use of alcohol-based hand sanitizer; maintaining a safe distance from anyone who is coughing or sneezing; wearing of facemask; avoid touching eyes, nose and mouth; covering of nose and mouth with bent elbow or a tissue when coughing or sneezing; staying at home when feeling unwell; and seeking medical attention when having fever, cough, or difficulty in breathing.⁷

While observing these preventive measures, vaccination is key to ending this pandemic. There are about 52 candidate vaccines in human trials as at December 2020.⁸ Pfizer/ bioNTech vaccine was the first to receive emergency validation by the WHO on the 31st of December, 2020.⁹ Others approved by WHO for emergency use include: AstraZeneca/Oxford vaccine, Johnson and Johnson vaccine, Moderna, Sinopharm, and Sinovac.¹⁰

Nigeria received her first batch of COVID-19 vaccine (AstraZeneca/Oxford vaccine) on the 2nd of March 2021.¹¹ As of 29th June, 2021, Two million, Two Hundred and Ninety-nine Thousand, Nine Hundred and Eighty-seven (2,299,987) people have received their first dose of COVID-19 vaccine while 1,209,591 people have received their second dose in Nigeria.¹² Bayelsa state flagged off COVID-19 vaccination on 12th of March 2021.¹³

Like other vaccines, COVID-19 vaccines are associated with some side effects which are mostly mild to moderate in severity and short lasting.^{14,15} This include pain, redness and swelling at injection site; fever; fatigue; headache; muscle pain; chills; nausea; vomiting; diarrhoea; mild flu-like symptoms; joint pain; weakness; loss of appetite; malaise; cough; and facial swelling.^{15,16} Others include transverse myelination, disorientation, hypersensitivity to light and noise, personality behaviour change. Severe allergic reactions have occurred though rarely to some COVID-19 vaccines.¹⁴⁻¹⁶ Severe allergic reaction such as anaphylaxis is a potential but rare side effect with any vaccine.¹⁶

Medical personnel, the military and other essential workers in Nigeria were among the first to be given the vaccine because of the limited supply of the vaccine to the country.¹⁶ It would be necessary to study the side effect profile of COVID-19 vaccine amongst vaccinated medical doctors because they would be in a good position to tell their experiences to other health care workers as well as their public. This study would also act as a reference tool for further research on COVID-19 vaccine uptake and side effects in Africans.

The study aimed at determining the uptake of, and the incidence and pattern of side effects to the COVID-19 vaccine amongst medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria.

METHODOLOGY

This was a descriptive cross-sectional study carried out among medical doctors at the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria, from 15th June to 30th July, 2021. The hospital is one of the two tertiary hospitals in the state. It resides within the suburbs of the state and 0.69 km from the state capital, Yenagoa. There are approximately 154 medical doctors working at the facility.

Using the formula for calculating sample size in cross-sectional studies¹⁷ and based on a prevalence of COVID-19 vaccination among health workers of 27.4% from another study in Yenagoa,¹⁸ a sample size of 102 was derived for this study after adjusting for an anticipated non-response rate of 30%. A semi-structured questionnaire was used to obtain data from 102 participants by consecutive sampling.

RESULTS

Comment [L4]: Add research objectives to find out relationship between the side effects of the Covid-19 vaccine and chronic medical conditions. Consistent with research methods and results

Comment [L5]: Explain what data is collected, how the data is collected, the classification of each research variable, the statistics used (descriptive and inferential)

Socio-demographic characteristics of participants

A total of one hundred and two medical doctors from the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa state, Nigeria participated in the survey. Of the 102 participants, about three-fifths were males (62.7%) while females were 37.3% (Table1). Thirty-nine (38.2%), 45 (44.1%), and 12 (11.8%) medical doctors were in the third, fourth and fifth decades of life respectively (Table 1). Majority of the participants were single (62.7%) and Christians by religion (93.1%).

Chronic medical conditions among participants

Chronic non-communicable diseases (NCD) were reported among seventeen participants (16.7%). The NCDs included hypertension (11.8%), diabetes mellitus (2.9%) and asthma (2.9%). Other conditions (2.9%) reported were chronic inflammatory synovitis and rheumatoid arthritis (Table 2).

Table 1: Socio-demographic characteristics of participants

Characteristics	Frequency N = 102	Percent (%)
Sex		
Male	64	62.7
Female	38	37.3
Age group		
20 - 29 years	39	38.2
30 - 39 years	45	44.1
40 - 49 years	12	11.8
> 50 years	6	5.9
Marital Status		
Single	64	62.7
Married	38	37.3
Religion		

Christian	95	93.1
Others	7	6.9

Table 2: Chronic medical conditions among respondents

Characteristics	Frequency N = 102	Percent (%)
Chronic medical condition		
No chronic medical condition	85	83.3
Chronic medical condition present	17	16.7
Type of chronic medical condition		
Hypertension	12	11.8
Diabetes	3	2.9
Asthma	3	2.9
Sickle Cell Disease	1	1.0
Others*	3	2.9

*Others – Allergic drug reaction, rheumatoid disease, chronic inflammatory arthritis

COVID 19 vaccine uptake and reported side effects among respondents

Thirty (30) out of 102 medical doctors (29.4%) had taken the vaccine about 3 months after it was provided at the hospital for vaccination of medical personnel(Figure 1).

Of the 30 medical doctors who had taken the vaccine, 26 of them (86.7%) reported different forms of side effects (Table 3). The most common reported side effect was pain at the injection site, which was reported in 23 participants (76.7%). Other side effects included fever (60.0%), muscle and joint pain (46.7%), fatigue (40.0%) and headache (40.0%). Most of those (46.7%) who reported side effects reported it on day 1 after the injection (Table 3).




Figure 1: Vaccine uptake among Participants

Comment [L6]:
Writing decimal numbers use the notation "dot" not "comma"

Table 3: Reported side effects among respondents.

Side effects	Frequency N = 30	Percent (%)
Side effects reported	26	86.7
No side effect reported	4	13.3
Reported Site effects		
Pain in injection site	23	76.7
Fever	18	60.0
Muscle and joint pain	14	46.7
Fatigue	12	40.0
Headache	12	40.0
Chills	9	30.0
Nausea	3	10.0
Vomiting	1	3.3
Allergy	1	3.3

Day COVID-19 Side effects started

Comment [L7]: Total percentage not 100%

Day 1	14	46.7
Day 2	11	36.7
Day 3 to 7	1	3.3

Association between chronic medical conditions and vaccine side effects

There was no significant difference ($p > 0.05$) in the occurrence of side effects among participants who reported chronic medical conditions and those who did not (Table 4)

Table 4: Association between chronic medical conditions and vaccine side effects

Characteristic	Total N = 30 (%)	Chronic medical condition		Chi square	P-value
		Present N = 3 (%)	Absent N = 27 (%)		
Side effect reported	26 (86.7)	3 (100.0)	23 (85.2)	0.37	.54
No side effect reported	4 (13.3)	0 (0.0)	4 (14.8)		
Pain at injection site	23 (76.7)	3 (100.0)	20 (74.1)	1.01	.31
No Pain at injection site	7 (23.3)	0 (0.0)	7 (25.9)		
Fever	18 (60.0)	3 (100.0)	15 (55.6)	2.22	.26
No fever	12 (40.0)	0 (0.0)	12 (44.4)		
Muscle/Joint pain	14 (46.7)	1 (33.3)	13 (48.1)	0.24	.63
No muscle/Joint pain	16 (53.3)	2 (66.7)	14 (51.9)		
Fatigue	12 (40.0)	1 (33.3)	11 (40.7)	0.06	.80

Comment [L8]: For all p-values add 0 in front of the dot

No Fatigue	18 (60.0)	2 (66.7)	16 (59.3)		
Headache	12 (40.0)	2 (66.7)	10 (37.0)	0.99	.55
No headache	18 (60.0)	1 (33.3)	17 (63.0)		
Chills	9 (30.0)	2 (66.7)	7 (25.9)	2.13	.53
No Chills	21 (70.0)	1 (33.3)	20 (74.1)		
Vomiting	1 (3.3)	0 (0.0)	1 (3.7)	0.12	.74
No Vomiting	29 (96.7)	3 (100.0)	26 (96.3)		
Nausea	3 (10.0)	0 (0.0)	3 (11.1)	0.37	.54
No nausea	27 (90.0)	3 (100.0)	24 (88.9)		

DISCUSSION

COVID-19 vaccine is that veritable tool in the war against COVID-19 pandemic, needed to break the chain of transmission of the infection and establish herd immunity, but its uptake is hindered by the fear of its side effects. COVID-19 vaccine side effects are yet to be well understood as the disease and its vaccine are novel.

More males than females participated in this study. This was due to the fact that more males than females presently work at the facility. Most of the participants were single and Christians as the study was carried out in a Christian-dominated locality.

From this study, most of the medical doctors had no chronic medical condition while the nature of chronic medical conditions reported by 16.7% of them was predominantly hypertension and diabetes. This is in keeping with similar studies done in Bayelsa, Lagos, China and Saudi Arabia respectively.¹⁹⁻²² Presence of co-morbidities predisposes to adverse clinical outcomes in patients with COVID-19 and may result in greater disease severity when compared with those with no co-morbidity.²¹ Thus, approved and authorized COVID-19 vaccines are especially important in people with underlying medical conditions.

COVID-19 vaccine uptake among medical doctors was low in this study; only 29.4% of the respondents took the vaccine. This low uptake rate is congruent with 27.7% recorded in Congo,²³ 37.4% in Jordan²⁴ and 39.3% in Ghana.²⁵ It is however not in consonance with 64.9%

Comment [L9]: There is a need for justification why the coverage of covid-19 vaccination for doctors is low, and there is a referral that strengthens it.

Comment [L10]: No reference yet

and 86.2% recorded in Saudi Arabia and China respectively.^{26,27} The low uptake rate among medical practitioners in this study is worrisome as it will likely affect the uptake of the vaccines by their patients. Maintaining confidence in vaccination depends on the interaction between patients and providers.²³ Attitude and utilization of vaccination by healthcare professionals (HCP) is a major factor that is consistently associated with patient acceptance and vaccination, adherence to vaccination schedules, and reduced hesitation/aversion.²⁸ In addition, vaccinated HCPs also have a noticeable effect on patients' decision to take a vaccine. The low uptake rate among the top echelon of the health care team may be related to the side effects of the vaccine and its overall safety. Thus, more vaccine clinical trials have to be done to provide more convincing data about the safety of this vaccine and settle all controversies about the drug once and for all.

The finding that 86.7% of those who took the vaccine in this study reported different forms of side effects is similar to studies done by Khaleduzzaman et al²⁹ and Oriji et al¹⁹, however slightly higher than the rate in studies done by Hatmal et al³⁰ and Menni et al.³¹ The most common reported side effect of pain at the injection site from this study was in consonance with a similar study among health workers in a sister facility by Oriji et al.¹⁹ It however disagrees with other studies.^{29,30} Other side effects include fever (60.0%), muscle and joint pain (46.7%), fatigue (40.0%) and headache (40.0%). Vaccines are designed to give immunity without the dangers of getting the disease however developing mild to moderate side effects is common. This is because the immune system is instructing the body to react in certain ways increasing blood flow so more immune cells can circulate, and it raises the body temperature in order to kill the virus. Mild-to-moderate side effects, like a low-grade fever or muscle aches, are normal and not a cause for alarm: they are signs that the body's immune system is responding to the vaccine, specifically the antigen and is gearing up to fight the virus. These side effects usually last for a few days.^{31,32}

In this present study, most of the side effects (46.7%) were reported within the first day of receiving the vaccine. This is consistent with several other studies²⁹⁻³² but inconsistent with a study done in Bangladesh²⁹ where most symptoms were reported after the third day of vaccination. Spike proteins are viral protein found on the surface of viruses and also apply to vaccines which follow the natural process of viral infection to mount an immune response. An immediate response is elucidated by the innate immune system initiating an inflammatory response resulting in fever and pain at injection site while the adaptive immunity which produces T cells and antibodies provides protection on subsequent exposure hence a delayed response.

From this study, there was no association between COVID-19 vaccine side effects and presence of chronic medical conditions. Although side effects were reported among all participants with chronic medical condition in this study, it is likely due to the small number of vaccinated respondents and was not of statistical significance. This is consistent with the knowledge that COVID-19 vaccines may be administered to most people with underlying medical conditions.¹

Experiencing side effects after getting vaccinated means the vaccine is working and the immune system is responding as it should.¹⁶ After vaccination, it usually takes a few weeks for the body to build immunity against SARS-CoV-2. So, it's possible a person could be infected with SARS-CoV-2 just before or after vaccination and still get sick with COVID-19. This is because the vaccine has not yet had enough time to provide protection.

CONCLUSION

Comment [L11]: Need an explanation why the relationship is not significant and other references need to be added

Uptake of COVID-19 vaccine among medical doctors was low and is worrisome, as it is likely to affect the uptake of the vaccines by their patients. Incidence of COVID-19 vaccine side effects was high; however, the symptoms were mild to moderate. The most common side effect was pain at injection site, observed especially within 24 hours of vaccination. In the face of this challenging pandemic, COVID-19 vaccination still remains a safe means of ensuring protection against the disease.

ETHICAL APPROVAL

This was obtained from the Research and Ethics Committee of the Niger Delta University Teaching Hospital, Okolobiri, Bayelsa.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. CDC 2019 Novel Coronavirus, Wuhan, China. CDC. Published January 26, 2020. Accessed 2 August 2021. Available: <https://www.cdc.gov/coronavirus/2019-ncov/about/index.html>. 2020.
2. Gallegos A. WHO Declares Public Health emergency for novel coronavirus. Medscape Medical News. Published January 30, 2020. Accessed 2 August 2021. Available: <https://www.medscape.com/viewarticle/924596>. Published January 30, 2020.
3. Ramzy A, McNeil NG. W.H.O. Declares Global Emergency as Wuhan Coronavirus Spread. The New York Times. January 30, 2020. Accessed 2 August 2021. Available: <https://nyti.ms/2RER70M>.
4. Coronavirus Live Updates: W.H.O Declares Pandemic as Number of Infected Countries Grows. The New York Times. March 11, 2022 Accessed 4 August 2021. Available: <https://www.nytimes.com/2020/03/11/world/coronavirus-news.html#link-682e5b06>.
5. WHO Coronavirus (COVID 19) Dashboard. World Health Organization. Accessed 13 September 2021. Available: <https://covid19.who.int>.
6. Covid-19 Nigeria. Nigeria Centre for disease control. Accessed 13 September 2021. Available: <https://covid19.ncdc.gov.ng>.
7. Coronavirus Disease (COVID19) Advice for the Public. WHO. Accessed 13 September 2021. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>.
8. The different types of COVID-19 vaccines. World Health Organization. Accessed 14 September 2021. Available: <https://www.who.int/news-room/features-stories/detail/the-race-for-a-covid-19-vaccine-explained>.
9. WHO issues its first emergency use validation for a COVID-19 vaccine and emphasizes need for equitable global access. World Health Organization. Accessed 30 September 2021. Available: <https://www.who.int/news/item/31-12-2020-who-issues-its-first-emergency-use-validation-for-a-covid-19-vaccine-and-emphasizes-need-for-equitable-global-access>.

Comment [L12]: It is necessary to write down the ethical approval number and an explanation that the respondent has received informed consent before data collection

Comment [L13]: Some of his written references are incomplete and inconsistent. Complete the reference meta data, journal name, volume, number, page, DOI

Comment [L14]: Need to add a primary reference 80% of the total reference

10. WHO validates sinovac COVID- 19 vaccine for emergency use and issues interm policy recommendations. World Health Organization. Accessed 21 August 2021. Available: <https://www.who.int/news/item/01-06-2021-who-validates-sinovac-covid-19-vaccine-for-emergency-use-and-issues-interim-policy-recommendations>.
11. WHO Africa. COVID- 19 vaccines shipped by COVAX arrive in Nigeria. World Health Organization Nigeria. Accessed August 30 2021. Available: <https://www.afro.who.int/news/covid-19-vaccines-shipped-covac-arrive-nigeria>.
12. COVID- 19 Vaccination Update. National Primary Health Care Development Agency. Accessed 30 August 2021 Available: <https://m.facebook.com/NPHCDA/posts/4451541584870460>.
13. Bayelsa Begins Covid-19 Vaccination, Deputy Governor, Orthers Vaccinated. Ministry of Health News. Accessed 29 August 2021. Available: <https://mohbayelsa.gov.ng/bayelsa-begins-covid-19-vaccination-deputy-governor-others-vaccinated/>.
14. Possible Side effects After Getting a COVID-19 Vaccine. Centre for Disease Control and Prevention. Updated June 21, 2021. Accessed 20 August 2021. Available: <https://www.cdc.gov/coronavirus/2019-ncov/vaccine/expect/after.html>.
15. Poole J. Covid vaccine side effects. Nemeton Research Foundation. Accessed 20 August 2021. Available: <https://www.researchgate.net/publications/348327013>.
16. Side Effects of COVID-19 Vaccines. WHO/ World Health Organization. Accessed 20 August 2021. Available: <https://www.who.int/news-room/feature-stories/detail/side-effects-of-covid-19-vaccines>.
17. Oyidarin OT, Usman SA, Osoba ME, Olukorode SO, Lucero-Prisno III DE. Towards effective and efficient covid-19 vaccination in Nigeria. *Journal of Global Health Reports*. 2021; 5e2021023. doi:10.29392/001c.21404.
18. Araoye MO. Subject Selection. In: Research Methodology with statistics for Health and Social sciences. Ilorin, Nigeria: Nathedex Publishers, 2003: 115-129.
19. Orij PC, Allogoa DO, Wagio TJ, Obagah L, Tekenah ES, Ozori ES. Hesitancy of COVID-19 Vaccination among Health Workers (other than Doctors) in a Tertiary Health Hospital in South-South, Nigeria. *Asian Journal of Research in Infectious Diseases*, 2021;7(1):21-31.
20. Orij PC, Allagoa DO, Obagah L, Oguiche OI, Ohaeri OS, Tekenah ES et al. Side Effect Profile of COVID-19 Vaccine among Health Workers in a Tertiary Health Institution in South-South Nigeria. *European Journal of Medical and Health Sciences*. 2021; 3(3): 59-64.
21. Osibogun A, Balogun M, Abayomi A, Idris J, Kuyinu Y, Odukoya O, et al. Outcomes of COVID-19 patients with comorbidities in southwest Nigeria. *PLoS One*.2021;16(3): e0248281. Published 2021 Mar 15. doi:10.1371/journal.pone.0248281
22. Guan WJ, Liang WH, Zhao Y, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: a nationwide analysis. *Eur Respir J* 2020; 55(5):2000547. Published 2020 May 14. doi:10.111183/1393003.00547-2020.
23. Fathi M, Vakili K, Sayehmiri F, Mohamadkhani A, Hajjesmaeili M, Rezaei-Tavirani M. The prognostic value of comorbidity for the severity of COVID-19: A systematic review and meta-analysis study. *PLoS ONE*. 2021;16(2):e0246190. Published 2021 Feb 16. doi:10.1371/journal.pone.0246190.
24. Nzaji MK, Ngombe LK, Mwamba GN, Ndala DB, Miema JM, Lungoyo CL, et al. Acceptability of Vaccination Against COVID-19 Among Healthcare Workers in the Democratic Republic of the Congo. *Pragmatic and Observational Research*. 2020; 11: 103-110.
25. El-Elimat T, AbuAISamen MM, Almomani BA, Al-Sawalha NA, Alali FQ. Acceptance and attitudes toward COVID-19 vaccines: A cross-sectional study from Jordan. *PLoS One*. 2021;16(4): e0250555. Published 2021 Apr 23. doi:10.1371/journal.pone.0250555.

26. Agyekum MW, Afrifa-Anane GF, Kyei-Arthur F, Addo B. Acceptability of COVID-19 Vaccination among Health Care Workers in Ghana: *Advances in Public Health*, vol. 2021:1-8. Published 2021 Apr 20. Accessed 20 October 2021. Available: <https://doi.org/10.1155/2021/9998176>.
27. Elharake JA, Galal B, Alqahtani SA, Kattan RF, Barry MA, Temsah M, et al. COVID-19 Vaccine Acceptance among Health Care Workers in the Kingdom of Saudi Arabia. *International Journal of Infectious Diseases*. 2021;109: 286–293.
28. Xu B, Gao X, Zhang X, Hu Y, Yang H, Zhou Y-H. Real-World Acceptance of COVID-19 Vaccines among Healthcare Workers in Perinatal Medicine in China. *Vaccines*. 2021; 9(7):704. Published 2021 Jun 27. doi.org/10.3390/vaccines9070704.
29. Asma S, Akan H, Uysal Y, et al. Factors affecting influenza vaccination uptake among health care workers: a multi-center cross-sectional study. *BMC Infect Dis*. 2016;16:192. Published 2016 May 4. doi:10.1186/s12879-016-1528-9.
30. Khaleduzzaman HM, Nafisa Jabin Mishu NJ. Frequency of Side Effects after First Dose of Vaccination against COVID-19. Among the People of Bangladesh. *European Journal of Medical and Health Sciences*. 2021;3(3):22-24.
31. Hatmal MM, Al-Hatamleh MA, Olaimat AN, Hatmal M, Alhaj-Qasem DM, Olaimat TM et al. Side Effects and Perceptions Following COVID-19 Vaccination in Jordan: A Randomized Cross-Sectional Study Implementing Machine Learning for Predicting Severity of Side Effects. *Vaccines*. 2021;9(6):556. doi.org/10.3390/vaccines9060556.
32. Menni C, Klaser K, May A, Polidori L, Capdevila J, Louca P et al. Vaccine side-effects and SARS-CoV-2 infection after vaccination in users of the COVID Symptom Study app in the UK: a prospective observational study. *The Lancet Infectious Diseases*. 2021; 21(7): 939-949.
33. Djanas D, Yusirwan, Martini RD, Rahmadian, Putra H, Zanir A et al. Survey data of COVID-19 vaccine side effects among hospital staff in a national referral hospital in Indonesia. *Data Brief*. 2021;36:107098. doi: 10.1016/j.dib.2021.107098.

Comment [L15]: Many references are incomplete Meta data