

EFFECT OF REPLACING BONE ASH WITH FRESH WATER SNAIL (*Pila ampullacea*) SHELL ASH ON THE PERFORMANCE OF WEANER RABBITS.

Comment [H1]: No dot

Abstract

Twenty five (25) weaned male rabbits of mixed breeds were used to investigate the effect of replacing bone ash with fresh water snail (*Pila ampullacea*) shell ash on the performance of weaned rabbits. The experiment was conducted at the Teaching and Research Farm, University of Agriculture, Makurdi, Benue State, Nigeria for a period of 12 weeks. The rabbits were randomly assigned to the five dietary treatments and replicated five times giving a total of one rabbit per replicate in a completely randomized design (CRD). Rabbits were raised intensively in hutches of about 4ft × 4ft. Five experimental diets tagged T₁ to T₅ were formulated -such that fresh water snail (*Pila ampullacea*) shell ash replaced bone ash at 0%, 25%, 50%, 75% and 100%. Feed and water were served *ad libitum*. Bio sanitary and bio security measures were strictly adhered to. The study showed that the growth parameters were not affected (P=.05) by the dietary treatments. This study had shown that fresh water snail (*Pila ampullacea*) shell ash can serve as a substitute for bone ash in weaned rabbits diet up to 100% without -compromising their performances.

Comment [H2]: This can be omitted

Comment [H3]: This can be removed

Comment [H4]: Too?

Comment [H5]: This can be removed

Key Words: Rabbit, *fresh water snail shell ash, bone ash, performance indices.*

Comment [H6]: Check italic use

Comment [H7]: No dot

INTRODUCTION

Animal protein consumption in Nigeria is about 7.4 g per capita/day [1], instead of the 28 g per person per day recommended by the [2]. The wide gap between the estimated protein requirement and the actual protein consumed is as a result of poor nutritional situation in Nigeria [3]. This indicates the need to intensify livestock production. Viable options need to be explored and evaluated in order to achieve this objective [4]. The use of livestock species that are yet to play a major role in animal production is among such alternative. Rabbit production has the potentials of alleviating animal protein deficiency in Nigeria [5].

Comment [H8]: Please identify the reference (authors name)

The rabbit has great potentials and good attributes such as; fast growth rate, high efficiency in converting forage to meat, short generation interval, high prolificacy, relatively low cost of production, and high nutritional quality of rabbit meat which includes; low fat, sodium and cholesterol levels. Rabbit meat contains about 20.8% of protein and its consumption is bereft of cultural and religious biases [6].

There is an urgent need to explore the potentials of non-conventional calcium sources that do not compete with human for food. One of such alternative feedstuff that is cheap, locally available, less competitive with humans and livestock is the fresh water snail (*Pila ampullacea*) shells. *Pila ampullacea* in river Benue and its tributaries has been studied by [7]. The abundance of fresh water snails is as a result of availability of food, shelter and oviposition sites along the said river water body. One of the most successful methods of catching is hand picking done as a community effort on regular basis. Snail shell is a mineral ingredient that contains about 98% of calcium carbonate [8]. It is therefore a biological source of calcium that can be used in animal feeding. Investigations have been done on the use of many sources of calcium such as gypsum, limestone and oyster shell in layers and broilers diets [9] but there is a dearth information on the use of snail shells especially fresh water snail (*Pila ampullacea*) shells in animal feeds.

Comment [H9]: Can be substituted by 'fresh water snail'

Comment [H10]: Please identify the name of author

Comment [H11]: Already explained, This can be removed and check throughout manuscript

MATERIALS AND METHODS

Experimental Site

The experiment was conducted in the rabbitry unit at the Teaching and Research Farm, College of Animal Science University of Agriculture Makurdi Benue State. Benue State lies within the lower River Benue trough in the middle belt region of Nigeria. Its geographic coordinates are longitude 7° 47' and 10° 0' East. Latitude 6° 25' and 8° 8' North [10]

Comment [H12]: Space adds

Source of Bone Ash

Comment [H13]: Of?

Bone ash was bought at God 4 us livestock consult, beside SRS junction, new bridge road, north bank Makurdi.

Sources and Collection of Freshwater Snail (*Pila ampullacea*) Shells

Comment [H14]: Add space between fresh and water

Comment [H15]: remove

Freshwater snails are in abundance in River Benue and its tributaries. The test ingredient was sourced locally at Gbajimba and Iyeh in Guma Local Government Area and Makurdi metropolis, the flesh is usually removed and the shells are thrown away by the consumers.

Comment [H16]: Add space between fresh and water

Processing of Fresh Water Snail (*Pila ampullacea*) Shell

Comment [H17]: remove

Fresh water snail (*Pila ampullacea*) shell were collected between the months of February and March. The shells were thoroughly washed, dried and burnt for about 1 hour until they became whitish in appearance; they were then crushed into fine powder as shell ash and used in the diet. The mineral composition of the shell was analyzed by the procedure of Association of Official Analytical Chemists (AOAC) [11].

Comment [H18]: remove

Comment [H19]: correct this

Comment [H20]: How many oC?

Comment [H21]: Add space

Table 1: Mineral Composition of Fresh Water Snail (*Pila Ampullacea*) Shell Ash .

Comment [H22]: Please check font type and size throughout table, it's not consistent

Parameter	% composition
Ash	92.45
Calcium	41.60
Phosphorus	0.01

Pasa = Pila ampullacea shell ash

Experimental Design

A Completely Randomized Design (CRD) was used for this experiment. A total of twenty five (25) weaned male rabbits of mixed breeds at five weeks of age with an initial average weight of about 664.00 – 667.00g were obtained from Dagwom Farm, National Veterinary Research Institute (NVRI) Vom, Jos Plateau State for the research. The rabbits were allowed for a preliminary feeding period of seven days to enable them acclimatize after which they were randomly assigned to five (5) dietary treatments designated as T₁ to T₅. Each of the dietary treatments had five (5) rabbits with each rabbit serving as a replicate.

Comment [H23]: This can be omitted

Comment [H24]: This can be omitted

Comment [H25]: This can be omitted

Housing and Management of Experimental Animals

The rabbits were housed individually in the hutches and labeled according to the treatment and replicate assigned to them. The dimension of the hutches was 40cm × 30cm × 30cm, (this was to enable it accommodate the feeders and drinkers). The initial weight of each rabbit was taken before assigning them to one of the five dietary treatments. Prophylactic medication was given against any infection before the commencement of the experiment. Each rabbit was observed daily to ensure good health. A known quantity of diet that will ensure the birds feed *ad-libitum* was served daily to each replicate, left over feed was weighed daily and quantity consumed was determined by the difference. Fresh clean water was also provided every morning. The experiment lasted for 12 weeks.

Comment [H26]: Please add space

Experimental Diets

Five experimental diets were formulated tagged T₁ to T₅. T₁ served as a control diet. Fresh water snail (*Pila ampullacea*) shell ash replaced bone ash at 0%, 25%, 50%, 75% and 100% respectively. T₁ containing 100% bone ash while T₅ contained 100% fresh water snail (*Pila ampullacea*) shell ash. These were mixed with other ingredients as presented in Table 2 below;

Comment [H27]: remove

Comment [H28]: remove

Table 2: Composition of Experimental Diets with Fresh Water Snail (*Pila ampullacea*) Shell Ash as a Replacement for Bone Ash (%)

Feedstuff	Experimental Diets				
	(0%Pasa)	(25%Pasa)	(50%Pasa)	(75%Pasa)	(100%Pasa)

Maize	30.00	30.00	30.00	30.00	30.00
Full fat soyabean	20.00	20.00	20.00	20.00	20.00
Groundnut cake	12.00	12.00	12.00	12.00	12.00
Maize offal	14.00	14.00	14.00	14.00	14.00
Rice offal	20.05	20.05	20.05	20.05	20.05
Bone ash	3.00	2.25	1.50	0.75	0.00
Pasa	0.00	0.75	1.50	2.25	3.00
Methionine	0.20	0.20	0.20	0.20	0.20
Lysine	0.20	0.20	0.20	0.20	0.20
Table Salt	0.30	0.30	0.30	0.30	0.30
Vit/min. premix	0.25	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00	100.00

Analysed Nutrient:

(%).Crude protein 16.86 16.86 16.86 16.86 16.86

M.E (Kcal/kg) 2637.10 2637.10 2637.10 2637.10 2637.10

Comment [H29]: add table line

Note: Pasa = Pila ampullacea shell ash

Performance Parameters

The parameters evaluated during the trial include; Initial weight (g), Final body weight gain (g), Average daily feed intake (g), Average weekly weight gain (g), Feed efficiency and Protein efficiency ratio.

Average Daily Feed Intake (ADFI) (g)

The rabbits in each treatment were fed weighed quantity of their group diets daily, feed intake was determined by obtaining the differences between the quantity of feed offered and the left over. Average daily feed intake was determined by dividing the total feed intake of the rabbits by the number of the experimental day

$$ADFI = \frac{\text{Total feed intake (g)}}{\text{Number of experimental days}}$$

No of days

Comment [H30]: Check line spacing

Average Daily Weight Gain (ADWG) (g)

The initial weight of the rabbits was recorded and they were then weighed on daily basis, therefore, the average daily body weight gain was determined using the formula below:

$$\text{ADWG} = \frac{\text{Final weight} - \text{Initial weight (g)}}{\text{No. of days}}$$

Feed Efficiency (FE)

This was determined by dividing feed intake by the weight gain

$$\text{FE} = \frac{\text{Feed intake (g)}}{\text{Weight gained (g)}}$$

Final Body Weight (g)

The final body weight of the rabbits was determined by weighing the rabbits at the end of the experiment, it was recorded in grams (g).

Protein Efficiency Ratio (PER)

The weight gain per unit of protein consumed was calculated using the formula below:

$$\text{PER} = \frac{\text{Weight gained(g)}}{\text{Gram of protein taken}}$$

Statistical Analysis

The data collected were subjected to one way Analysis of Variance (ANOVA) using Minitab statistical software version [16][12]. The separation of means was effected using Duncan's Multiple Range Test (DMRT) as outline by [13].

Comment [H31]: add space

Comment [H32]: Please specify the name of author

RESULTS AND DISCUSSION

The result of the proximate analysis of fresh water snail (*Pila ampullacea*) shell ash is presented in Table 2. It was observed that fresh water snail (*Pila ampullacea*) shell ash contained high percentages of ash (92.45%) and calcium (41.60%) but has negligible amount of phosphorus (0.01%) when compared to bone ash. The result showed that fresh water snail (*Pila ampullacea*) shell ash is a potential source of calcium in livestock feed.

Comment [H33]: remove

Comment [H34]: remove

Comment [H35]: remove

The nutrient composition of the five experimental diets is shown on Table 3. The findings revealed that the proximate fractions were not different from each other significantly. The diets were observed to have met the nutritional requirements for weaned rabbits across treatments.

Comment [H36]: in

The result of average daily feed intake was not significantly ($P>0.05$) affected by the dietary treatments. The study recorded daily feed intake values of 84.95 – 90.32 g, which were generally within the concentrate feed intake ranges of 61.08 to 133.9 g earlier reported for rabbits [16]. The high dietary inclusion levels of fresh water snail (*Pila ampullacea*) shell ash did not affect feed intake.

Comment [H37]: check

The average final weight recorded in this study ranged from 2163.40 to 2237.00(g) and were not significantly ($P>0.05$) affected by the dietary treatments. The result indicated that fresh water snail (*Pila ampullacea*) shell ash did not influence the final weight of the rabbits. This was in line with the findings of [15] who asserted that oyster shell and gypsum are sources of calcium which aids metabolic activities and not principal nutrient for growth in broiler chickens.

Comment [H38]: g

Comment [H39]: remove

Comment [H40]: Please specify the name of author

The average daily body weight gain in this finding showed no significant differences ($P>0.05$) among the treatment groups. The ADWG of the rabbits ranged from 17.82 to 18.70(g). This agrees with [15] who reported that oyster shell and gypsum are sources of calcium which aids metabolic activities and not principal nutrient for growth in broiler chickens. This study revealed that fresh water snail (*Pila ampullacea*) shell ash did not influence the final weight and average daily weight gain of rabbits.

Comment [H41]: g

Comment [H42]: Please specify the name of author

Comment [H43]: remove

Mortality	0.00	0.00	0.00	0.00	0.00	0.00

- Formatted: Font: 11 pt
- Formatted: Font: 11 pt
- Formatted: Font: 11 pt
- Formatted: Font: 11 pt
- Formatted: Font: 11 pt
- Formatted: Font: 11 pt
- Formatted: Expanded by 0.05 pt
- Formatted: Condensed by 0.5 pt

SEM= Standard Error of Means; NS= Not Significantly Different (P=.05); Pasa = Pila ampullacea shell ash
ADWG=Average daily weight gain; ADFI= Average daily feed intake
P E R = P r o t e i n e f f i c i e n c y

C O N C L U S I O N A N D R E C O M M E N D A T I O N S

C o n c l u s i o n

There were no significant (P=.05) differences across dietary treatments in all the performance parameters investigated. It is therefore concluded that fresh water snail shell ash can replace bone ash up to 100% without any adverse effect on the performance of weaned rabbits.

R E C O M M E N D A T I O N

It was recommended that supplemental levels of Phosphorus be included in weaned rabbit's diet.

C O M P E T I N G I N T E R E S T S D I S C L A I M E R :

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather

it was funded by personal efforts of the authors. *SEM= Standard Error of Means; NS= Not Significantly Different (P=.05); Pasa = Pila ampullacea shell ash*

- Formatted: Character scale: 58%, Not Expanded by / Condensed by

ADWG=Average daily weight gain; ADFI= Average daily feed intake

P E R = P r o t e i n e f f i c i e n c y

C O N C L U S I O N A N D R E C O M M E N D A T I O N S

C o n c l u s i o n

There were no significant ($P=0.05$) differences across dietary treatments in all the performance parameters investigated. It is therefore concluded that fresh water snail shell ash can replace bone ash up to 100% without any adverse effect on the performance of weaned rabbits.

E T H I C A L A P P R O V A L

"All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

R E F E R E N C E S

1. F.A.O. (Food and Agricultural Organization). 2006. Statistical Data base. Rome, Italy. Pp. 234-236.
2. British Medical Association, 1993. *Basic food and nutrient requirement for Africa*. In the World Book of Encyclopedia International, 7: 310-311.
3. Nworgu, F. C., Adebawale, E. A., Oredin, O. A. and Oni, A. 1999. Prospects and Economic of Broiler Production Using Two Plant Protein Sources. *Tropical Journal of Animal Science*. 2(1): 159 – 166.

4. Owen, O.J., Alawa, J.P., Wekhe, S.N., Isirimah, N.O., Chukuigwe, E.C., Aniebo, A.O., Ngodigha, E.M. and Amakiri, A.O. (2008). Incorporating poultry litter in animal feed: a solid waste management strategy. *Egyptian Journal of Animal Production* (In press).
5. Ajala, M.K. and Balogun, J.K. 2004. Economics of rabbit production in Zaria, Kaduna State. *Tropical Journal of Animal Science*, 7(1): 1-10
6. Biobaku, W.O. and Oguntona, E.B. 1997. The effects of feeding multi nutrient mini blocks and pelleted diet on the growth of rabbits. *Nigeria. Journal of Animal Production*, 24(2): 147-149.
7. Omudu, E. A. and Iyough, A. 2005. "Ecological Studies of the Gastropod Fauna of Some Minor Tributaries of River. *Research International*, 2(2): 306-310
8. Cobbinah, J. R., Vink, A. and Onwuka, B. 2008. Snail farm: Production, Processing and Marketing. Agrodok- Series No. 47, Agromisa Foundation, CTA, Wageningen, Pp. 84.
9. Saunders – Blades, J. L., Maclsaac, J. L., Korver, D. R. and Anderson, D. M. 2009. The effect of calcium source and particle size on the production performance and bone quality of laying hens. *Poultry Science*, 88: 3338 – 353.

10. Wikipedia 2015. Benue State. Geology and Environment. Wikipedia the free encyclopedia. "Retrieved From" <https://en.wikipedia.org>.

Formatted: Default Paragraph
Font, Font: (Default) Calibri, 11 pt

11. AOAC. 1995. Official Methods of Analytical chemists 16th edition, Arlington, Virginia, USA.

12. MINITAB Statistical Software 2014. V.16, Minitab Inc. P.A., US.

13. Obi, I. U. 1990. *Statistical Methods of Detecting Differences Between Treatment Means*. 2nd Edition. Snaap Press. Enugu, Nigeria Pp.25-85.

14. Pauzenga, U. 1985. Feeding parent stock. *Zootec. Int.*, 22,22-24.

15. Omole, A. J., Ogbosuka, G. E., Salaka, R. A. and Ajayi, O. O. 2005. Effect of replacing oyster shell with gypsum in broiler finisher diet. *Journal of Applied Sciences Research* 1 (2) : 2 4 5 – 2 4 8 .

16. Okorie, K.C. 2003. The Effect of palmitic acidfortified maize wet milling by-product on the performance of weaner rabbits. *Czech. Journal of Animal Science*, 48(9): 365-370.

17. Ayers, A.C., Barrett, R.P. and Cheeke P.R. 1996. Feeding value of tree leaves (Hybrid Poplar and black locust) evaluated with sheep, goats and rabbits.

Animal Feed Science Technology, 57: 51 - 52

Comment [H49]: check font type and size in table 3 for consistency

Formatted: Expanded by 0.65 pt