

## Original Research Article

# Antioxidant activities of vitamin C and bi-carbonate buffers on hormones secretion and serum metabolites of heat-stressed rabbit buck

### ABSTRACT

**Aims:** To evaluate the antioxidant activities of vitamin C and bicarbonate buffers on hormones secretion and serum metabolites of heat-stressed rabbit buck.

**Methodology:** A total of twenty-five (25) Adult rabbit [buckbucks](#) (New Zealand White crosses) of 12 months old were used in this study. The rabbits were randomly allotted into the experimental treatments of five treatment groups with five (5) rabbits per treatment in a Completely Randomized Design (CRD). The treatment groups consisted of: Control, Sodium Bicarbonate ( $\text{Na}_2\text{HCO}_3$ ) and Potassium bicarbonate ( $\text{KNO}_3$ ), Vitamin C, and baobab fruit pulp meal (BFPM) as supplements respectively. The experiment lasted for 9 weeks. Temperature humidity of the pen house and thermoregulatory condition of the rabbits were evaluated. Blood samples (5 ml) were collected from the ear vein at 10.00 h from four rabbits chosen randomly from each group of rabbits respectively before and the end of the experiment for serum metabolite, thyroxine and testosterone hormones evaluation.

**Results:** It was found that Vitamin C and BFPM significantly ( $P<0.05$ ) reduced thermoregulatory parameters and increased hormonal secretions. The pattern of secretion of serum metabolites was not significantly ( $P<0.05$ ) consistent

**Conclusion:** It was concluded that Vitamin antioxidants were [more](#) effective to ameliorate heat stress than bicarbonate buffers.

**Key Words:** Antioxidants, Heat stress, Hormones, Thermoregulation

### 1. INTRODUCTION

Rabbits are hoped to play an important role in solving meat production deficiency, particularly in Nigeria. Nigeria is localized in the tropical regions; rabbits reared in Nigeria are suffered from a variety of problems related to heat stress. Poor reproductive efficiency is one of the factors militating against the increase in the production output of rabbits. The rabbit reproductive bucks are prone to heat stress because of their poor functional sweat gland. High ambient temperature has been shown to increase the free radicals and other reactive oxygen (ROS) species production in body fluids and tissues [1-2]. The accumulation of these free radicals due to over-production or a decreased antioxidant defense leads to damage of biological macromolecules and disruption of normal cell metabolism [3]. Heat stress is responsible for the increased production of ROS [4], decrease feed intake and weight gain [5] increase rectal temperature and heart rate [5, 6], and also increases the scrotal temperature which disrupts spermatogenesis and ultimately causes infertility [7-9]. Heat stress has been reported to negatively affect the testicular function and also lead to low sperm quality and viability [10] this has led to a decrease in the rate of reproduction of young rabbits.

Several methods including nutritional and physiological means to [alleviatingalleviate](#) the negative effects of high environmental temperature on the performance of rabbits have been documented [11]. Studies [12-14] have shown that antioxidant nutrient supplementation, especially vitamin C, E, and A, zinc, and chromium, can be used to mitigate the negative effects of environmental stress. Vitamin C is also present in some plants; plants and their parts could serve as [phytobioticsprobiotics](#) and antioxidants to the

livestock [15-16]. Some phytochemicals in plants improve antioxidant, anti-microbial, feed flavor, and palatability which could result in increased feed intake and performance in animals [16]. NaHCO<sub>3</sub> was found to improve oxidative stress and heat tolerance by immunomodulation. Sodium bicarbonate in feed or water has shown potential benefits on production performance [17-19], egg characteristics [20-21], and blood profile [22] in poultry birds and rabbits exposed to heat stress. It is cheap, easily available, and easy to handle, therefore, can be safely used to ameliorate the adverse effects caused by heat stress. The influence of antioxidants on hormone secretion and serum metabolites of heat-stressed reproductive bucks was therefore investigated.

## 2. Materials and methods

### 2.1 Experimental site

This study was carried out at the Rabbit Unit of the National Animal Production Research Institute (NAPRI) Shika, Zaria. Shika lies between 11° 12' 42" N and 7° 33' 14" E at an altitude of 691 m above sea level [23]. Zaria has an average [rain-fall](#) of 1100mm which starts from late April and early May to mid-October and an average temperature of 37°C and average relative humidity of 75%

### 2.2 Preparation of buffer solution

Potassium bicarbonate, sodium bicarbonate, and carbonate anhydrous salts were purchased from a laboratory equipment and chemicals vendor in Samaru-Zaria Nigeria. Distilled water was prepared in the Multiuser Laboratory of the Department of Chemistry, Ahmadu Bello University Zaria. The buffer solution was prepared according to the methods of [24] at a pH of 7.5 in the Department of Biochemistry, Ahmadu Bello University Zaria.

### 2.3 Experimental animals, diets, and design

A total of thirty (25) Rabbit Bucks (New Zealand White crosses) of 12 months old were used in this study. The rabbits were randomly allotted into the experimental treatments of five treatment groups with five (5) rabbits per treatment in a Completely Randomized Design (CRD). The treatment groups consisted of Control (Basal diet with no supplementation), Sodium Bicarbonate (NaHCO<sub>3</sub>) and Potassium bicarbonate (KNO<sub>3</sub>) solution with basal diet respectively, Basal diet- Vitamin C, and BFPM as supplement respectively (designated T1, T2, T3, T4, and T5 respectively). The basal diet composition (kg) was as follows: Maize 30, Groundnut haulm 25, Groundnut cake 10, soybean meal 10, rice bran 15, Bone meal 9.2, common salt 0.35, mineral-vitamin premix 0.25, dl-methionine 0.1, and lysine 0.1. The basal diet was formulated to meet the nutrient requirements of growing rabbits according to the recommendations of [25]. The calculated analysis of the nutrient composition of the basal diet is shown in Table 1. The water was offered ad libitum but changed daily in the morning. All recommended managerial practices were dully observed and the study lasted for 9 weeks.

**Table 1: Calculated composition of the experimental diets**

Nutrients	Composition
Metabolizable Energy (kcal/kg)	2300
Crude Protein (%)	17
Crude fiber (%)	14

### 2.4 Housing

The animals were housed in perforated metallic hutches measuring 75 X 75 X 75 cm and raised 80 cm from the floor level in a naturally ventilated building. The hutches were thoroughly washed and disinfected with a locally made disinfectant and allowed to dry for one week before the animals were brought. Feed and watering troughs which were made of burnt clay were provided in each hutch. The rabbits were placed individually in clearly labeled cells.

### 2.5 Meteorological data of rabbit microclimate

The microclimate (ambient temperature and relative humidity values) within the rabbit house were recorded twice daily at 08:00 h and 15.00 h during the study period using a digital thermometer (Cocet, Shenzhen-Guangdong, China). The data collected was used to compute the temperature-humidity index

(THI), an indicator of the thermal comfort level of the rabbits. The THI was calculated using the modified formula for the rabbit by Marai et al. [5] as follows:  $THI = t - [(0.31 - 0.31 \times RH) (t - 14.4)]$

Where RH = relative humidity /100.  
t = ambient temperature.

The values of THI obtained were compared to that classified for tropical regions as shown below:  
1) < 27.8 = Absence of heat stress, 2). 27.8 - 28.9 = Moderate heat stress, 3)28.9 – 30 = Severe heat stress and 4) above 30 = Very severe heat stress.

## 2.6 Measurement of Thermoregulatory Parameters

Parameters measured included rectal temperature (RT), ear temperature (ET), respiratory rate (RR), and heart rate (HR). Measurements were taken at 14.00 h to 15.00 h of the day. Rectal and ear temperature were measured with a digital thermometer. The ear temperature was measured by placing the digital thermometer in direct contact with the central area of the auricle. RR was measured by visually counting the flank movements for one minute with the help of a stopped clock. HR was measured by counting the heartbeat for one minute with the help of a stethoscope.

## 2.7 Serum evaluation

Blood sampling was done at the beginning and at the end of the experiment; the sampling was done at 10.00 h. Four rabbits were randomly selected from each treatment group and 5 ml of blood was collected from their ear veins into sample bottles without anticoagulants. The blood sample was allowed to clot and the serum was harvested after centrifuging the samples at 3000 rounds/minute for 15 minutes. The serum harvested was stored at -10°C until when analyzed. Serum thyroxine and testosterone concentrations were determined by using commercially-available ELISA Kits (Diagnostic Procedure Corp., Los Angeles, CA; The USA) according to the manufacturer's instructions. Serum glucose, total protein, albumin, and cholesterol concentration were evaluated using an auto-analyzer and Chemical Commercial Kits from Stanbio Laboratory Inc. San Antonio, Texas, USA, according to the manufacturer's instruction. The detection ranges of the  $T_4$  were 26-58.4 g/mg.

## 2.8 Statistical analysis

Data obtained from the study were subjected to analysis of variance using the general linear model procedure of SAS [26]. Significant differences among treatment means were separated using the pairwise difference (Pdfff) in the SAS package.

# 3. RESULTS AND DISCUSSION

## 3.1 Monthly temperature-humidity index (THI)

The monthly temperature-humidity index (THI) inside the rabbitry during the experimental period is shown in Figure 1. THI in the mornings averaged 26.44°C while the Afternoon THI averaged 28.74°C. The graph also shows that the THI values kept increasing from the month of February with a peak in May. There was a decline in THI in the month of June. The THI value of 27°C (Feb) indicated that there was no heat stress in the month of February, while the THI values of 28°C (March), 29.5°C (April), 31.2°C (May), and 28°C (June) are indications that the rabbit house was moderately thermally stressful, severely thermally stressful and very severely thermally stressful [5] in these months. The averaged THI 28.74°C during the experimental period indicated that the rabbit house was thermally stressful and may have had adverse effects on the rabbits [5]. Overall data obtained indicated that THI in the afternoon was higher by 1.24 % than THI in the morning.

## 3.2 Thermoregulatory Response of Adult Rabbit Bucks

The result of the thermoregulatory response (Table 2) shows that Vitamin C and BFPM significantly ( $P < 0.05$ ) reduced heart rate, rectal, and ear temperature in adult bucks compared to  $KHCO_3$  and  $Na_2CO_3$  or the control. Antioxidant vitamins were reported to be effective to alleviate heat load in rabbits [27, 13-14]. The administration of ascorbic acid [28], glutathione supplementation [12], during exposure to high environmental temperatures reduces the body temperature in chickens. In heat-stressed sheep, selenium injection decreased rectal temperature and body weight loss [29]. The BFPM treatments gave the best performance, which agrees with the fact that the presence of phytochemical compounds in plants

may facilitate the ability of animals to maintain their body homeostasis including body temperature by provoking endogenous cellular defense mechanisms to cope with oxidative stress and inflammation induced by heat stress [30].  $\text{NaHCO}_3$  could increase respiratory alkalosis severity [31], which may be responsible for the high values of heart rate, rectal and ear temperature recorded in rabbits treated with the buffers.

### 3.3 Serum Metabolites of Adult Rabbit Bucks

Serum metabolites of rabbits administered the antioxidants (Table 3) showed that triglyceride and phosphorous were significantly ( $P < 0.05$ ) different among the treatment groups. The final serum values in BFPM-treated rabbits significantly ( $P < 0.05$ ) improved albumin; the control recorded significantly higher serum glucose while  $\text{Na}_2\text{CO}_3$  significantly increased serum calcium level. The significant increase in final serum albumin in male rabbits administered vitamin C and BFPM, may be due to the fact that vitamins are essential parts of some of the enzymes or co-enzymes [32] and enzymes are important in food digestion, making micronutrients to be available for growth and other body functions. The significant increase in final serum calcium in the treatment with  $\text{NaHCO}_3$  may be due to the increase in body metabolism and excitation caused by the buffers, thereby increasing blood calcium [32]. The trend in serum glucose recorded in this experiment agrees with the reports of Ondruska et al. [33]. It was attributed to being due to an increase in glucose utilization during muscular movements required for high respiratory activity [34-35], or due to increases in corticosteroid concentrations [36].


### 3.4 Thyroxine Levels in Adult Rabbit Bucks

The effect of buffer, vitamin C and BFPM on thyroxine levels of adult male rabbits (Fig. 2) revealed that initial thyroxine levels of the rabbits did not differ significantly when compared to those of the treatment groups. The control group and  $\text{Na}_2\text{CO}_3$  group recorded the lowest thyroxine levels compared to the rest of the treatments. On termination of the experiment, the values of thyroxine obtained from the serum showed that vitamin C and BFPM increased significantly ( $P < 0.05$ ) thyroxine levels compared to other treatments, and the control group recorded the least value. The lower thyroxine levels recorded in the control ~~reveals~~ reveal that the control group was ~~the~~ heat-stressed and metabolic activities may have been lowered. Previous studies reported a decrease in thyroid hormones with increasing temperature [37-38], a decrease in T3 level, and a concurrent increase in T4 level [39] during heat stress. Yahav et al. [40] reported reduced concentrations of thyroid hormones (triiodothyronine and thyroxine) in the blood plasma and a decreased metabolic rate of pullets and cocks reared under a high ambient temperature. The reduction in thyroxine levels after the experiment compared to the initial values can be attributed to two factors; acclimation of the animals for the control and amelioration of heat stress by the buffers and the vitamin. The values are similar to the report of [41]. Acclimation increases the adaptive ability of birds to subsequent thermal stress by reducing the level of triiodothyronine. The buffers especially  $\text{KCHO}_3$  may have triggered the stimulation of the thyroid glands leading to the secretion of more thyroxine compared to the control while the exogenous antioxidant vitamin sources (vitamin C and BFPM) may have reduced oxidative stress and improved body metabolism, leading to the increase in serum thyroxine [42]. Vitamin C synthesized in rabbit liver has been demonstrated to protect the animal from heat stress and improve disease resistance in rabbits by optimizing the function of the immune system [43], but during stress vitamin C produced is rapidly consumed and the amount synthesized falls below animal requirements. Supplemental vitamin C may therefore reinstate normal metabolic functions during heat stress. Serum concentrations of T4 increased by increasing dietary VC or VE levels of heat-stressed Japanese quails [44] and rabbits [42]. Thyroid hormones are the key hormones in the regulation of metabolism and adaptation of animals to stress [45].


### 3.5 Testosterone Levels in Adult Rabbit Bucks

The initial testosterone concentration of rabbits treated with buffer, vitamin C and BFPM in Fig. 3, did not show any significant difference among the treatments. It was observed that vitamin C and BFPM increased significantly ( $P < 0.05$ ) testosterone concentration compared to the rest of the treatments whose values were lower than that recorded initially. The treatment with  $\text{KCHO}_3$  buffer recorded significantly ( $P < 0.05$ ) lower testosterone levels than the rest of the treatments. The final testosterone values of the control that was lower than the initial values may be attributed to heat stress, which might have affected the Leydig cells. Jelodar and Zare [46] reported a decrease in serum testosterone

concentration of rats exposed to radiation from phones; and attributed the decrease to the effect of radiation on Leydig cells, pituitary, or hypothalamus and alteration of gonadotropin secretion. It is also possible that the buffers, especially  $\text{KHO}_3$ , also had a negative effect on the Leydig cells, responsible for the production of testosterone. Stress can also cause structural and pathological changes in the Leydig cells. Apoptosis associated with nuclear damage of the cells can lead to a decrease in testosterone and estrogen production [47]. It is also possible that the buffers interfered with the transfer of free cholesterol to mitochondria of Leydig cells, which is an important step in steroidogenesis, and also disrupted the conversion of cholesterol to testosterone by impairing the activity of key regulatory enzymes in steroidogenesis [48]. The significant improvement recorded in rabbits treated with vitamin C and BFPM could be the vitamins counteracted free radicals that may limit the normal functioning of the Leydig cells from functioning properly. Vitamin C has been reported to be an effective antioxidant.



Formatted: Font: (Default) Arial, 12 pt, Bold



Formatted: Font: (Default) Arial, 12 pt, Bold

Figure 1: Monthly Temperature Humidity Index of the Pen House

Table 2 Effect of Bicarbonate Buffers, Vit C and BFPM on Thermoregulatory Response of Adult Rabbit Bucks

Treatments

Parameters	Control	KHCO <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	Vit. C	BFPM	SEM
Respiratory Rate (counts/min)	158.34	154.02	148.91	150.58	141.03	3.78
Heart Rate (beat/min)	141.03 <sup>b</sup>	143.23 <sup>ab</sup>	143.53 <sup>a</sup>	140.99 <sup>b</sup>	141.03 <sup>ab</sup>	0.67
Rectal Temperature (°C(°C))	38.73 <sup>a</sup>	38.04 <sup>a</sup>	38.29 <sup>a</sup>	37.97 <sup>ab</sup>	37.04 <sup>b</sup>	0.28
Ear Temperature (°C(°C))	36.79 <sup>a</sup>	35.59 <sup>ab</sup>	36.13 <sup>ab</sup>	35.23 <sup>ab</sup>	34.99 <sup>b</sup>	0.47

Means within rows with different superscripts are significantly different: P<=0.05


Vit C = Vitamin C, BFPM = Baobab fruit pulp meal

**Table 3 Effects of Bicarbonate Buffers, Vit C and BFPM on Serum Metabolite of Adult Rabbit Bucks**


Parameters	Treatments					SEM
	Control	KHCO <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	Vit. C	BFPM	
<b>Initial</b>						
Glucose (mg/dl)	4.17	5.03	4.12	4.60	4.70	0.33
Total Protein (mg/dl)	65.00	67.00	63.37	66.80	67.23	1.67
Albumin (mg/dl)	35.00	36.67	39.00	37.33	40.00	1.53
Cholesterol (mg/dl)	1.37	1.10	1.23	1.20	1.30	0.56
Triglyceride (mg/dl)	0.70 <sup>b</sup>	1.07 <sup>ab</sup>	0.98 <sup>ab</sup>	0.87 <sup>ab</sup>	1.17 <sup>a</sup>	0.09
Calcium (mg/dl)	2.38	2.32	2.37	2.38	2.32	0.06
Phosphorous (mg/dl)	0.87 <sup>b</sup>	1.15 <sup>a</sup>	1.07 <sup>ab</sup>	1.00 <sup>ab</sup>	1.00 <sup>ab</sup>	0.04
<b>Final</b>						
Glucose (mg/dl)	4.70 <sup>a</sup>	4.13 <sup>b</sup>	4.20 <sup>b</sup>	4.47 <sup>b</sup>	4.50 <sup>b</sup>	3.57
Total Protein (mg/dl)	64.00	64.00	67.33	69.33	70.67	1.58
Albumin (mg/dl)	33.67 <sup>b</sup>	32.67 <sup>c</sup>	33.33 <sup>b</sup>	38.66 <sup>ab</sup>	40.00 <sup>a</sup>	1.33
Cholesterol (mg/dl)	1.50	1.37	1.38	1.55	1.53	0.07
Triglyceride (mg/dl)	2.90	1.33	1.27	1.27	1.37	0.71
Calcium (mg/dl)	2.30 <sup>b</sup>	2.29 <sup>bc</sup>	2.44 <sup>a</sup>	2.25 <sup>c</sup>	2.42 <sup>ab</sup>	0.03
Phosphorous (mg/dl)	1.00	1.07	0.90	1.00	1.06	0.04

Means within rows with different superscripts are significantly different: P<=0.05

Vit C = Vitamin C, BFPM = Baobab fruit pulp meal




Formatted: Font: (Default) Arial, 12 pt, Bold




Formatted: Font: (Default) Arial, 12 pt, Bold

**Fig. 2 Effects of Bicarbonate Buffers, Vit C and BFPM on Thyroxine Levels in Adult Rabbit Bucks**  
 Vit C = Vitamin C, BFPM = Baobab fruit pulp meal

UNDEI



Formatted: Font: (Default) Arial, 12 pt, Bold



Formatted: Font: (Default) Arial, 12 pt, Bold

Vit C = Vitamin C, BFPM = Baobab fruit pulp meal

**Fig. 3 Effects of Bicarbonate Buffers, Vit C and BFPM on Testosterone Levels in Adult Rabbit Bucks**

#### 4. CONCLUSION

The vitamin antioxidants displayed superiority in ameliorating heat stress on most of the parameters that were evaluated compared to the bicarbonate buffer solution. Baobab fruit pulp meal treatments gave the best performance. The use of a plant source of vitamin C in rabbit diets is therefore encouraged and recommended to be during hot environmental conditionconditions.

#### Ethical approval

All authors hereby declare that the procedures for care and handling of animals in this study were strictly followed in accordance with the code of ethics for animal experimentexperiments as stated in [http://ec.europa.eu/environment/chemicals/lab\\_animal/legislation\\_en.htm](http://ec.europa.eu/environment/chemicals/lab_animal/legislation_en.htm).

#### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly useused products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use

these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by the personal efforts of the authors.

## REFERENCE

1. El-Ghaffar SKA, Aly M, Fatma Moustafa A, Mahmoud AZ. Pathological studies on the rabbit viral hemorrhagic disease (RVHD) with special reference to the use of vitamin A, E and C as prophylaxis. *Assiut-Vet. Med. Journal*. 2000; 43: 85, 251-274.
2. Akiyama M. In vivo scavenging effect of ethylcysteine on reactive oxygen species in human semen. *Nippon Hinyokika Gakkai Zasshi*. 1999; 90(3):421-428.
3. Spurlock ME, Savage JE. Effects of dietary protein and selected antioxidants on fatty haemorrhagic syndrome induced in Japanese quails. *Poult. Sci*. 1993; 72: 2095-2105.
4. Greco E, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Tesarik J. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *Journal of Andrology*. 2005; 26: 349-353.
5. Marai IFM, Ayyat MS, Abd El-Monem UM. Growth performance and reproductive traits at first parity of New Zealand white female rabbits as affected by heat stress and its alleviation under Egyptian conditions. *Tropical Animal Health Production*. 2001; 33: 451-462.
6. Marai IFM, Habeeb AAM, Gad AE. Rabbits' productive, reproductive and physiological traits as affected by heat stress (A Review). *Livestock Production Science*. 2002; 78: 71-90.
7. Paul, C, Murray, AA, Spears, N, Saunders, PTK. A single, mild, transient scrotal heat stress causes DNA damage, subfertility and impairs formation of blastocysts in mice. *Reproduction* 2008; 136, 73-84
8. Jung, A, Leonhardt, F, Schill, WB, Schuppe, HC. Influence of the type of undertrousers and physical activity on scrotal temperature. *Hum. Reprod*. 2005; 20, 1022-1027.
9. Lue, Y, Hikim, APS, Wang, C, Im, M, Leung, A, Swerdlof, RS. Testicular heat exposure enhances the suppression of spermatogenesis by testosterone in rats: The "two-hit" approach to male contraceptive development. *Endocrinology*. 2000; 141, 1414-1424
10. Rasooli, A., Taha Jalali, M., Nouri, M., Mohammadian, B., Barati, F. Effects of chronic heat stress on testicular structures, serum testosterone and cortisol concentrations in developing lambs. *Anim. Reprod. Sci*. 2010; 117, 55-59.
11. Sharaf AK, El-Darawany AA, Nasr AS, Habeeb AAM. Recent techniques for amelioration the effect of heat stress conditions on male rabbits. *Zagazig Journal of Agriculture Research*, 2019; 46 (2): 501-514
12. Sahin K, Onderai M, Sahin N, Gursu, MF, Kucuk O. Dietary vitamin C and folic acid supplementation ameliorate the detrimental effects of heat stress in Japanese quail. *Journal of Nutrition* 2003; 133, 1852- 1886.

13. In-Surk J, Ko Y, Moon Y, Sohn S. Effects of vitamin C or E on the proinflammatory cytokines, heat shock protein 70 and antioxidant status in broiler chicks under summer conditions. *Asian-Australian Journal of Animal Science* 2014; 27, 749–756.
14. Prabsattroo T, Wattanathorn J, Iamsaard S, Muchimapura S, Thukhammee W. Moringa oleifera leaves extract attenuates male sexual dysfunction. *American Journal of Neuroscience*. 2012; 3: 17–24
15. Dhama, K, Latheef SK, Manin S, Samad HA, Karthik K, Tiwari R, Khan RU, Alagawany M, Farag MR, Alam GM, Laudadio V, Vincenzo T. Multiple beneficial application and modes of action of herbs in poultry health and production. A review. *Int J Pharmacol* 2015; 11(3):152–176
16. Valenzuela-Grijalva NV, Pinelli-Saavedra A, Muhlia-Almazan A, Dominguez-Diaz D, Gonzalez-Rios H. Dietary inclusion effects of phytochemicals as growth promoters in animal production. *Journal of Animal Science Technology*. 2017; 59:8.
17. Ahmad T, Sarwar M, Nisa M, Haq A, Hasan Z. Influence of varying sources of dietary electrolytes on the performance of broilers reared in a high temperature environment. *Animal Feed Science and Technology*. 2005; 120: 277-298.
18. Khattak FM, Acamovic T, Sparks N, Pasha TN, Hussain MH, Joiya M. Comparative efficacy of different supplements used to reduce heat stress in broilers. *Pakistan Journal of Zoology*; 2012; 44: 31-41.
19. Peng Y, Wang Y, Ning D, Guo Y. Estimation of dietary sodium bicarbonate dose limit in broiler under high ambient temperatures. *The Journal of Poultry Science*. 2013; 50(4):346-353.
20. Kaya I, Karademir B, Ukar O. The effects of diet supplemented with sodium bicarbonate upon blood pH, blood gases and eggshell quality in laying geese. *Veterinary Medicine Czech*. 2004; 49: 201-206.
21. Jiang MJ, Zhao JP, Jiao HC, Wang XJ, Zhang Q, Lin H. Dietary supplementation with sodium bicarbonate improves calcium absorption and eggshell quality of laying hens during peak production. *British Poultry Science* 2015; 56(6):740-747.
22. Kurtoglu V, Kurtoglu F, Balevi T. Effects of sodium bicarbonate, potassium chloride and sodium chloride supplementation on some blood biochemical parameters in laying hens. *Proceedings of the European Symposium on Poultry Nutrition*. 2007; Aug 26-30; Strasbourg, France. Pp.189-192
23. Ovimaps. Ovi location map; Ovi earth imagery 2014; March 5th,
24. Chandra M. Buffer: A guide for the preparation and use of buffers in biological systems. EMD Biosciences, Inc., an affiliate of Merck KGaA, Darmstadt, Germany, 2006.
25. NRC, Nutrient Requirement for Rabbits. 9th Ed, (National Academy Press, Washington D. C., U.S.A. 1994.
26. S.A.S. Statistical Analysis System Institute, User's Guide. 6<sup>th</sup> Edition. North Carolina, USA, 2002.

27. Anoh KU. Alleviating heat stress with Baobab fruit pulp meal: Effect on testosterone concentration and gonadal and epididymal morphometry of adult rabbit bucks. *Acad. J. Agric. Res.* 2017; 5(8): 178-182.
28. Mckeeand JS, Harrison PC. Supplemental ascorbic acid does not affect heat loss in broilers exposed to elevated temperature. *Journal of Thermal Biology.* 2013; 38, 159– 162.
29. Alhidary IA, Shini S, Al Jassim R AM, Gaughan JB. Effect of various doses of injected selenium on performance and physiological responses of sheep to heat load. *Journal of Animal Science.* 2012; 90, 2988– 2994.
30. Akbarian A, Michiels, J, Degroote J. Association between heat stress and oxidative stress in poultry; mitochondrial dysfunction and dietary interventions with phytochemicals. *J Animal Sci Biotechnol* 2016; 7: 37.
31. Teeter, RG, Smith, MO, Owens, FN, Arp, SC, Sangiah, S, Breezile, JE. Chronic heat stress and respiratory alkalosis: Occurrence and treatment in broiler chicks. *Poultry Science.* 1985; 64: 1060-1064.
32. Burger A G. Environment and thyroid function. *Journal of Clinical Endocrinology and Metabolism.* 2004; 89:1526.
33. Ondruska L, Rafay J, Okab, AB Ayoub, MA, Al-Haidary, AA, Samara, EM, Parkanyi, V, Chrastinova, L, Jurcik, R, Massanyi, P, Lukac, N, Supuka, P. Influence of elevated ambient temperature upon some physiological measurements of New Zealand White rabbits. *Veterinari Medicina.* 2011; 56 (4): 180–186.
34. Habeeb AA, Marai IFM, Kamal TH. Heat stress. In *Farm Animals and the Environment*, edited by C. Philips and D. Piggins. C.A.B. International, U.K. 1992; pp. 27-47.
35. Okab, AB, El-Banna, SG Koriem, AA. Influence of environmental temperatures on some physiological and biochemical parameters of male New-Zealand rabbits. *Slovak Journal of Animal Science.* 2008; 41: 12–19.
36. Habeeb AAM, Marai IFM, El-Maghawry AM, Gad AE. Growing rabbits as affected by salinity in drinking water under winter and hot summer conditions of Egypt, *Egyptian Journal of Rabbit Science.* 1997; 7: 81-94.
37. Williams J, Njoya J. Plasma concentrations of thyroid hormones in growing pullets and laying hens reared under two tropical climates and fed on diets with different energy concentrations. *British Poultry Science.* 1998; 39: 579-582.
38. Sokolowicz Z, Herbut E. Effect of chronic high temperature stress on thyroid activity and metabolic rate of broiler pullets cockerels. *Annals of Animal Science.* 1999; 26, 377-383.
39. Marai IFM, Habeeb AAM, Gad AE. Growth performance traits and the physiological background of young doe rabbits as affected by climatic conditions and lighting regime, under sub-tropical conditions of Egypt. *Proceedings - 8th World Rabbit Congress –September 7-10, 2004 – Puebla, Mexico* pp. 288 – 297.

40. Yahav, S., A. Straschnow, I. Plavnik, Hurwitz SV. Effects of diurnally cycling versus constant temperatures on chicken growth and food intake. *British Poultry Science*. 1996; 37: 43-54.
41. Garriga et al, In: Usama T Mahmoud Mootaz AM Abdel-Rahman and Madeha HA Darwish Effects of Propolis, Ascorbic Acid and Vitamin E on Thyroid and Corticosterone Hormones in Heat Stressed Broilers. *Journal of Advanced Veterinary Research*. 2014; 4(1) 18-27.
42. Daader AH, Al-Sagheer, AA, Gabr HA, Abd El-Moniem EA. Alleviation of heat-stress-related physiological perturbations in growing rabbits using natural antioxidants. *Spanish Journal of Agricultural Research*. 2018; 16: (3):e0610. <https://doi.org/10.5424/sjar/163-13184>
43. Bain, BS. The role of Vitamin C in stress management. *World's Poultry Science*. 1996; 12 (4): 34-41.
44. Sahin K, Kucuk O, Sahin N, Sari M. Effects of vitamin C and vitamin E on lipid peroxidation status, serum hormone, metabolite, and mineral concentrations of Japanese quails reared under heat stress (34° C). *Int J Vitam Nutr Res*. 2002; 72: 91-100. Google Scholar
45. Brecchia G, Zerani M, Bonano A, Boiti C. Metabolic adaptations in neonatal mother-deprived rabbits. *World Rabbit Sci*. 2010 18: 229-236.
46. Jelodar GA, Zare Y. Effect of radiation leakage of microwave oven on rat serum testosterone at pre and post pubertal stage, *Journal of Human reproduction*. 2008; 15(4): 64-68.
47. Hamada AJ, Singh A, Agarwal A. Cell phones and their impact on male fertility: Fact or fiction. *Open Reproductive Science Journal*. 2011; 5: 125-137.
48. Ebomoyi, MI, Ahumibe KC. Serum testosterone and morphology of the testes in wistar rats following chronic garlic feeding. *Physiology Pathophysiology*, 2010; 1(3): 39-43.