

# Influence of weather factors on Succession and incidence of insect pests in soybean

## ABSTRACT

**Background** The present investigation was conducted at Rajmata Vijayaraje Scindia Krishi Vishwa Vidhyalaya Zonal Agricultural Research Station, Morena (M.P.) to studies the succession and influence of weather factors on incidence of insect pests and their natural enemies of soybean. On the basis of pest populations and their succession level, nine species of the insect pests was observed during the crop season. The activity of aphid, whitefly, jassid, tobacco caterpillar, green semilooper, stem fly, girdle beetle, green stink bug and lady bird beetle was observed from the last week of July to second week of **October**. The peak population of aphid (2.20/ aphids/ 3 leaves), whitefly (1.98/3 leaves), jassid (3.25 jassid/3 leaves) and tobacco caterpillar (8.15 larvae/mrl), green semilooper (3.00 larval/mrl), stem fly (5.20 percent), girdle beetle (4.80 percent/ plant), green stink bug (5.35/ mrl) and lady bird beetle (1.40/plant) was observed in between 31<sup>st</sup> to 35<sup>th</sup> **SMW**. With regards to correlation with weather factors, aphid, whitefly, jassid, tobacco caterpillar, green semilooper, girdle beetle, green stink bug and lady bird beetle population was found positively correlated, with maximum temperature, relative humidity and rainfall whereas stem fly found negatively correlated with maximum **temperature**.

**Key words:** Incidence, insect pests, **soybean**, succession, weather factors

## **COMPETING INTEREST**

## INTRODUCTION

**Soybean** is one of the most important oil seed crops in the country grown for oil and protein production in both the **rabi** and **kharif** seasons. It is a rich source of animal feed protein and cooking oil (Schmutz *et al.* 2010). It is known as the “GOLDEN BEAN” of the 20th Century. Though, Soybean is a legume crop, yet it is widely used as oilseed. The area under soybean cultivation has increased from 10.60 million ha to 11.25 million ha 2017-18 to 2019-20 and production from 10.98 MT to

**Comment [WU1]:** Describe in brief the problem of the study at least in a single statement as background information.

**Comment [WU2]:** Include data collected and which statistical data analysis method used here.

**Comment [WU3]:** Abstracts should be free from abbreviations.

**Comment [WU4]:** At last write the main conclusion of the study

**Comment [WU5]:** Write competing interest like The article has no any conflict interest and has not been published in another publication and is not being submitted simultaneously to another journal.

**Comment [WU6]:** In the first paragraph introduce the morphology of the soybean to be easily known by any reader.

**Comment [WU7]:** These local names should be written in bracket after the English name to be understand by International readers. Is it summer, winter, autumn or spring?

11.32 MT. In India, soybean has acquired third position among the oil consumption after groundnut and mustard. Madhya Pradesh contributes about 67% area and 58% production in the country and is called as “Soya state” (Anon. 2005 and Anon. 2007). In Madhya Pradesh, soybean occupies an area of 5.24 m ha with production of 6.72 MT and productivity of 1286 kg/ha (Anon. 2019). Soybean is infested by more than 275 insect pests on different plant parts of soybean throughout its growth stage and about a dozen of them have been reported causing serious damage from sowing to harvesting (Ramesh Babu, 2010). The grub of girdle beetle, [*Obereopsis brevis* (Swed)] bores the main stem and branches resulting in stunting plant growth and sometime whole plant succumb to injury. The stem fly, (*Melanagromyza sojiae*) attacks the soybean throughout the growing season, but the most vulnerable period is within 3-4 weeks after germination. The maggot may reduce the grain yield up to 33 per cent (Singh and Singh, 1992). This crop suffers a lot due to the attack of number of insect pests (Lal *et al.* 1981). It is mainly attacked by gram podborer, *Helicoverpa armigera* Hubner; leaf eating caterpillar, *Spodoptera litura* Fabricius; green semilooper, *Chrysodeixis acuta* Walker; grey semilooper, *Amynaocto* Guenee; leaf miner, *Aproeremamodicella* Deventer; whitefly, *Bemisia tabaci* Gennadius; stem fly, *Ophiomyia phaseoli* Tryon; thrip, *Caliothrips indicus*; aphid, *Aphis glycine* Koch and jassid, *Empoasca kerri* Pruthi (Ahirwari *et al.* (2015). Population density of insect pests and their natural enemies fluctuates with changing weather conditions. Abiotic factors regulate seasonal incidence, population count and development rates of the pests and natural enemies. As the cultivation of soybean has expanded around the world, crops become susceptible to different environmental and biotic stress which has increased the pest infestations. Among these temperature and relative humidity play key role that regulates population dynamics, development rates and seasonal incidence of pest and their natural enemies.

**Comment [WU8]:** Reference. Specially such types of data needs updated citation.

**Comment [WU9]:** Updated it

**Comment [WU10]:** Reference

## MATERIALS AND METHODS

The present studies were conducted during the kharif season 2018 at Rajmata Vijayaraje Scindia Krishi Vishwa Vidhyalaya Zonal Agricultural Research Station,

**Comment [WU11]:** Write its name in English and kharif in bracket and italic. Local names should be italic.

Morena (M.P.). The experimental area is having uniform topography, gentle slope and adequate drainage. The soybean variety RVS2007-6 was used in the experiment with plot size 8.0 m X 3.0m, no of row/plot 20, row length 3.0m and row to row spacing 40 cm.

**Observations recorded:**

**Stem fly:**

Observation on stem fly was recorded on the randomly selected five plants at weekly interval from germination till harvest of the crop. To record the tunneling caused by the maggot of stem fly the plants were uprooted and open vertically. Plant height and tunnel length were also measured for calculating percent tunneling.

**Girdle beetle:**

To record the seasonal incidence of girdle beetle, observations on randomly selected five plants were made at weekly interval starting from the initiation of infestation till harvest of the crop. The pest incidence was recorded by counting healthy and damaged plant by girdle beetle and the data ~~would be were~~ calculated in percentage.

**Foliage feeders and their natural enemies:**

For recording the pest succession of foliage feeders and their natural enemies, the observations were recorded once in a standard week at 5 sites of one meter row length. The insect pests appearing on the crop right from seedling up to harvest were recorded. The crop was kept unprotected for this purpose. The sequence in which the pests appeared was also noted.

**Sucking pests:**

Observation on sucking pests were recorded on 3 leaves/plant (top, middle and bottom) in 10 randomly selected plants.

Observations on the incidence of major insect pests were recorded from the first appearance of the pest and it was continued till maturity of the crop at weekly intervals and data was correlated with meteorological parameters.

## **RESULTS AND DISCUSSION**

The results of present studies influence of weather factors on succession and incidence of insect pests in soybean are given in table 1, 2, 3 and 4.

**Comment [WU12]:** Table number should be written at the end of each result. Not like this.

Succession of insects and field incidence revealed that eight species of insect pests and one species of natural enemies were observed which were associated with various stages of the soybean crop during kharif 2018-19. The major groups (sucking pest and foliage feeder) of insect pests attacked at the various stages of the crop were aphid, whitefly, jassid, tobacco caterpillar, green semilooper, stem fly, girdle beetle and green stink bug. Only one natural enemy's lady bird beetle was observed on the soybean crop (insetTable number).

**Comment [WU13]:** It will be better to write according to their damaged or infestation level from highest to lowest

#### **Aphid, *Aphis gossypii* (Hemiptera: Aphididae)**

Aphid was first appeared during last week of July *i.e.* from 30<sup>th</sup> standard week in soybean during vegetative stage and the population of aphid increased and reached the peak level (2.20 aphid/ three leaves) on first week of August *i.e.* 31<sup>th</sup> standard week, subsequently, it declined gradually and disappeared during fourth week of September *i.e.* 38<sup>th</sup> standard week. Present findings are in accordance with Suyalet *al.* (2018) stated that.....

Among the different weather parameters studied, maximum temperature showed highly significant positive correlation with aphid population ( $r = 0.627$ ) and relative humidity showed non-significant but positive correlation with aphid population ( $r = 0.369$ ), while, rainfall showed non-significant positive correlation with aphid population ( $r = 0.323$ ). Whereas, minimum temperature reflected non-significant positive correlation with the aphid population (0.160). During the peak aphid population, the maximum and minimum temperature were 30.71<sup>0</sup>C and 24.14<sup>0</sup>C respectively, whereas relative humidity and rainfall were 88.68% and - 03.79 mm respectively (Table no). The result was somehow similar with the result obtained by Patel (2012).

**Comment [WU14]:** It is better to write petal (2012) result

#### **White fly, *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae)**

**Formatted:** Font: Italic

First incidence of white fly was observed in vegetative stage when the crop age was about 25 days old. Results showed that the pest was present-observed on the crop during the entire cropping season and remained active from fourth week of July *i.e.* from vegetative stage of the crop to second week of October (till maturity stage).; population of whitefly reached the peak (1.98/three leaves) at 34<sup>th</sup> MSW. Incidence of

**Comment [WU15]:** Write in full for the first time

white fly was observed from first week of ~~august~~ August to second week of October and the whitefly reached the peak level during last week of August, Similar findings was also reported by Chaudhary *et al.* (2018) and Garg *et al.* (2014).

The results indicated that the maximum temperature ( $r = 0.690$ ) and relative humidity ( $r = 0.543$ ) showed highly significant positive correlation with whitefly population while, minimum temperature and rainfall showed non-significant positive correlation with the whitefly population. During the peak whitefly population, the maximum and minimum temperature were  $30.57^{\circ}\text{C}$  and  $25.79^{\circ}\text{C}$  respectively, whereas relative humidity and rainfall were 95.57% and 07.79 mm respectively and it confirms with the finding of Patel (2012), and the result was also in accordance with Kujur (2011). From the result it was clear that maximum temperature and relative humidity showed highly significant positive correlation with whitefly population. ~~the~~ [The](#) result confirms the findings of Chaudhary *et al.* (2018).

#### **Jassid, *Empoasca kerri* Pruthi (Hemiptera; Cicadellidae)**

First appearance of Jassid was observed in the vegetative stage when the crop age was about 32 days old. From the data it ~~is~~ [was](#) evident that the pest was present in the crop from first week August i.e. from vegetative stage to third week September till reproductive stage. The population of Jassid increased and reached the peak level (3.25 Jassid/ three leaves) on third week of August i.e. 33<sup>rd</sup> standard week, subsequently, it declined gradually and disappeared during second week of October i.e. 40<sup>th</sup> standard week. The incidence of jassid was observed from first week of august to second week of October, and the peak population of jassid was obtained in third week of august. The result was [somewhat](#) similar to Chaudhary *et al.* (2018).

The results of jassid population presented in Table [4.4 and 4.6](#) indicated that the maximum temperature showed non-significant [positive](#) correlation with jassid population ( $r = 0.458$ ) while all other weather parameter also showed non-significant positive correlation with jassid population. During the peak jassid population maximum and minimum temperature were  $32.64^{\circ}\text{C}$  and  $25.57^{\circ}\text{C}$  respectively, whereas relative humidity and rainfall were 88.43% and 03.57 mm respectively. [Sutaria et al.](#) (2010a) ~~also~~ reported the positive correlation of leaf hopper population with minimum temperature and relative humidity.

#### **Tobacco caterpillar, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae).**

**Comment [WU16]:** Compare with other works in detail

**Comment [WU17]:** Tables 4.4 and 4.6 are not indicated in the article. Check it

Tobacco caterpillar first appearance was observed in the vegetative stage when the crop age was about 39 days old. The data showed that the pest was present in the crop from second week August *i.e.* from vegetative stage to second week September till reproductive stage. The population of tobacco caterpillar increased and recorded peak (8.15 larvae/mrl) at 35<sup>th</sup> SMW (First week of September). First incidence of tobacco caterpillar was observed at vegetative stage of soybean crop when crop age was about 27 days old and incidence continued till the maturity of the crop, insect was damaging the foliage of soybean crop. Present findings are in accordance with Gangrade et al., (1975) they also reported that the tobacco caterpillar was a serious pest in Madhya Pradesh. Chaudhary, (2009) has also reported the dispersion of *Spodoptera obliqua* (Walker) and *Spodoptera litura* (F.) on soybean crop from August to October.

Formatted: Font: Italic

The results of tobacco caterpillar indicated that the maximum temperature showed significant positive correlation with Tobacco caterpillar population ( $r = 0.725$ ) and relative humidity also showed significant positive correlation with Tobacco caterpillar population ( $r = 0.627$ ). ~~whereas~~ Whereas, minimum temperature, and rainfall showed non-significant positive correlation with the Tobacco caterpillar population. and this result was in accordance with Patel (2012). During the peak population the maximum and minimum temperature were 30.93<sup>0</sup>C and 25.64<sup>0</sup>C respectively, Whereas relative humidity and rainfall were 95.57% and 07.79 mm respectively. This result was also in accordance with Kujur j. (2011).

#### **Green Semilooper, *Chrysodeixis acuta* (Walker) (Lepidoptera; Noctuidae)**

First incidence of green semilooper was observed in the vegetative stage when the crop age was about 25 days. The data showed that the pest was present on the crop during the entire cropping season *i.e.* first week of August to second week of September. The population of green semilooper increased and reached the peak level (3.0 larvae/mrl) on last week of August *i.e.* 34<sup>th</sup> standard week, subsequently, it declined gradually and disappeared during third week of September *i.e.* 37<sup>th</sup> standard week. In the present study the incidence of green semilooper was observed from first week of ~~august~~ August to second week of September and reached the peak during last week of August. Present findings are in accordance with Netam *et al.* (2013) and Garg *et al.* (2014).

The maximum temperature and relative humidity showed significant positive correlation with Green Semilooper population ( $r = 0.680$ ) and ( $r = 0.538$ ) respectively, while minimum temperature, and rainfall showed non-significant positive correlation with the Green Semilooper population. During the peak population of green semilooper the maximum and minimum temperature were  $30.57^{\circ}\text{C}$  and  $25.79^{\circ}\text{C}$  respectively, whereas relative humidity and rainfall were 94.43% and 14.00 mm respectively. The result confirms the findings of Kushram (2016). The maximum temperature and relative humidity showed significant positive correlation with Green Semilooper population, while minimum temperature, and rainfall showed non-significant positive correlation with the Green Semilooper population this result was in accordance with Patel (2012).

#### **Stem fly, *Melanagromyza sojae* (Zehntner) Diptera: Agromyzidae:**

First incidence of stem fly was observed immediately after seedling stage when the crop age was about 32 days old. From the table it is evident that the pest was present on the crop during the entire cropping season *i.e.* from first week of August to reproductive stage of the crop *i.e.* third week of September. The incidence of the stem fly was reached the peak (45.25 percent/mrl) at 33<sup>rd</sup> SMW. In the present investigation, the incidence of stem fly was recorded from first week of August to second week of September and reached the peak level during last week of August. Aske *et al.* (2007) also reported the similar results of stem fly incidence in soybean crop.

Comment [WU18]: Which table?

The results indicated that the maximum temperature showed highly significant negative correlation with stem fly population ( $r = -0.523$ ) while relative humidity and rainfall showed non-significant but positive correlation with stem fly population ( $r = 0.364$  and  $r = 0.366$ ). Minimum temperature showed non-significant but positive correlation with the stem fly population. At the time when the population of stem fly reach at peak, maximum and minimum temperature were  $32.64^{\circ}\text{C}$  and  $25.57^{\circ}\text{C}$  respectively, whereas relative humidity and rainfall were 88.43% and 03.57 mm respectively. This result matches with the finding of Singh and Singh (1992). The maximum temperature showed significant negative correlation with stem fly population it confirms the finding of Motaphale *et al.* (2017).

### **Girdle beetle, *Obereopsis brevis* (Swed.) (Coleoptera; Lamidae)**

First appearance of Girdle beetle was observed in the vegetative stage when the crop age was about 39 days old. The data showed that the pest was present on the crop from second week of August till reproductive stage of the crop *i.e.* third week of September. Girdle beetle infestation reached the peak (4.80 percent) at 35<sup>th</sup> SMW. The incidence of Girdle beetle was found from second week of August and it continued upto second week of October. The peak population of girdle beetle was observed in first week of September. Similarly, incidence of girdle beetle was also seen by Garg *et al.* (2014) and Kushram (2016).

The results indicated that the maximum temperature showed significant positive correlation with Girdle beetle population ( $r = 0.653$ ) and relative humidity showed highly significant positive correlation with Girdle beetle population ( $r = 0.529$ ) while, minimum temperature, and rainfall showed non-significant positive correlation with the Girdle beetle population. At the time of peak population of girdle beetle the maximum and minimum temperature were 30.93°C and 25.64°C respectively, whereas relative humidity and rainfall were 95.57% and 07.79mm respectively ([Table no](#)). This result confirms with the finding of Patel (2012).

### **Green stink bug, *Nezara viridula* (Linn.) (Hemiptera; Pentatomidae)**

Green stink bug first appeared during first week of August *i.e.* during vegetative stage from 31<sup>st</sup> standard week in soybean and the population of Green stink bug increased and reached the peak level (5.35 Green stink bug /mrl) on last week of August *i.e.* 34<sup>th</sup> standard week, subsequently, it declined gradually and disappeared during second week of September *i.e.* 36<sup>th</sup> standard week. Thus the population of green stinkbug ranged between 0.00 and 5.35 bug/mrl on soybean crop during the crop period. Incidence of green stink bug was from second week of August to first week of September and the highest population of green stink bug was observed during full developmental stages the result was in accordance with Smith *et al.* (2009).

The results indicated that the maximum temperature and relative humidity showed significant positive correlation with green stink bug population ( $r = 0.672$ ) and ( $r = 0.554$ ) respectively. However, minimum temperature and rainfall were non-significant, but these were positively correlated to the incidence of green stink bug. And at the time of peak population *i.e.* 5.35 bug infested /mrl. Maximum and

minimum temperature were 30.57<sup>0</sup>C and 25.79<sup>0</sup>C respectively, whereas relative humidity and rainfall were 94.43% and 14.00 mm respectively ([Table no](#)). This result was in accordance Gupta, P.K. (2006).

**Lady bird beetle, *Coccinella septempunctata* (Fabricius) (Coleoptera; Coccinellidae)**

Lady bird beetle was found active on soybean crop from first week of august to last week of ~~september~~September. The data indicated that Lady bird beetle activity started from vegetative stage of soybean crop and it was more active during reproductive stage of the crop. Lady bird beetle population reached peak (1.40/plant) at 35<sup>th</sup> SMW, declined subsequently and disappeared on 39 SMW. Lady bird beetle is the important predator in soybean crop. Lady bird beetle was observed from second week of august to last week of September and the highest population of lady bird beetle was observed during first week of September([Table no](#)). The result was more or less similar to Garg *et al.* (2014) and Suyal *et al.* (2018).

The results of lady bird beetle population on soybean crop indicated that the maximum temperature and relative humidity showed highly significant positive correlation with lady bird beetle population ( $r = 0.734$ ) and ( $r = 0.621$ ) respectively, while minimum temperature, and rainfall showed non-significant positive correlation with the lady bird beetle population. During the peak population of lady bird beetle, the maximum and minimum temperature were 30.93<sup>0</sup>C and 25.64<sup>0</sup>C respectively, Whereas relative humidity and rainfall were 95.57% and 07.79 mm respectively ([Table no](#)). ~~and the~~The present findings ~~is~~are more or less similar with Gupta (2006).

### **CONCLUSIONS**

On the basis of pest populations and their succession level, nine species of the insect pests i.e. aphid, whitefly, jassid, tobacco caterpillar, green semilooper, stem fly, girdle beetle and green stink bug with one natural enemy's lady bird beetle was observed during the crop season. The activity of aphid, whitefly, jassid, tobacco caterpillar, green semilooper, stem fly, girdle beetle, green stink bug and lady bird beetle was observed from the last week of ~~july~~July to second week of October. During the [study](#) period the Peak population of aphid (2.20/ aphids/ 3 leaves), whitefly

**Comment [WU19]:** The conclusion is almost a photo copy of the Abstract. It needs rewriting. What is the significant of this study for future management of these pests? There should be recommendation

(1.98/3 leaves), jassid (3.25 jassid/3 leaves) and tobacco caterpillar (8.15 larvae/mrl), green semilooper (3.00 larval/mrl), stem fly (5.20 percent), girdle beetle (4.80 percent/ plant), green stink bug (5.35/ mrl) and lady bird beetle (1.40/plant) was observed in between 31<sup>st</sup> to 35<sup>th</sup> SMW. With regards to correlation with abiotic factors, aphid, whitefly, jassid, tobacco caterpillar, green semilooper, girdle beetle, green stink bug and lady bird beetle population was found positively correlated, with maximum temperature, relative humidity and rainfall whereas stem fly found negatively correlated with maximum temperature.

#### REFERENCES

- Ahirwar, Rambihari, Devi, Payal and Gupta, Rajeev (2015). Seasonal incidence of major insect- pests and their biocontrol agents of soybean crop (*Glycine max* L. Merrill). *Academic Journals*, **10**(12): 402-404.
- Anonymous (2005). Directors Reports of experiments, National Research Centre for Soybean, Indore (M.P.). pp. 210-212.
- Anonymous (2007). Directors Reports of experiments, National Research Centre for Soybean, Indore (M.P.). pp. 225-226.
- Anonymous (2019). Annual Report on Soybean. Indian Institute of Soybean Research (ICAR), Indore (M.P.). pp. 2-4.
- Aske, S., Khandwe, N. and Singh, K. J. (2007). Incidence and damage of major pest of soybean in Madhya Pradesh. *Insect Environ.*, **12**(4): 156-159
- Bhattacharya A.K. and Rathore Y.S. (1979). Soybean insect problem in India. Proceeding of World Soybean Research Conference-II (Ed. Corbin, C.T.), March 26<sup>th</sup> to 29<sup>th</sup>, 1979, North Carolina State University, Raleigh, U.S.A. pp. 291-303.
- Chaturvedi, S., Singh, K.J., Singh, O.P. and Dubey, M.P., 1998, Seasonal incidence and damage of major insect pests of soybean in Madhya Pradesh. *Crop Res., Hissar.*, **15** (2/3): 260-264.
- Chaudhary, V. K. (2009). Dispersion of *Spilosoma obliqua* (Walker) and *Spodoptera litura* (F) in soybean field. *Pest Management and Eco. Zoology.* **17**(2): 133-139.
- Chaudhary, D.M., Chaudhary, N.J., Chaudhary, F.K. and Chauhan, R.P. (2018). Succession of pests and their natural enemies on soybean. *International Journal of Chemical Studies*, **6**(6): 2667-2673.

**Comment [WU20]:** There are some inconsistency in writing the references like 1. Journal name are italicized but not all  
2. author/s naming  
If possible add DOI for the references

- Gangrade G.A. (1976). Terminal Technical Report on the Project “Assessment of effects on yield and quality of soybean caused by major arthropod pests, *Department of Entomology, Jawaharlal Nehru Krish Vishwa Vidyalaya, Agricultural University, Jabalpur*. pp. 143.
- Garg, V.K.; Patel, Y. and Nayak, M.P. (2014). Response of *kharif* legumes to soybean girdle beetle, *Obereopsis brevis* (GAHAN). “ABSTRACT” Proceeding of Soycon - 2014 International Soybean Research Conference on “mitigating productivity constraints in soybean for sustainable agriculture held from 22-24<sup>th</sup> February, 2014. pp: 255.
- Gupta, P.K. 2006. Studies on seasonal incidence and effect of modern insecticides on major insect pests of soybean [*Glycine max* (L.) Merrill] M. Sc (Ag.) Thesis Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur.
- Kujur, J. (2011). Population dynamics of major insect-pests of soybean and management of defoliators and girdle beetle. M.Sc. (Ag.) Thesis, I.G.K.V.V., Raipur, India, **32**(2): 13-16.
- Kushram, T. (2016). Spectrum of insect pest complex and management of major insect pests on soybean (*Glycine max*). M.Sc. (Ag.) thesis Entomology, I.G.K.V., Raipur, **16**(1): 26-28.
- Lal S.S., Yadav C.P. and Dias C.A. (1981). Insect pests of pulse crops and their management. *Pesticide*, **21**: 66-67.
- Motaphale, A.A., Bhosle, B.B., Surwase, S.R. and Khan, F.S. (2017). Seasonal incidences of stem fly on soybean in relation to weather parameters. *Intl. J. adv. Sci. Engg. Tech.*, **5**(1): 69-71.
- Mundhe D.R. (1980). Insect pest complex on soybean, [*Glycine max* (L.) Merrill] in Marathwada Region. *Journal of Marathwada Agriculture University*, **5**(3): 259-260.
- Netam, H.K., Gupta R. and Soni, S. (2013). Seasonal incidence of insect pests and their biocontrol agents on soybean. *J. Agri. and Vet. Sci., (IOSR-JAVS)*, **2**(2): 11
- Patel, A. 2012. Studies on insect pest complex of soybean [*Glycine max* (L.) Merrill] and their management with bioagents. M. Sc (Ag.) Thesis. Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur.
- Ramesh Babu (2010). Literature on hepatitis (1984-2003): A bibliometric analysis. *Annals of Library and information Studies*, **54**: 195-200.

- Schmutz J., Cannon S.B., Schlueter J., Ma J. and Mitros T. (2010). Genome sequence of the palaeo polyploid soybean. *Nature*. 463: 178-183.
- Singh, K.J., and Singh, O.P. (1992). Influence of stem tunneling by the maggots of *Melanagromyza sojiae* (Zehnt) on yield of soybean. *J. Insect Sci.*, 5(2):198-200.
- Smith, J.F., R.G. Lutrell and J.K. Greene (2009). Seasonal abundance, species composition and population dynamics of green stink bug in production field of early and late soybean in south Arkansas. *J.Economic Ent.*, 102(1): 229-36.
- Suyal, Parul, Gaur, Neeta, Rukesh Pramod K.N. and Devrani, Ashish (2018). Seasonal incidence of insect pests and their natural enemies on soybean crop. *Journal of Entomology and Zoology Studies*, 6(4): 1237-1240.

Comment [WU21]: ?

**Table 1: Insect-pests recorded on soybean variety RVS-2007-6 at ----- during kharif season-2018 kharif season (for all tables)**

S. No.	Affected plant part	Common Name	Scientific Name	Order	Family
1.	Foliage	Aphid	<i>Aphis gossypii</i>	Hemiptera	Aphididae
2.		Whitefly	<i>Bemisia tabaci</i> ( <i>Gennadius</i> )		Aleyrodidae
3.		Jassid	<i>Empoasca kerri</i> ( <i>Pruthi</i> )		Cicadellidae
4.		Tobacco caterpillar	<i>Spodoptera litura</i> ( <i>Fabricius</i> )	Lepidoptera	Noctuidae

5.	Foliage, flowers & Pods	Green Semilooper	<i>Chrysodeixisacuta</i> (Walker)		
6.	Stem	Stem fly	<i>Melanagromyza-sojae</i> (Zehntner)	Diptera	Agromyzidae
7.		Girdle beetle	<i>Obereopsis brevis</i> (Swed.)	Coleoptera	Cerambycidae
8.	Foliage	Green stink bug	<i>Nezaraviridula</i> (Linn.)	Hemiptera	Pentatomidae

UNDER PEER REVIEW

**Table 2: Incidence of Insect pest complex of soybean during kharif season 2018**

SMW	Duration	Crop Age (days)	Crop stage	population/plant/3leaves			No. of larvae/mrl*		Stem fly		Girdle beetle inf (%)plant	Green Stink bug inf./mrl	Lady bird beetle/plant
				Aphid	White fly	Jassid	Tobacco caterpillar	Green semilooper	%plant inf./mrl	Stem tunneling (%)			
29	22-28Jul	18	0	0	0	0	0	0	0	0	0		
30	29Jul-4Aug	25	Vegetative	1.75	0.25	0	0	0.5	0	0	0	0	0
31	5-11Aug	32		2.2	1.2	0.75	0	1.75	20.3	2.25	0	0.95	0.25
32	12-18Aug	39		2	1.5	2.55	0.85	2.2	38.82	3.5	1.5	2.3	0.6
33	19-25Aug	46	Reproductive	1.8	1.75	3.25	1.6	2.5	45.25	5.2	2.4	3.25	0.85
34	26Aug-1Sep	53		1.15	1.98	2	3.25	3	40.28	4.5	3.5	5.35	1.1
35	2-8Sep	60		0.8	1.5	1.5	8.15	2.65	41.1	3.2	4.8	3.8	1.4
36	9-15Sep	67		0.25	1.25	1.15	4.5	1.2	31.2	1.75	3.25	0	1.2
37	16-22Sep-	74		0.2	1.2	0.9	0	0	18.27	0.8	2.75	0	0.95
38	23-29Sep	81		0.15	0.9	0.75	0	0	0	0	0	0	0.5
39	30Sep-6Oct	88	Maturity	0	0.8	0.5	0	0	0	0	0	0	0
40	7-13Oct	95		0	0.25	0.23	0	0	0	0	0	0	0

**Table 3: Influence of weather parameters on incidence of Insect pest complex of soybean during kharif season 2018**

SMW	Duration	population / plant / 3 leaves			No. of larvae/(mrl)		Stem fly		Girdle beetle infestation (%)	Green Stink bug infestation /(mrl)	Lady bird beetle/plant	Temp. ( <sup>0</sup> C)		Relative Humidity (%)	Rainfall (mm)
		Aphid	White fly	Jassid	Tobacco caterpillar	Green semilooper	%plant inf./mrl	Stem tunneling (%)				Max.	Min.		
28	15-21July	0	0	0	0	0	0	0	0	0	0	37.36	27.57	78.57	3.03
29	22-28 July	0	0	0	0	0	0	0	0	0	0	35.07	26	93.86	13.5
30	29 July - 4 Aug	1.75	0.25	0	0	0.5	0	0	0	0	0	31.14	25.5	90.71	10.9
31	5-11Aug	2.2	1.2	0.75	0	1.75	20.3	2.25	0	0.95	0.25	30.71	24.14	78.86	3.79
32	12-18Aug	2	1.5	2.55	0.85	2.2	38.82	3.5	1.5	2.3	0.6	32.07	25.43	83.29	6.57
33	19-25Aug	1.8	1.75	3.25	1.6	2.5	45.25	5.2	2.4	3.25	0.85	32.64	25.57	88.43	3.57
34	26Aug-1Sep	1.15	1.98	2	3.25	3	40.28	4.5	3.5	5.35	1.1	30.57	25.79	94.43	14
35	2-8Sep	0.8	1.5	1.5	8.15	2.65	41.1	3.2	4.8	3.8	1.4	30.93	25.64	95.57	7.79
36	9-15Sep	0.25	1.25	1.15	4.5	1.2	31.2	1.75	3.25	0	1.2	30	24.86	93.29	6.86
37	16-22Sep	0.2	1.2	0.9	0	0	18.27	0.8	2.75	2.25	0.95	32.43	24.43	83.71	0
38	23-29Sep	0.15	0.9	0.75	0	0	0	0	1.75	0.85	0.5	33.64	24.93	79.14	1.57
39	30 Sep-6 Oct	0	0.8	0.5	0	0	0	0	0.9	0	0	33.79	24.07	76.14	1.29
40	07-13 Oct	0	0.25	0.23	0	0	0	0	0	0	0	36.71	23.29	58.29	0

**Table 4: Correlation between pests' population and weather parameters during kharif season 2018**

S. No	Insect Pests	Weather parameters							
		Temperature ( $^{\circ}$ C)				Relative Humidity (%)		Rainfall (mm)	
		Min.		Max.		R	$b_{yx}$	R	$b_{yx}$
		r	$b_{yx}$	R	$b_{yx}$				
1.	Aphid	0.160 NS	--	0.627*	0.0107	0.369 NS	--	0.323 NS	--
2.	Whitefly	0.250 NS	--	0.690*	0.0023	0.543*	0.0361	0.410 NS	--
3.	Jassid	0.235 NS	--	0.458 NS	--	0.411 NS	--	0.450 NS	--
4.	Tobacco caterpillar	0.234 NS	--	0.725*	0.0025	0.627*	0.0120	0.220 NS	--
5.	Green Semilooper	0.242 NS	--	0.680*	0.0043	0.538*	0.0382	0.384 NS	--
6.	Stem fly	0.171 NS	--	-0.523*	0.0450	0.364 NS	--	0.366 NS	--
7.	Girdle beetle	0.235 NS	--	0.653*	0.0049	0.529*	0.0422	0.412 NS	--
8.	Green stink bug	0.243 NS	--	0.672*	0.0042	0.554*	0.0323	0.424 NS	--
9.	Lady bird beetle	0.233 NS	--	0.734*	0.0006	0.621*	0.0134	0.280 NS	--

\* Significant (at 5%), NS – Non-significant