

DETERMINATION AND EVALUATION OF HEAVY METAL CONCENTRATION IN SURFACE WATER AND SEDIMENT FROM ORASHI RIVER IN RIVERS STATE, NIGERIA.

ABSTRACT

The research was intended to study the influence of human and industrial activities on the Orashi River. Three groups are observed to have impacted the environment – Oil/gas industries, tyre burning from abattoir, untreated human and animal waste from settlers and the abattoir. The study was carried out from September 2019 to August 2020. The mean concentrations of the parameters studied in some samples were close to, or exceeded World Health Organization (WHO) and Federal Ministry of Environment (FMEnv) recommended limits for drinking water and seafood. The results from this study have provided information on the heavy metals profile on the surface water and sediment of the river. Essential heavy metals detected in water were in the range of Cu (1-5.4mg/l), Zn (1-4.4mg/l) and Fe (0.1 – 1.8 mg/l) which were within the WHO (1989), USEPA (1986), FEPA (1999) permissible limit of 3mg/l. The non-essential metals investigated were Cd, Cr and Pb but only Cd was detected. Cd levels (0.1-0.8mg/l) were slightly high, above 0.01mg/l above the permissible limit. Heavy metals in sediment were higher than those in the water body, occurring in the sequence Zn>Cu>Fe >Cd. The values ranges were Zn (1.4-5mg/kg), Cu (0.8-3mg/kg), Fe (0.4-4.9mg/kg) and Cd (0.5-1.8mg/kg). All relationship were statistically significant (p=0.05). With regards to seasonal variation, dry season recorded higher values than rainy season.

KEYWORDS: Metal, Determination, Evaluation, Industries, Seafood, Contamination.

Comment [MOU1]: How does tyre burning (smoke) gets into river water pollution. Why is tyre burning related to abattoir?

Comment [MOU2]: Nigerian Federal Ministry...???

Comment [MOU3]: Would this be the only conclusion that gets to the abstract?

INTRODUCTION

BACKGROUND INFORMATION

Water is the most important natural resources and there are many conflicting demand for them. Skillful management of water bodies is required if they are to be used for such diverse purpose as domestic and industrial supply, crop irrigation, transport, recreation, sports, commercial fisheries, power generation and waste disposal. Water bodies are vulnerable to contamination accident and bioterrorism attacks because they are relatively

unprotected, easily accessible, often isolated and their various use by human pre-disposes them to contamination. (Gullick *et al*; 2003). Environmental exposure to toxic metals is a critical issue in environmental and public health.

The modern world is aware of the relationship between water and water borne diseases, as a vital public health issue. (WHO 1989, Olashansky, *et al*, 1997., Asonye *et al*, 2007.

Heavy metals are well known pollutants in aquatic systems where industrial wastes are discharge, petroleum production and refining, gas flaring, gas processing plants and conveyance pipelines (Okoye *et al*, 1991, Izonfuo and Bariwanli 2001, Olarije 2003). Gas flares are operated in relatively uncontrolled manner in Niger Delta region. These effluents discharge and atmospheric emissions from flow stations and refineries often settle in the aquatic environment. When contamination reaches levels in excess of the assimilative capacity of the receiving waters, they may affect the survival, reproduction capacity, growth and behavioral condition of organisms. (Asonye *et al*, 2007).

In the attempt to define and measure the effect and presence of pollutants on aquatic ecosystems, biomarkers or bio-indicators have attracted a great deal of interest. The increase level of heavy metals in human has often been traced to heavy metal contamination in the aquatic system (Schuwerack *et al*, 2009).

The heavy metals of interest include the non-essential trace element Cd, Cr, and Pb) and the essential metals such as Cu, Zn, and Fe which have important biochemical functions to the organism at very low concentration (Rainbow 2007). These heavy metals are blacklisted in EEC Directive as dangerous substances in the aquatic environment. These heavy metals can be hazardous to human, even in very small amount. They are taken up by aquatic organisms and passed up the food chain through the process known as bio-magnification. Species varies in their degree of tolerance with the result that under polluted conditions, a reduction in species diversity is the most obvious effect (Edokpayi *et al* 2000). The intense industrial activity in Niger Delta region has attracted a lot of research interest but no systematic study has so far been conducted in Orashi River of ONELGA to ascertain its level of metal concentration and the health effects on the inhabitants. This may be attributed but

Comment [MOU4]: Please, specify that you are talking about "interest in this study". The toxic metals list is much longer.

It is not clear why this study would be interested only on these few metals.

Comment [MOU5]: Need a quote and reference here.

Comment [MOU6]: "much" instead of "a lot of"???

not limited to the restive nature of the inhabitants of the area as researchers may be vulnerable to attack by the youths who always vent their anger on soft targets.

It is therefore very important to conduct this study in view of the rapid growth of population, exploration and exploitation of natural resources, lack of environmental regulations, industrialization, urbanization, clearing of bank vegetations, annual dredging of the river to contain yearly flooding, construction of roads and bridges, drains and embankment walls and Agricultural activities of the inhabitant of Ogba/Egbema/Ndoni LGA (ONELGA) to record the range of the concentration within commercial as well as non-commercial species so that assessment about potential hazardous levels from the human nutritional standpoint can be made.

AIM: To determine the concentrations of heavy metals in the surface water and sediment from Orashi River. This is to enable the assessment of potential hazardous levels from human nutritional stand point.

MATERIALS AND METHODS

THE STUDY AREA

The study area is Ogba/Egbema/Ndoni LGA in Rivers State of Nigeria. (Fig: 1a, b & c). The area has a number of oil wells and major flow stations within the Niger Delta region of Nigeria. Nigerian Agip Oil Company (NAOC) and Total E & P Nigeria Limited explores, exploit crude oil and flare gases indiscriminately in three locations - Ebocha, Obrikom and Obitte.

The inhabitants of the area are predominantly farmers and fishermen which is their basic source of livelihood. The area has a growing population of 283,294 in 2006 and a projection of 398,000 in 2016 (National population commission of Nigeria (web), National Bureau of statistics (web)).

The site of the study is Orashi river, a non-tidal freshwater of the lower Niger basin that runs through some communities in Imo State, Egbema, Ndoni and Ogba communities in Rivers State. The river is a freshwater swamp forest river with several tributaries and originate from River Niger and empties into Sombrieiro river in Ahoada. The area is

Comment [MOU7]: This appears as a political statement, rather than a sound explanation. Is there much funding for this type of research, for example?

May I suggest this paragraph to be removed?

Comment [MOU8]: This "(never closes; I mean, there is no corresponding ")"

tropical with two seasons- the rainy (April – October) and dry season (November – March) which are usually flooded in the rainy seasons.

Fig 1: maps showing the study area




Fig 1a. Map of Nigeria Showing Rivers State



Fig.1b Map of Rivers State Showing Ogba/Egbema/Ndoni L.G.A

Comment [MOU9]: Please revise this figure. It does not get to current standards.

Fig.1c Map of Ogba/Egbema/Ndoni L.G.A



SAMPLING STATIONS

A reconnaissance survey was carried out in the study area on November 2018 and then sampling stations were established at five locations along the Orashi River, 5km distance from each other using Global positioning system navigator (GPS) as shown in Table 1 and represented by station 1 – 5.

Table 1: Geographical positioning system (GPS)

STATIONS	LOCATIONS	COORDINATES	ELEVATION
----------	-----------	-------------	-----------

Station 1	Okwuzi	N05°29'08.3" E006°42'26.3"	21M
Station 2	Ebocha	N05°27'49.3" E006°42'06.6"	24M
Station 3	Ndoni	N05°27'24.6" E006°40'27.8"	12M
Station 4	Obrikom	N05°23'31.0" E006°39'03.0"	22M
Station 5	Omoku	N05°20'18.7" E006°38'34.6"	16M

FIELD STUDY

Water and Sediment were collected from the 5 sampling stations monthly, beginning from September 2019 during rainy and dry seasons respectively. The sampling of the 5 stations started at about 8.30am each day

FIELD COLLECTION OF WATER SAMPLES

Water samples were collected from all stations at a depth of 15-25cm with pre-rinsed containers. Duplicate samples of surface water from the five (5) sampling station were collected for twelve (12) sampling periods (September 2019 to August 2020).

Water sample were collected with pre-rinsed 100ml plastic container and treated with 2ml concentrated HNO₃ in order to stabilize the oxidation state of the metals and stored in a refrigerator. (APHA-AWWA-WEF, 2005).

COLLECTION OF SEDIMENT SAMPLE

Sediment samples were collected at each of the five (5) sampling stations for the specified period using BENTHIC GRAB of about 1.0cm³ in replicate for each sampling station for each sampling period. The sample were put in a polythene bag, kept in a cooler and taken to the laboratory deep freezer (°C) prior to sediment analysis.

Comment [MOU10]: was??

Comment [MOU11]: Please specify the temperature

LABORATORY ANALYSIS OF SAMPLES

In the laboratory, the following analysis was done on the duplicate samples collected.

Comment [MOU12]: performed?

HEAVY METAL ANALYSIS IN SURFACE WATER (Fe, Cd, Pb, Cu, Cr and Zn)

Samples previously collected in the field and fixed with 2 drops of concentrated HNO₃ in order to preserve the metals and also to avoid precipitation were used for heavy metals analysis.

Metal analysis was performed using Unicam 969 Atomic Absorption spectrophotometer (AAS) in accordance with (APHA 1998). This involves direct aspiration of the sample into an air/acetylene or nitrous oxide/acetylene flame through which a light source emitting a narrow spectral line of characteristic energy. A hollow cathode lamp peculiar only to the metal under investigation was used to excite the free atoms of metal of interest in the flame. The concentration of the excited metal atoms calculated by comparison with a standard curve of the metal. Prior to analysis, the AAS

SEDIMENT SAMPLE ANALYSIS

The sediment samples were air-dried at room temperature for 5 days and grounded into powdery form.

1g of each was digested with Equia-Regia (mixture of HCL and HNO₃ in the ratio 3:1). The digestates were filtered with 20ml of de-ionized water and the filtrates were stored in clean acid-washed and appropriately labeled 30ml polyethylene sample containers for analysis by Atomic Absorption Spectrometric method using Atomic Spectrometer (Model 210 VGP Buck Scientific, USA).

Comment [MOU13]: aqua???

DATA ANALYSIS

All statistical analysis and presentation of results was done using Microsoft excel and Minitab 16 software.

Raw data was subjected to a two way analysis of variance (ANOVA) with replication using MINITAB.

Comment [MOU14]: This is redundant. You could write "minitab" only once without any loss of comprehension.

RESULTS OF HEAVY METALS IN SURFACE WATER

The following heavy metals were monitored during the study period – Chromium (Cr), Cadmium (Cd), Lead (Pb), Copper (Cu), Zinc (Zn) and Iron (Fe). Of all these metals, chromium and lead was not detected during the sampling period.

CADMIUM (Cd) IN SURFACE WATER

The mean values recorded in appendix 1 are station 1(0.35mg/l), station 2(0.53mg/l). Station 3 (0.4mg/l), Station 4(0.46mg/l) and station 5(0.27mg/l) and represented in fig 1. The values recorded show that it ranged from 0.1 to 0.8mg/l.

Seasonal variations in cadmium levels in water are presented in Table 3 and showed that dry seasons has higher value (0.43mg/l) than Rainy season (0.37mg/l).

The analysis of variance (ANOVA) result showed that there was significant difference (P<0.05) observed for the water sample during the monitoring period (P=0.000).

COPPER IN SURFACE WATER

The mean levels of Cu is presented in appendix 1- station 1(2.73mg/l), station 2(4.08mg/l), station 3(1.2mg/l), station 4(1.4mg/l) and station 5(0.91mg/l) and represented in fig 2. The value ranged from 1.0 to 5.4mg/l. The highest value recorded was in February and March in station 2.

Seasonally, the result in table 3 showed that Rainy season (2.24mg/l) recorded a higher value while the Dry season recorded a lower value (1.89mg/l) during the sampling period.

The analysis of variance (ANOVA) results showed there is significant difference (P< 0.05) for copper concentration in the water sampled during the monitoring period (P= 0.000).

Comment [MOU15]: Difference between what values? Is it comparing one station to the other? Comparing one month to ne next? Comparing depth concentration? This is unclear and needs a mor rigorous description

Comment [MOU16]: Have you looked into temperature related solubility?




Fig 2: Cadmium (Cd) in surface water




Fig 3: Copper (Cu) in surface water

ZINC IN SURFACE WATER

The mean concentration of zinc in water during the study period is in appendix 1 thus station1 (2.16), station 2(1.67), station 3(3.52), station 4(2.2) and station 5(1.35mg/l) and represented in fig 3. The result indicates that zinc range from 1.0 to 4.4mg/l in the surface water. The highest level of zinc was observed in the month of March followed by February. Seasonally, higher values were recorded for rainy season (2.31mg/l) while lower values (1.97mg/l) were recorded for dry season.

The analysis of variance (ANOVA) showed that there was significant difference ($P < 0.05$) between location and time.

IRON (Fe) IN WATER

The result obtained for iron is presented in appendix 1 and represented in fig 4. The result indicates that iron range from 0.1 to 1.8. The mean value levels were station 1(0.49), station 2(0.75), station 3(0.55), station 4(1.08) and station 5(1.25).

Seasonally, higher values (0.86mg/l) were recorded for dry season while lower levels (0.79mg/l) were recorded for rainy season. The highest level of iron was observed in the month of March followed by April, February and January.

The analysis of variance (ANOVA) shows that there is significant difference ($P < 0.05$) between locations and time. $P=0.000$.

Comment [MOU17]: ????




Fig 4: Zinc in Surface water.




Fig 5: Iron (Fe) in Surface water.

HEAVY METALS IN SEDIMENT

The following heavy metals were sampled from the 5 stations and analyzed during the monitoring period:

CADMIUM (Cd) IN SEDIMENT.

The concentration of Cadmium recorded in sediment is presented in appendix 2 and shown in fig. 5. The results indicate that Cadmium level in this study ranged from 0.5 to 1.8mg/kg in all the stations. The mean values obtained were station 1(0.805), station 2(1.42), station 3(1.04), station 4(0.83) and station 5(0.73mg/kg).

Seasonal variations as presented in table 4 show that total Cadmium level was higher during the dry season (1.14mg/kg) than the rainy season (0.99mg/kg).

The analysis of variance (ANOVA) result showed that there is significant difference ($p < 0.05$) in time ($P = 0.000$).

Comment [MOU18]: This is letter o but it should be number 0

COPPER (Cu) IN SEDIMENT.

The concentration of Copper recorded in sediment is presented in appendix 2 and shown in fig. 6. The results indicate that Copper level in this study ranged from 0.8 to 3mg/kg in all the stations. The mean values observed for copper was station 1(1.93), station 2 (2.70), station 3(1.24), station 4(1.48) and station 5(1.24mg/kg).

Seasonal variations show that Copper level was higher during the rainy season (1.90mg/kg) than the dry season (1.8mg/kg).

The analysis of variance (ANOVA) result showed that there was significant difference ($p < 0.05$) between locations and time ($P = 0.000$).




Fig 6: Cadmium (Cd) in Sediment




Fig 7: Copper in sediment

ZINC (Zn) IN SEDIMENT.

The concentration of Zinc recorded in sediment is presented in appendix 2 and shown in fig.7. The results indicate that Zinc level in this study ranged from 1.4 to 5mg/kg in all the stations. The mean values observed for Zinc was station 1(3.68), station 2(2.24), station 3(4.49), station 4(2.42) and station 5(1.91mg/kg).

Seasonal variations recorded in table 4 show that Zinc level was higher during the rainy season (2.74mg/kg) than the dry season (2.72mg/kg).

The analysis of variance (ANOVA) result showed that there was significant difference ($p < 0.05$) between locations and time ($P=0.000$).

IRON (Fe) IN SEDIMENT.

The concentration of Iron recorded in sediment is presented in appendix 2 and shown in fig. 8. The results indicate that Iron level in this study ranged from 0.4 to 4.9mg/kg in all the stations. The mean values observed for Iron was station 1(1.37), station 2(1.39), station 3(2.17), station 4(1.84) and station 5(1.62).

Seasonal variations show that Iron level was higher (1.91mg/kg) during the dry season than the rainy season (1.41mg/kg).

The analysis of variance (ANOVA) result showed that there was significant difference ($p < 0.05$) between locations and time ($P=0.005$).




Fig 8: Zinc in Sediment




Fig 9: Iron (Fe) in Sediment

DISCUSSION

HEAVY METALS IN WATER

The essential heavy metals detected in the surface water in the present study were copper (Cu), Zinc (Zn) and Iron (Fe) and the non-essential heavy metals is cadmium (Cd).

The values obtained from the 5 stations for the 4 heavy metals detected vary in composition in different station and at different time (month).

CADMIUM (Cd) IN WATER

Cadmium (Cd) was widely distributed in low levels in the environment. The mean value recorded ranged from 0.29 – 0.8 with specific mean record of station 1 (0.55), station 2 (0.53), station 3 (0.4), station 4 (0.46) and station 5 (0.29). These values of Cd are high compare to FMEnv, WHO/USEPA limit of 0.1mg/l. This high cadmium concentration differs from the low metal concentration reported by earlier workers which include Obire *et al*, (2003) on Elechi creek, Omoigherale and Ogbeibu, (2005) on Osse River, Southern Nigeria and Ekweozor and Agbozu (2001)

This may be attributed to intense oil exploration and exploitation in the area which ranked ONELGA as the highest oil producing LGA in Nigeria. These activities involved the use of dynamites, chemicals; effluent discharges in the area which will incidentally washed into the river during rain.

COPPER (Cu) IN WATER

The values of copper ranged from 0.1 – 5.4mg/l with specific mean record of 2.73, 4.08, 1.2, 1.4 and 0.91mg/l for station 1, 2, 3, 4 and 5 respectively. The results also show that dry season recorded a higher value than the rainy season. Copper is an essential element observed in low levels in stations 3, 4 and 5. Notwithstanding, the discharge of waste containing copper should be discouraged since its presence in the environment is hazardous, as copper salts are poisonous. Station 2 recorded high level (4.08mg/l) which is above the 3mg/l FMEnv (1999) maximum permissible limit. The low copper concentration in some stations is in agreement with low metal concentration reported by Obire *et al*, (2003) on Elechi creek, Ekweozor and Agbozu (2001) in Azuabie creek.

ZINC (Zn) IN WATER

The concentration of Zinc in surface water ranged from 1 – 4.4mg/l during the study period with specific mean values of station 1(2.16), station 2(1.67), station 3(3.52), station 4(2.2) and station 5(1.35) mg/l.

Rainy season recorded higher values (2.31mg/l) than dry season (1.97mg/l) with the highest value (3.5mg/l) in March followed by February and January in station 3. The low zinc concentration in stations 1, 2, 4 and 5 is in agreement with international standards – WHO (1989), USEPA, (1986) and FEPA (1999) permissible limit of 3mg/l. The high level of Zn (3.52mg/l) in station 3 may be attributed to the tyre ash in abattoir located near the Shore of Orashi River in station 3. Zinc makes up 51.48% of tyre ash (Bay *et al*, 2005). This indicate an impact from runoff of tyre ash and animal blood which is rich in zinc and Iron

Comment [MOU19]: There could not be "agreement"; the author might prefer to talk about "below limits", or expressions of that sort.

Comment [MOU20]: Is there a basis for this observation?

IRON (Fe) IN WATER

The concentration of Iron in surface water ranged from 0.1 – 1.8mg/l during the study period with specific mean values of station 1(0.49), station 2(0.75), station 3(0.55), station 4(1.08) and station 5(1.25) mg/l.

Dry season recorded higher values (0.86mg/l) than rainy season (0.79mg/kg) with the highest value observed in the month of March in station 5. The low Iron concentration in all stations is in agreement with international standards – WHO (1989), USEPA, (1986) and FEPA, (1999) permissible limit of 3mg/l. The presence of Iron in the stations may be attributed to the tyre ash in abattoir located near the Shore of Orashi River and other anthropogenic activities in the area. Iron makes up 6.33% of tyre ash. This indicates an impact from runoff of tyre ash and animal blood which is rich in Iron Nzenwa *et al*, 2018).

Comment [MOU21]: ibid

SEDIMENT ANALYSIS IN SEDIMENTS

Sediment is an important component of aquatic ecosystem because it provides a habitat for a wide range of benthic and epi-benthic organisms (Adeyemo *et al*, 2008) and act as sink to contaminants and pollutants. There is exchange between sediments and the overlying water which depends on the chemical characteristics of water and sediment both phases (Adeyemo *et al*, 2008).

Therefore, exposure to certain substances in sediments represents a potentially significant hazard to the health of these organisms. An assessment of this hazard would require an understanding of the relationships between concentration of chemicals in sediment and occurrence of adverse biological effects.

Comment [MOU22]: Even if the health of these organisms is not under threat, the metal bioaccumulation would endanger those organisms that are higher in the food chain!!!

The heavy metals analyzed for in the sediments ~~are~~ were Cadmium, Copper, Lead, Zinc and Iron.

Comment [MOU23]: Effects that must include bioaccumulation and trophic level.

CADMIUM (Cd) IN SEDIMENT

The results obtained in this study showed a range of 0.5 – 1.8mg/kg and mean values of station 1(0.805), station 2(1.42), station 3(1.04), station 4(0.83) and station 5(0.73) mg/kg.

The level recorded in sediment is higher than that recorded in water and exceeded the maximum level of 0.01mg/l allowed for discharge into inland waters (EPA, 1991). The levels were higher than the reported values of below detection limit (BDL) and 0.080mg/kg in Taylor creek (Agbozu, 2001).

Cadmium is toxic even at trace level and has been identified as one of the trace metals contained in crude oil in varying concentrations and has been shown to be capable of accumulating in food chain (Nwadinigwe and Nworgu (1999).

Seasonally, Cadmium level were generally higher in the dry season (**1.14mg/kg**) than rainy season (**0.99mg/kg**) and exceeded sediment toxicity guideline value.

Cadmium has both long and short term effects on specific tissues/organs of the body. Kidney and ocular damages are long term effects while testicular damage is short term (Asonye *et al*, 2007)

Comment [MOU24]: Mammals? Plants? Benthonic? Bacteria? Please be more precise as what you mean.

COPPER (Cu) IN SEDIMENTS

The level of copper recorded in this study were relatively high and ranged from 0.8 – 3.0mg/kg and mean values of stage 1(1.93), stage 2(2.70), stage 3(1.24), station 4(1.48) and stage 5(1.24) mg/kg. The level recorded in sediment is higher than that recorded in water but is within the FMENV recommended limit of 3mg/kg allowed in sediment.

Copper is not often considered a threat to human health except when present at abnormal values.

Seasonally, the levels were slightly higher in the rainy season (1.90mg/kg) than dry season (1,8mg/kg) and are within sediment guideline value (Asonye *et al*, 2007).

ZINC (Zn) IN SEDIMENTS

The level of Zinc recorded in this study ranged from 1.4 – 4.5mg/kg and mean values of station 1(3.68), station 2(2.24), station 3(4.49), station 4(2.42) and station 5(1.91) mg/kg. The level recorded in sediment is higher than that recorded in water and are within the FMENV recommended limit of 3mg/kg allowed in sediment except in station 4 (4.49mg/kg) which exceeded sediment toxicity guideline value. This could be attributed to effluents from abattoir with tyre ash and residue as well as animal blood waste which have high percentage of Zinc (Nzenwa *et al*, 2018). Zinc is not often considered a threat to human health except when present at abnormal values.

Seasonally, the values were slightly higher in the rainy season (2.79mg/kg) than observed in dry season (2.72mg/kg).

IRON (Fe) IN SEDIMENTS

The concentration of Iron recorded in sediment ranged from 0.4 – 4.9mg/kg and mean values of station 1(1.37), station 2(1.39), station 3(2.17), station 4(1.84) and station 5(1.62) mg/kg.

These values recorded in sediments are within the FMENV recommended limit of 3mg/kg allowed in sediment. It is interesting to note that Iron is one of the beneficial metals and can become toxic if optimum level of 3mg/kg is exceeded.

Seasonally, Iron levels were generally higher during the dry season (1.91mg/kg) than observed in rainy season (1.41mg/kg). This could be attributed to sedimentation of pollutants from continuous leaching of rusty iron fragment from pipeline, tyre burning sites in abattoir, residue and animal blood waste.

Statistically, there is significant different ($p < 0.05$) between locations and time ($p = 0.005$).

Comment [MOU25]: difference?

CONCLUSION

The research was intended to study heavy metal concentration as a result of the influence of human and industrial activities on the Orashi River and to provide information on the heavy metals profile of the river. The average mean concentrations of the heavy metals studied in some samples were close to or exceeded World Health Organization (WHO) and Federal Ministry of Environment (FMEnv) recommended limits for drinking water and seafood.

Essential heavy metals detected in water were Copper (Cu), Zinc (Zn) and Iron (Fe) and were low when compared with international standard (WHO 1989, USEPA 1986, FEPA 1999, FMEnv 1992) recommended limits of 3mg/l. The non-essential metals detected were Cadmium (Cd) which exceeded the international standard of 0.01mg/kg but do not constitute health risk. Heavy metals concentration was generally elevated at and around abattoir and gas flare sites.

The high level of Heavy metals in the water calls for concern as prolonged exposure can have some health – risk implication on the biota and humans who are the final consumers.

Comment [MOU26]: prolonged?

RECOMMENDATION

This study has shown that Heavy metals accumulate over time within the Orashi aquatic ecosystem. The Orashi River have been shown to be the major source of domestic water supply and protein supplement to the indigenous community who are exposed to the hazardous effects of these pollutants. It is therefore recommended that:

(i) The companies operating in the area should adopt improved waste management plans to reduce the levels of the pollutants discharge in the environment.

(ii) Comparative studies in other biota of the environment would be useful in monitoring the rate and mechanism of uptake of the metals in the food. This will provide data for informed decision on uptake of what is available in a polluted environment.

(iii) The local communities should be enlightened about the adverse effects of anthropogenic activities, oil pipeline vandalization/sabotage and bunkering activities now rampant in the communities to make money.

Comment [MOU27]: There is no data on "over time" phenomena. This seems to be a "one point in time" study. Please provide details.

Comment [MOU28]: has?

Comment [MOU29]: There is no data regarding this observation

Comment [MOU30]: There is no data on this observation. Please provide detailed evidence

REFERENCES

- Adeyemo, O.K. Adedokun, O.A Yusuf, R.K. and Adeleye E.A. (2008). Seasonal Change In physicochemical Parameters and Nutrient Load of River Sediments in Ibadan City Nigeria. *Global NEST Journal* 10 (3) 326-336
- Agbozu, I.E, Ekweozor. I.K.E & Opuene K (2007). Survey of heavy metals in the catfish *Synodontis clarias* *International Journal of Environmental science and Technology* 4(1) 93-97
- American Public Health Association (APHA,1988). Standard methods for the examination of water and waste water. 20th Edition, Washington DC, New York.
- APHA-WWWA-WEF (2005) Standard methods for the examination of water and waste water (20th ed) Greenberg A.E and Eaton, A.D (Eds). American Public Health Association, Alexandria, American water work Association, water Environment Federation, Washington DC
- Asonye, C.C. Okolie, N.P.Okenwa, E.E and Iwuanyanwu U.G (2007). Some physicochemical characteristics and Heavy metal profiles of Nigerian Rivers, Streams and water ways. *African Journal of Biotechnology*, 6(5):617-624
- Chukwu, L.O and Odunzeh, C.C (1993).Toxic and sub-lethal effect of spent lubricant oil on the gill morphology of the tropical Teleost- *oreochromis niloticus*. *Journal of Nigerian Environmental society* 1(3), 334-339.
- Edokpayi, C.A., Okenyi J.C., Ogbelibu A.E & Osimen E.C (2000). The effect of human activities on the macro-benthic invertebrate of Ibiekum stream Ekpoma, Nigeria. *Bioscience Research community*, 12, 79-87.
- European Economic Commission (EEC) (2005). Commission Regulation (EC) No.78/2005(pp.L16/43EL16 /45) *Official journal of European Union*.
- FAO (1983). Compilation of legal limits for hazardous substances in fish and fishery products. FAO Fishery Circular No 464. Food and Agriculture Organization; pp. 5- 100
- FAO//WHO (1989). Evaluation of certain food additives and the contaminant mercury, lead and cadmium; WHO Technical Report series No 505.
- FDA (2001). Fish and fisheries product Hazards and control guidance. 3rd Edition, Center for Food safety

and Applied Nutrition, US Food and Drug Administration
 Federal Environmental Protection Agency, (1999). Guidelines and standards for Environmental Pollution in Nigeria, FEPA. Lagos, Nigeria.
 FMEnv (2001). National Guidelines and standard for waste water quality in Nigeria. FMEnv. Lagos Nigeria

Formatted: Spanish (Chile)

Gullick, R.W., Grayman, W.M., Deininger, R & Males, R.M (2003). Design of early warning monitoring system of source waters. *Journal of American water work Association*, 95, 58-72.

Izonfuo, L.W.A. and Bariweni, P.A (2001). The effect of Urban Runoff water and human activities on some physico-chemical parameters of the Epie Greek in the Niger Delta. *Journal of applied science and Environmental management*, 5(10:47-55

Nwadingwe, C.A and Nwaorgu, O.N. (1997). Metal Contaminants in some Nigerian well- Head curdes. Comparative Analysis. *Journal of Chemistry society of Nigeria*, 24: 118- 121.

Nwaedozie, J.M. (1998). The Determination of Heavy Metal pollution in some fish samples from River kaduna. *Journal of Chemical society of Nigeria*, 23;21-23.

Nzenwa, P. O., Okorie, P.U., Ukaga, C.N., Chinedu-Eleonu, P.O., Obasi, C.C., Ibe, F.C., Akujobi, T.C., **Okwodu, N. E.** & Ugochukwu, M.G. (2018). Microbial and Heavy Metal Efficient Characteristics of Government Abattoir in Owerri Municipal, Imo State. *Global Research Journal of Science* ISSN: 2270-8300. GRJS Online

Obire, O., Tamuno, D.C & Wemedo, S.A (2003). Physico-chemical quality of Elechi Creek in Port-Harcourt Nigeria. *Journal of Applied Science and Environmental Management*. 7, 43-49.

Okoye, B.C.O., Afolabi, O.A and Ajao, E.A. (1991). Heavy Metals in the Lagos Lagoon Sediments. *International Journal of Environment Studies*, 38:131- 136

Olajire, A.A., Ayodele, E.T., Onyedirdan, G.Q. and Oluyemi, E.A.(2003). Levels of speciation of Heavy Metals in soils of Industrial Southern Nigeria. *Environmental Monitoring and Assessment*, 85(2):135-155.

Olashan sky, S. Carnes, B., Roger, R and Smith, L. (1997). Infectious Diseases- New and Ancient threat to World Health. *Population Bulletin*, 52(2):2-43.

Omoigberale, M.O., Oboh, I. P., Erhunmwunse, N.O., Ezenwa, I.M & Omoruyi, S.O (2014). An assessment of the trace metal contents of Owen River, Edo state, Nigeria. *European International Journal of Science and Technology*. 3(5), 88-98

Rainbow, P.S. (2007). Trace metal bioaccumulation models, metabolic availability and Toxicity. *Environmental International*, 33, 576-582

Schuwercak, P.M.M., Lewis, J.W & Jones, P.W (2007) Interactive Cellular and Physiological Response in the Freshwater Crab; *Polamonautes Warreni* to cadmium and microbial gill infestation. *Ecotoxicological and Environmental safety*. 67(4), 302- 310

United State Environment Protection Agency (USEPA) (2002). Current drinking water standards. Office of Groundwater and Drinking water, Government printing office, Washington 19, 162-166.

World health organization(WHO) (1984). *Guidelines for drinking water quality* Geneva.

World Health Organization (WHO) (1988). *IPCS Environmental Health criteria 61 Chromium* WHO, Geneva. P2693-2698.

Appendix 1: Raw data of Heavy Metals obtained in water samples

Cd	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	0.4	0.6	0.2	0.2	0.6	0.6	0.3	0.2	0.3	0.2	0.3	0.3
St 2	0.8	0.8	0.3	0.3	0.8	1	0.4	0.3	0.4	0.4	0.4	0.4
St 3	0.5	0.6	0.2	0.2	0.7	0.6	0.3	0.3	0.3	0.3	0.4	0.4
St 4	0.7	0.4	0.4	0.3	0.8	0.5	0.4	0.4	0.4	0.4	0.4	0.4
St 5	0.3	0.3	0.1	0.1	0.4	0.4	0.2	0.2	0.3	0.3	0.3	0.3

Cu	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	1	2	1.2	2	3	3.2	3.8	3.4	3.5	3.3	3.2	3.2

St 2	2	2.5	1.8	3	5	5.4	5.4	5	5	4.8	4.6	4.5
St 3	0.6	0.7	0.68	0.8	1.2	1.2	1.24	1.18	1.6	1.6	1.8	1.8
St 4	1.2	1.18	1	1.2	1.3	1.4	1.4	1.3	1.8	1.74	1.7	1.6
St 5	0.8	0.8	0.6	0.8	0.9	1	1	0.9	0.9	1	1.2	1

Zn	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	2	1.8	1.2	1.4	2.54	2.1	2.6	2.2	2.5	2.52	2.4	2.6
St 2	1.4	1.4	0.8	1.12	1.8	1.6	2.42	2	1.9	1.95	1.6	2
St 3	3.6	3.8	2	2	3.9	4.2	4.4	4	3.7	3.6	3.6	3.4
St 4	1.8	1.7	1.4	1.4	2	1.8	2	1.6	2.2	2.3	2.4	2.4
St 5	1	1.2	1	1.2	1.3	1.4	1.6	1.4	1.4	1.5	1.6	1.6

Fe	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	0.2	0.24	0.18	0.2	0.8	1	1.2	1	0.24	0.28	0.28	0.24
St 2	0.5	0.6	0.3	0.5	1	1.02	1.3	1.18	0.6	0.7	0.7	0.6
St 3	0.3	0.3	0.1	0.25	0.85	0.9	1.04	1	0.4	0.5	0.5	0.4
St 4	0.9	0.94	0.6	0.6	1.2	1.4	1.6	1.5	1	1.12	1.12	1
St 5	1.14	1.2	0.8	0.84	1.4	1.6	1.8	1.6	1.14	1.2	1.2	1.14

Appendix 2: Heavy Metal in Sediment

Cd	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	0.6	0.6	0.4	0.7	0.8	1	1.16	1.1	0.8	0.8	0.9	0.8
St 2	1.2	1.18	1	1.2	1.6	1.76	1.8	1.5	1.4	1.4	1.6	1.4
St 3	0.8	0.8	0.6	0.9	1	1.2	1.4	1.38	1	1	1.2	1.2
St 4	0.9	0.8	0.8	0.7	0.9	1.14	1.2	1.2	0.9	0.9	0.9	0.8
St 5	0.5	0.6	0.5	0.5	0.8	1	1.15	1	0.6	0.6	0.8	0.7

Cu	Sept	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
St 1	1.2	2.2	2.18	2.24	2.3	2.4	2.5	2.48	1.3	1.4	1.4	1.5
St 2	2.4	2.7	2.66	2.7	2.84	2.9	3	3	2.5	2.6	2.6	2.6
St 3	0.8	1.14	1	1.2	1.26	1.4	1.6	1.8	1	1.2	1.2	1.3
St 4	1.4	1.24	1.2	1.26	1.34	1.6	1.8	1.46	1.6	1.6	1.6	1.6
St 5	1	1	0.9	1	1.2	1.5	1.7	1.6	1.2	1.2	1.2	1.4

Zn	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	2.6	2.62	2.5	2.6	2.7	2.76	2.8	2.7	2.7	2.7	2.7	2.8
St 2	2	2	1.8	2	2.4	2.48	2.5	2.4	2.2	2.3	2.4	2.4
St 3	4.2	4.3	4.2	4.2	4.5	4.8	5	4.6	4.4	4.5	4.6	4.6
St 4	2	2	2	2.1	2.6	2.9	3.18	3.2	2.2	2.2	2.3	2.4
St 5	1.4	1.46	1.4	1.4	2	2.3	2.5	2.5	1.6	1.8	1.8	1.8

Fe	SEPT	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG
St 1	0.5	0.54	0.6	0.6	2.7	2.76	2.8	2.7	0.7	0.8	0.8	0.9
St 2	0.8	0.9	0.4	0.9	2.4	2.48	2.5	2.5	0.9	1	0.9	1
St 3	0.6	1.24	0.9	0.9	4.5	4.8	4.9	4.8	0.8	0.9	0.8	0.9
St 4	1.2	1.5	1.1	1.24	2.6	2.9	3	3.2	1.3	1.3	1.4	1.3
St 5	1.44	0.48	0.5	1.26	2	2.3	2.5	2.6	1.5	1.6	1.6	1.6

Appendix 3: Seasonal Variation in Heavy Metal Concentration in water

Heavy Metal	DRY SEASON							RAINY SEASON						
	OCT	NOV	DEC	JAN	FEB	MAR	\bar{x} conc.	APR	MAY	JUN	JUL	AUG	SEP	\bar{x} conc.
Cd	0.54	0.24	0.22	0.66	0.62	0.32	0.43*	0.28	0.34	0.32	0.36	0.36	0.34	0.37
Cu	1.44	1.07	1.56	2.28	2.44	2.57	1.89	2.36	2.56	2.49	2.5	2.42	1.12	2.24*

Zn	1.98	1.28	1.42	2.31	2.22	2.60	1.97	2.24	2.34	2.37	2.32	2.6	1.96	2.31*
Fe	0.66	0.4	0.48	1.05	1.18	1.39	0.86*	1.26	0.68	0.76	0.76	0.68	0.61	0.79

Appendix 4: Seasonal Variation in Heavy Metal Concentration in Sediment

Heavy Metal	DRY SEASON							RAINY SEASON						
	OCT	NOV	DEC	JAN	FEB	MAR	\bar{x} conc.	APR	MAY	JUN	JUL	AUG	SEP	\bar{x} conc.
Cd	0.78	0.66	0.8	1.02	1.22	1.34	1.14*	1.2	0.94	0.94	1.08	0.98	0.8	0.99
Cu	1.66	1.59	1.68	1.79	1.96	2.12	1.8	3.65	1.52	1.6	1.6	1.68	1.36	1.90*
Zn	2.40	2.38	2.46	2.84	3.05	3.20	2.72	3.08	2.62	2.7	2.8	2.8	2.44	2.74*
Fe	0.93	0.7	0.82	2.84	3.05	3.14	1.91*	3.16	1.02	1.12	1.1	1.14	0.91	1.41

UNDER PEER REVIEW