

Original Research Article

QUALITY EVALUTION OF NIGERIA BASE -MASA FROM BROKEN RICE ENRICHED WITH AFRICAN YAM BEAN AND CRICKET FLOUR BLENDS

ABSTRACT

Production and nutritional quality of Nigeria base -*Masa* from broken rice enriched with African yam bean and cricket blends was investigated. Broken rice, African yam bean and cricket were processed to flour using standard procedures. Blends of different proportions of broken rice, African yam bean and cricket were prepared at 100 %, 80/0/20 %, 80/20/0 % and 80/10/10 % respectively. The control sample was 100 % broken rice. The proximate, vitamins and minerals, antinutrients contents and sensory attributes of the *Masa* products were determined. The data obtained showed that moisture, protein, fat, ash carbohydrates and energy contents ranged from 38.46-44.48 %, 5.60-15.32 %, 0.71-9.71 %, 0.51-3.17 %, 0.61-0.79 %, 26.71-54.13 % and 245.25-270.77 kcal/100g respectively. The range values for vitamins and minerals are β -carotene (0.34-12.33 μ g/100g), vitamin C (0.52-10.61 mg/100g), potassium (24.93-37.87 mg/100g), calcium (142.03-284.11 mg/100g), phosphorus (64.89-127.71 mg/100g), and iron (22.06-50.67 mg/100g). The antinutrients ranged from 0.095-1.860 mg/100g for alkaloids, 0.535-2.600 mg/100g for tannins, 0.040-1.550 mg/100g for phytates and 0.015-0.285 mg/100g for oxalates. The colour, aroma, taste, texture and general acceptability of the sensory attributes results ranged from 6.13-7.73, 6.60-7.07, 6.27-7.33, 5.93-6.73 and 6.67-7.73 respectively.

Keywords : broken rice; african yam bean; cricket; masa; vitamins; minerals.

1. INTRODUCTION

Masa is a fermented bread-like product, which is round in shape with brown smooth boy and crippling edges, made in Nigeria from pearl millet, maize or rice flour [1]; [2]; [3]. *Masa* is one of the varieties of fermented cereal-based foods and is a good source of income for the women who prepare this traditional product for sale [1]. '*Masa*' serves as breakfast and snack item [3]. *Masa*' is consumed in various forms by all aged groups in the middle belt and northern states of Nigeria. '*Masa*' results from frying of the fermented rice dough which is round in shape with brown smooth body and crippling edges.

Rice is a cereal which is used mainly for human food. The milling process of rice causes some loss of nutrients and breakage of some rice resulting in "broken rice." Little attention has been paid to the use of broken rice in Nigeria. Indeed, it is underutilized and often referred to as "poor man's food" in some societies because it is cheaper than unbroken rice and purchased mainly by those who cannot afford unbroken rice [4]. The use of flour from broken rice in products could significantly increase the utilization of broken rice grains; thereby reducing losses that are normally incurred with the locally processed rice grains in Nigeria.

Cricket powder may be used as a protein-rich additive. It should be pointed out that it is cheap and easy to produce, and is considered a good source of dietary protein [5]. Crickets are a valuable source of protein, unsaturated fatty acids, dietary fiber, vitamins and minerals [5]; [6]. Cricket powder obtained from these insects is characterized by high protein content (about 60–70 %), a lack of carbohydrates, and high iron and calcium content [6]. A FAO report explicitly recommends the use of insects as a source of easily digestible protein, especially in the face of rapid population growth in the world and the growing difficulty of providing sufficient food [7].

Comment [DM1]: Rice was replaced 2 times as research and control. Please correct.

Comment [DM2]: Please enter the name *Masa* with or without quotation marks. In my opinion, this is a proprietary name of the product, therefore the quotation marks are not needed.

Comment [DM3]: It was in the first sentence of the paragraph.

African yam bean is an herbaceous leguminous plant occurring throughout tropical Africa (United States Department of Agriculture [8]). African yam bean has attracted research interest because of its nutrient content. Amino acid analyses indicate that the lysine and methionine levels in the protein are equal to or better than, those of soybeans while most of the other essential amino acids corresponds to WHO/FAO recommendation. A protein content ranging between 20.2 and 21.2% has been reported for African yam bean by [9]. Its protein concentrate has been reported to be used in fortification of starchy foods like maize, cassava and *akamu* flours [9].

Fermented dough (Sourdough) is an important modern fermentation of cereal flours and water based upon an earlier spontaneous process [10]. The sourdough microflora is dominated by lactic acid bacteria and, along with yeast; they play a key role in cereals fermentation. The sourdough fermentation has a number of beneficial effects that include prolonged shelf life, accelerated volume gain, delayed staling, improved flavour, and good nutritional value. Sourdough has been reported to contribute to extended shelf life by inhibiting the growth of spoilage bacteria and moulds [10]. With the increasing number of people especially, the low income earners suffering from malnutrition and many health challenges, demand for product high in nutritional and medicinal value is increasing. The use of African Yam Bean flour and cricket powder in the enrichment of rice-based *Masa*, will enhance the nutritional and health status of the consumers. The use of flour from broken rice in products could significantly increase the utilization of broken rice grains; thereby reducing losses that are normally incurred with the locally processed rice grains. Therefore, the objective of this research was to evaluate the nutritional quality of *Masa* produced from broken rice enriched with African yam bean flour and cricket blends.

2. MATERIALS AND METHODS

2.1 Material Procurement

Broken rice (*Oryza sativa*), African Yam Bean (*Sphenostylis stenocarpa*), and cricket (*Acheta domestica*), bakers yeast (*Sacharomyces cerevisiae*), sugar, salt, oil, trona (sodium sesquicarbonate) were purchased from modern market Kaura-Namoda, Zamfara State Nigeria and were taken to the Department of Food Technology, Federal Polytechnic Kaura-Namoda for further processing.

2.2 Preparation of African Yam Bean Flour

African yam bean flour was prepared according to the method described by [11]. African Yam Bean was weighed, sorted, washed and sundried after fermentation for 48 hours. The roasted bean was coarsely grinded, winnowed to remove the seed coats and grinded into flour using attrition mill and sifted through 250 µm aperture sieve. Fine flour was then obtained and placed in sealed containers for further use. The flow chart showing the production of African yam bean flour is shown Figure 1.

2.3 Preparation of Cricket Powder

The method described by Adamina *et al.* [12] was used for the preparation of cricket powder. The cricket was cleaned, boiled then fried and was dried. The dried cricket was milled and finally packaged for further processing. This is as shown in Figure 1.

Comment [DM4]: Will the additives used be expensive?

Comment [DM5]: Please provide the chemical formula.

Comment [DM6]: This is the first name, not the author's surname. There are also such errors in the References section.

Comment [DM7]: Adding process parameters would improve work reception and understanding of manufacturing technology.

Comment [DM8]: It should be Figure 2. Check the references throughout the content of the manuscript.

African Yam Bean Seeds




Figure 1: Flow Chart Showing the Production of African Yam Bean Flour

Source: Method described by [11] with modification

Comment [DM9]: All technological diagrams in the manuscript should be made more carefully.





Figure 2: Flow Chart Showing the Production of Cricket Powder

Source: Method described by [12] with modification.

2.4 Preparation of *Masa*

Masa was produced according to the modified method described [13]. The composite flour was mixed together following the addition of rice grits, water, yeast, sugar and salt. The mixture was then fermented for 6 hours followed by the neutralization of the fermented mixture with *trona*. The portion was remixed to have even distribution of the *trona* and was then cooked at 5 mins.

Comment [DM10]: Add a reference to the drawing as you did above to maintain the aesthetics of the work.






Figure 3: Flow Chart Showing the Production of *Masa*
Source: Method described by [13] with modification.

2.5 Blend Formulation for the Preparation of *Masa*

The criteria used for selecting the cereal-legume combination were based on Food and Agriculture Organization/ World Health Organization/UNU specifications of the daily protein and energy requirements for people of different ages, sexes, and levels of physical exertion.

Blends with different proportions of broken rice, African yam bean and cricket powder were prepared as shown in Table 1.

Table 1: Blend Formulation for the Production of *Masa* from Broken Rice, African Yam Bean and Cricket Powder

Sample Code	Ingredient (%)		
	Broken Rice Flour	African Yam Bean Flour	Cricket Powder
250	100	0	0
251	80	0	20
252	80	20	0
253	80	10	10

Table 2: Recipe Formulation for *Masa*

Comment [DM11]: I believe that this recipe should be moved to chapter 2.4.

Component	Masa Composition
Flour *	1.25 kg
Sugar	15 g
Water	500 ml
Yeast and Baking powder	5 g
Trona	10 ml
Oil	12 ml
Salt	Pinch

***Broken Rice or Composite Flour**

Source: Method described by [1] with modification

2.6 Determination of the Proximate Composition of Masa

The proximate composition of masa from broken Rice, african yam bean and cricket were determined according to the methods described by [14] and Carbohydrate content was determined by difference according to [15].

2.6.1 Moisture Content Determination

Moisture content was determined using the air oven dry method. A clean dish with a lid was dried in an oven (Uniscope Surgifriend Medicals, England) at 100 °C for 30 min. It was cooled in desiccators and weighed. Two (2) grams of sample was then weighed into the dish. The dish with its content was then put in the oven at 105 °C and dried to a fairly constant weight. The loss in weight from the original sample (before heating) was reported as percentage moisture.

$$\% \text{ Moisture} = \frac{\text{Weight Loss } (W_2 - W_3)}{\text{Weight of Sample } (W_2 - W_1)} \times 100$$

Where:

W_1 = Weight of dish,

W_2 = Weight of dish + sample before drying,

W_3 = Weight of dish + sample before drying.

2.6.2 Ash Content Determination

Two (2) gram of sample was weighed into an ashing dish which had been pre-heated, cooled in a desiccator and weighed soon after reaching room temperature. The crucible and content was then heated in a muffle furnace at 550 °C for 6 hrs. The dish was cooled in a desiccator and weighed soon after reaching room temperature. The total ash was calculated as percentage of the original sample weight.

$$\% \text{ Ash} = \frac{(W_3 - W_1)}{(W_2 - W_1)} \times 100$$

Where:

W_1 = weight of empty crucible, W_2 = weight of crucible + sample before ashing,

W_3 = weight of crucible + content after ashing

2.6.3 Crude Fibre Determination

Two (2) grams of the sample was extracted using Diethyl ether. This was digested and filtered through the California Buchner system. The resulting residue was dried at 130 °C for 2 hrs, cooled in a dessicator and weighed. The residue was then transferred into a muffle furnace (Uniscope Surgifriend Medicals, England) and ignited at 550 °C for 30 min, cooled and weighed. The percentage crude fibre content was calculated as:

Comment [DM12]: Need enter

$$\% \text{ Crude Fibre} = \frac{\text{Loss in weight after incineration}}{\text{Weight of original food}} \times 100$$

2.6.4 Crude Fat Determination

Fat was determined using Soxhlet method. Samples were weighed into a thimble and loose plug fat free cotton wool was fitted into the top of the thimble with its content inserted into the bottom extractor of the Soxhlet apparatus. Flat bottom flask (250 ml) of known weight containing 200 ml of hexane was fitted to the extractor. The apparatus was heated and fat extracted for 8 hrs. The solvent was recovered and the flask (containing oil and solvent mixture) was transferred into a hot air oven (Uniscopie Surgifriend Medicals, England) at 105 °C for 1 hr to remove the residual moisture and to evaporate the solvent. It was later transferred into desiccator to cool for 15 min before weighing. Percentage fat content was calculated as

$$\% \text{ Crude Fat} = \frac{\text{weight of extracted fat}}{\text{Weight of Sample}} \times 100$$

2.6.5 Crude Protein Determination

The Kjeldahl method was used to determine the percentage crude protein. Two (2) grams of sample was weighed into a Kjeldahl digestion flask using a digital weighing balance (Uniscopie Surgifriend Medicals, England: Max. 180 g). A catalyst mixture weighing 0.88 g (96 % anhydrous sodium sulphate, 3.5 % copper sulphate and 0.5 % selenium dioxide) was added. Concentrated sulphuric acid (7 ml) was added and swirled to mix content. The Kjeldahl flask was heated gently in an inclined position in the fume chamber until no particles of the sample was adhered to the side of flask. The solution was heated more strongly to make the liquid boil with intermittent shaking of the flask until clear solution was obtained. The solution was allowed to cool and diluted to 25 ml with distilled water in a volumetric flask. Ten (10) ml of diluted digest was transferred into a steam distillation apparatus. The digest was made alkaline with 8 ml of 40 % NaOH. To the receiving flask, 5 ml of 2 % boric acid solution was added and 3 drops of mixed indicator was dropped. The distillation apparatus was connected to the receiving flask with the delivery tube dipped into the 100ml conical flask and titrated with 0.01 HCl. A blank titration was done. The percentage nitrogen was calculated from the formula:

$$\% \text{ Nitrogen} = \frac{(S-B) \times 0.0014 \times 100 \times D}{\text{Sample Weight}}$$

Where S = sample titre, B = blank titre, $S - B$ = corrected titre, D = diluted factor

$\% \text{ Crude Protein} = \% \text{ Nitrogen} \times 6.25$ (correction factor)

2.6.6 Carbohydrate Determination

Carbohydrate content was determined by difference according to Ihekoronye and Ngoddy (1985) as follows:

$$\% \text{ Carbohydrate} = 100 - (\% \text{ Moisture} + \% \text{ Ash} + \% \text{ Fibre} + \% \text{ Fat} + \% \text{ Protein})$$

2.7 Determination of the Mineral Content (mg/100g) of Masa

The minerals contents were determined by the method described by [14].

2.7.2 Determination of Calcium

Calcium was determined using the atomic absorption spectrophotometer. Calcium carbonate (2.495 g) was dissolved and diluted to 100 ml with de-ionized water. This solution contains 1000 mg Ca²⁺ ions and from this stock solution, calcium standard of the following concentration levels 0.0, 3.0, 6.0, 9.0 were prepared. The absorbance of both the sample and the standard working aliquot were determined in the atomic absorption spectrophotometer (Uniscopie Surgifriends, England) at 239.9 nm. The concentration of the test mineral in the sample was calculated with reference to the graph (standard curve) and obtained as follows:

$$\text{Calcium} = \frac{100 \times Y \times Vf \times D}{W \times 100 \times Va}$$

Where

W = weight of the sample analyzed

Y = Concentration of Calcium obtained from the standard curve,

Vf = Total volume of extract

Va = volume of extract used

D = Dilution factor

Comment [DM13]: This description is too laconic and does not say anything (What is the method? What minerals?)

2.7.3 Determination of Phosphorus

Phosphorus was determined using spectrophotometer. Phosphorus in the sample was determined by the molybdate method using hydroquinone as a reducing agent. Sodium sulphate (1.0 ml), 1.0 ml of ammonium molybdate and 1 ml of hydroquinone were added to 1 ml of the sample digest. The mixture was agitated and allowed to stand for 30 minutes for the blue colour to develop. The absorbance of the sample was determined using the spectrophotometer at 600 nm. The phosphorus standard was prepared by dissolving 1.1 g of monobasic potassium phosphorus (KA_2PO_4) into a 500 ml volumetric flask containing 500 ml of distilled water. Five drops of toluene were added to diminish microbial activity. Twenty millilitre of the Standard stock was collected and made up to 100 ml. This contained 100 ppm. Standard stock (0.1 ml) = 0.2 ppm. Zero to one millilitre of the 100 ppm phosphorus stock solution was poured into 100 ml volumetric flask separately and treated the same way as the sample. The reading of the standard was taken at 600 nm in UV/VIS spectrophotometer (Uniscope Surgifriend Medicals, England) and a standard curve was plotted.

$$P = \frac{100 \times Au \times C \times Vf}{W \times As \times Va}$$

Where

W = Weight of sample analyzed

Au = Absorbance of test sample

As = Absorbance of standard phosphorus solution

C = Concentration (in mg/ml) of sample

Vf = Total volume of extract

Va = Volume of extract analyzed

2.7.4 Determination of Potassium

Potassium determination was by Flame Photometry. One (1) gram of sample was dissolved in 20 ml of acid mixture (650 ml of concentrated HNO_3 ; 80 ml PCA; 20 ml conc. H_2SO_4) and aliquots of the diluted clear digest were taken for photometry using Flame analyzer.

2.7.5 Determination of Iron

Standard solution containing 100 mg/ml of Fe^{3+} ions was prepared from 1 g pure iron wire. The wire was dissolved in 20 ml concentrated HNO_3 , boiled in water bath and diluted to 1000 ml with distilled water. Standard solutions with concentrations 0, 0.5, 1.0, 2.0 and 4.0 ppm was prepared. Two milliliter of sample aliquot was diluted to 100 ml and was used to determine the absorbance of the sample using an atomic absorption spectrophotometer (Uniscope Surgifriends Medicals, England) at 510 nm. The standard and samples absorbance were noted and concentration of iron in the sample was determined from the standard curve.

2.8 Determination of Anti-Nutritional Factors of *Masa*

2.8.1 Alkaloid Determination

The alkaloid content was determined gravimetrically by the method described by [16]. 5 g of each sample was weighed using a weighing balance and dispersed into 50 ml of 10 % acetic acid solution in ethanol. The mixture was well shaken and then allowed to stand for about 4 h before it is filtered. The filtrate was then evaporated to one quarter of its original volume on hot plate. Concentrated ammonium hydroxide was added drop wise in order to precipitate the alkaloids. A pre-weighed filter paper was used to filter off the precipitate and it was then washed with 1 % ammonium hydroxide solution. The filter paper containing the precipitate was dried on an oven at 60 °C for 30 min, transferred into desiccators to cool and then reweighed until a constant weight was obtained. The constant weight was recorded. The weight of the alkaloid was determined by weight difference of the filter paper and expressed as a percentage of the sample weight analyzed. The experiment was repeated thrice for each food stuff sample and the reading recorded as the average of three replicates.

2.8.2 Determination of Tannins

Tannin was determined using the Folin-Denis spectrophotometer method described by [16]. The sample (0.5 g) was weighed into a conical flask and 100 ml of distilled water was added into it. This was gently boiled for one hour and then filtered using Whatman filter paper into a 100 ml capacity volumetric flask. The filter paper was re-washed with distilled water and the filtrate was diluted to 100 ml mark and then cooled. Fifty millilitre aliquot was put into each flask for the development of greenish-blue colour. Five millilitre of Folin – Denis reagent (100 g sodium tungstate, 20 g phosphomolybdic acid, 50 ml of 85 % phosphoric acid and 750 ml of water) and 10 ml of saturated sodium carbonate solution was added into it.

This was diluted to 100 ml mark with distilled water after a thorough mixing. The flask was allowed to stand in a water bath at 25 °C for one-half hour and the absorbance was measured in the UV/VIS spectrophotometer (Uniscopce Surgifriend Medicals, England) at 760 nm. Distilled water was used as blank for the calibration curve. A standard curve was plotted and concentration of each sample was obtained and used for the tannin calculation.

$$\text{Tannin mg/g} = \frac{A_n \times C \times 100 \times V_f}{A_s \times W \times V_a}$$

Where

A_n = Absorbance of test sample

A_s = Absorbance of standard solution

C = Concentration of standard

W = Weight of sample used

V_f = Total volume of extract

V_a = Volume of extract analyzed

2.8.3 Determination of Oxalate

Oxalate was determined by the method described by [16]. This determination involved three major steps: digestion, oxalate precipitation and permanganate titration

Digestion

Two gram of sample flour was suspended in 190 ml of distilled water in a 250 ml volumetric flask. Ten millilitres of 6 ml HCL was added and the suspension was digested at 100 °C for 1 hour. It was cooled and made up to 250 ml before filtration.

Oxalate Precipitation

Duplicate portions of 125 ml of the filtrate were measured into beakers and four drops of methyl red indicator was added. This was followed by the addition of concentrated NH₄OH solution (drop wise) until the test solution changed from pink to a faint yellow colour. Each portion was then heated to 90 °C, cooled and filtered to remove precipitate containing ferrous ion. The filtrate was again heated to 90 °C and 10 ml of 5 % CaCl₂ solution was added while stirred constantly. After heating, it was cooled and left overnight at 5 °C. The solution was then centrifuged at 2500 rpm for 5 minutes. The supernatant was decanted and the precipitate was completely dissolved in 10 ml of 20 % (v/v) H₂SO₄ solution and then filtered for titration.

Permanganate Titration

The filtrate was made up to 300 ml. Aliquot of 125 ml from the filtrate was heated to near boiling and then titrated against 0.05 M standardized KMnO₄ solution to a faint pink colour which persisted for 30 seconds. The calcium oxalate content was calculated using the formula below.

$$\text{Oxalate (mg/g)} = \frac{T \times (V_{me})(D_f) \times 10^5}{M_E \times M_F}$$

Where

T = Titre of KMnO₄

V_{me} = Volume mass equivalent

D_f = Dilution factor

M_E = Molar equivalent of KMnO₄

M_F = Mass of sample used

2.8.4 Determination of Phytate

Phytate was determined using the method described by [16]. The sample was first extracted with 0.2 HCl. One millilitre of the extract was poured into a test tube fitted with a ground glass stopper together with 1 ml of ferric solution. The ferric solution was prepared by dissolving 0.2 g ammonium iron (III) sulphate in 10 ml of 2 NHCl. The solution was then made up to 100 ml with distilled water. The tube was heated in a boiling water bath for 30 minutes, cooled in ice for 15 minutes and then allowed to reach ambient temperature. The content of the tube was centrifuged for 30 minutes (300 rpm). After centrifugation the supernatant (1 ml) was mixed with 1.5 ml of 2,2 bipyridine solution and the absorbance measured at 519 nm against distilled water using UV/VIS spectrophotometer (Uniscopce Surgifriend Medicals, England). Thus, the phytic acid content is calculated as shown below.

$$\% \text{ Phytic acid} = \frac{100 \times V_f \times C}{W \times V_a \times 100}$$

Where

Comment [DM14]: Please enter a full name.

C = Concentration of curve ms/mole
V_a = Total volume of extract analysed
V_f = Total volume of extract
W = Weight of sample

2.9 Sensory Evaluation of the Masa

Sensory evaluation of the *Masa* samples was carried out according to the method described by [17]. A panel of twenty (20) members consisting of students and members of staff from Food Science and Technology Department, University of Agriculture Makurdi, Nigeria. Panelists were chosen based on their familiarity and experience with *Masa* for sensory evaluation. *Masa* produced from each flour blend, along with the reference sample were presented in coded form (A-C) and were randomly presented to the panelists. The panelists were provided with portable water to rinse their mouth between evaluations. However, a questionnaire describing the quality attributes (colour, aroma, taste, texture and overall acceptability) of the *Masa* samples was given to each panelist. Each sensory attribute was rated on a 9-point hedonic scale (1 = dislike extremely and 9 = like extremely). *Masa* was produced from broken rice flour (100 %) as control.

2.10 Statistical Analysis

The GENSTAT Statistical Software (version 17.0) was used for data analyses. Data were subjected to analysis of variance (ANOVA) and the separation of means was done using Fisher's Least Significant Difference (LSD) at (P<0.05).

3. RESULTS AND DISCUSSIONS

Table 3: Proximate Composition (%) of Masa

Samples	Moisture	Protein	Crude fat	Ash	Fibre	Carbohydrate	Energy (kcal/100g)
250	38.46 ^d ±0.11	5.60 ^d ±0.07	0.71 ^d ±0.02	0.51 ^d ±0.03	0.61 ^b ±0.06	54.13 ^a ±0.16	245.25 ^d ±0.18
251	40.64 ^b ±0.42	14.46 ^b ±0.13	8.49 ^b ±0.04	1.49 ^c ±0.02	0.79 ^a ±0.01	34.14 ^c ±0.34	270.77 ^a ±1.55
252	39.77 ^c ±0.30	12.50 ^c ±0.08	7.59 ^c ±0.03	2.74 ^b ±0.01	0.79 ^a ±0.06	36.61 ^b ±0.28	264.75 ^b ±1.21
253	44.48 ^a ±0.22	15.32 ^a ±0.09	9.71 ^a ±0.04	3.17 ^a ±0.08	0.62 ^b ±0.04	26.71 ^d ±0.28	255.47 ^c ±1.21
LSD	0.79	0.88	0.09	0.12	0.12	0.76	3.21

Values are mean± standard deviation of triplicate determination. Values with different superscript within the same column are significantly different at (P<0.05).

Key: 250= 100 % broken rice flour, 00 % African yam bean and 00 % cricket flour
 251= 80 % broken rice flour, 00 % African yam bean and 20 % cricket flour
 252= 80 % broken rice flour, 20 % African yam bean and 00 % cricket flour
 253= 100 % broken rice flour, 10 % African yam bean and 10 % cricket flour

3.1 Proximate Composition of Masa

The proximate composition showed that the moisture content ranged from 38.46-44.48 %. The 253 sample had the highest moisture content (44.48 %) while sample 250 having the lowest (38.46 %). The protein contents of the *Masa* samples ranged from 5.60 % in the control sample to 15.32 % in the 253 sample. The highest protein content recorded in the 253 sample was due to the protein contents of cricket and African yam bean [18]. The results are similar with the findings of [19] and [1]. Fat acts as lubricating agent which improves the quality of food in terms of texture and flavour. Also, fat provides energy and is essential as it carries along fat soluble vitamins A, D, E and K [19]. The 253 *Masa* sample recorded the

Comment [DM15]: Please provide the values according to the principle of significant numbers (3 significant places).

Comment [DM16]: The name of the subsection should be placed above the table with the test results.

Comment [DM17]: Please indicate in the text which table the description relates to.

highest fat content. This was followed by the 251 sample then the 252 and the lowest was recorded in the control. The highest value recorded in the 253 sample was as result of fat contents of cricket and African yam bean as reported by [20] and [21]. The ash contents of the *Masa* samples are significantly ($P<0.05$) different from each other. Ash content gives an insight to the mineral content of the food hence, the 252 and 253 samples can be described as good sources of minerals hence the relatively high-value of minerals recorded in them. The results obtained in this study are in agreement with similar reports by [19] and [22] for rice based *Masa* enriched with grain amaranth/carrot powder and pearl millet/cowpea/groundnut respectively. The highest crude fibre content of *Masa* products was recorded in the 251 and 252 samples and showed no significant ($p>0.05$) different in their values. Fibre consumption has been linked to decreased incidence of heart disease, various types of cancer and diverticulosis [23]. Also, high levels of fibre in foods help in digestion of foods and contribute to the health of the gastrointestinal tract and system in man by aiding normal bowel movement thereby reducing constipation problems which can lead to colon cancer [24]. The high fibre contents of the 251 and 252 samples suggest that they would be ideal food for people suffering from obesity, diabetes, cancer and gastrointestinal disorders [25]. According to Schneeman [24], the crude fibre contributes to the health of the gastrointestinal system and metabolic system in man. Olamide *et al.* [19] and Nkama and Nagappa [22] reported close range of fibre contents in *Masa*. Carbohydrate provides heat and energy for all forms of body activities and as such its inadequacy can cause the body to divert proteins and body fat to produce needed energy and this might lead to depletion of body tissues [26]. There was a significant difference ($p<0.05$) in the carbohydrate contents of the *Masa* samples. Supplementation of the broken rice with African yam bean and cricket significantly decreased the carbohydrate content. The highest content (54.13 %) was recorded in the control sample. This was followed by the 252 sample, then the 251 sample and the lowest was recorded in the 253 *Masa* sample. The observed decreased value was a result of high and low carbohydrate contents of broken rice and African yam and cricket respectively [27]. The energy content of the 251 *Masa* sample (270.77 kcal/100g) was significantly ($p<0.05$) higher than the energy contents of other samples. The least was recorded in the control sample (245.25 kcal/100g). The energy content of the various *Masa* products is high when compared to that of cassava products (flour and garri) (3.1-3.9 Kcal/100g) [28], which is considered as one of the main dietary energy source in Nigeria. The basis for the highest energy content of the 251 *Masa* sample could be attributed to the values of protein, fat and carbohydrate contents of the 251 sample.

Table 4: Vitamins and Mineral Composition of *Masa*

Samples	B-carotene (µg/100g)	Vitamin C (mg/100g)	Potassium (mg/100g)	Calcium (mg/100g)	Phosphorus (mg/100g)	Iron (mg/100g)
250	0.34 ^d ±0.01	0.52 ^d ±0.08	24.93 ^d ±0.88	143.03 ^d ±0.86	64.89 ^d ±0.77	22.06 ^d ±0.09
251	4.06 ^c ±0.14	4.39 ^c ±0.09	28.89 ^c ±0.09	174.19 ^c ±0.79	98.92 ^c ±0.58	35.46 ^c ±0.21
252	12.33 ^a ±0.06	10.61 ^a ±0.08	33.11 ^b ±0.08	194.42 ^b ±1.55	122.99 ^b ±0.61	42.62 ^b ±0.06
253	9.50 ^b ±0.09	9.98 ^b ±0.04	37.87 ^a ±0.04	284.11 ^a ±1.41	127.71 ^a ±0.81	50.67 ^a ±0.79
LSD	0.25	0.22	2.21	3.33	1.95	1.15

Values are mean± standard deviation of triplicate determination. Values with different superscript within the same column are significantly different at ($P<0.05$).

Key: 250= 100 % broken rice flour, 00 % African yam bean and 00 % cricket flour
 251= 80 % broken rice flour, 00 % African yam bean and 20 % cricket flour
 252= 80 % broken rice flour, 20 % African yam bean and 00 % cricket flour
 253= 100 % broken rice flour, 10 % African yam bean and 10 % cricket flour

Comment [DM18]: Please provide the values according to the principle of significant numbers (3 significant places).

3.2 Vitamins and Mineral Composition of Masa

The result of the vitamins and mineral contents of Masa are presented above; Vitamins are organic substances required only in small amounts in the body for metabolism. Their requirements are in milligrams (mg), micrograms (μg), milliequivalents (mEq) and international units (IU) (Chidinma *et al.*, 2010). The 252 Masa sample (12.33 $\mu\text{g}/100\text{g}$) was significantly ($p<0.05$) higher than other samples. This was followed by the 253 sample (9.50 $\mu\text{g}/100\text{g}$) then the 251 sample (4.06 $\mu\text{g}/100\text{g}$) and the lowest content was recorded in the 250 Masa sample (0.34 $\mu\text{g}/100\text{g}$). The highest value recorded in the 252 sample was as a result of high β -carotene value of African yam bean [21], [18]. Also, the sample 252 recorded the highest vitamin C content among the samples due to its high value of Vitamin C. Abioye *et al.* [21] revealed significant presence of vitamin C in African yam bean. The values of the vitamin C recorded are significantly different ($p<0.05$).

Minerals are inorganic elements which are essential for the normal functioning of the body. They are required in smaller quantities in addition to proteins, carbohydrates, fats and vitamins, they are inorganic or "ash constituents" of foods which cannot be destroyed by heating [29]. Although they yield no energy, they have important roles to play in many activities in the body [15]. As ash content gives an insight to the mineral content of the food, hence, Masa produced from composite flour can be described as a rich source of minerals as seen from the significant ($P<0.05$) increased in the mineral contents of Masa with substitution levels of African yam bean and cricket.

Potassium activates several enzyme reactions and helps in the release of energy from carbohydrates, fats and proteins. It also functions with sodium and calcium to regulate neuromuscular excitability [29]. The potassium content of the Masa samples ranged from 24.93-37.87 mg/100g with sample 253 and 250 having the highest and lowest content respectively. Calcium is a mineral required by the body for a variety of physiological functions and the maintenance of bone tissues throughout life [30]. Calcium is necessary for supporting bone formation and growth; it also helps in the maintenance of healthy teeth, skeletal and soft tissue, mucous membranes and skin. The result of the calcium content of Masa samples showed that, the 253 sample was significantly ($p<0.05$) higher than all other Masa samples. Phosphorus works closely with calcium to build strong bones and teeth [29]. The highest phosphorus content of the Masa samples was found in the 253 Masa sample (127.71 mg/100g) and the lowest was found in the control sample (64.89 mg/100g). The iron content of the Masa products ranged from 22.06-50.67 mg/100g with the control and sample 253 having the lowest and highest iron values respectively. The results of the vitamins and mineral contents of Masa are higher than those reported by Olamide *et al.* [19] for rice-based Masa.

Table 5: Anti-nutrients Contents (mg/100g) of Masa

Samples	Alkaloids	Tannins	Phytate	Oxalate
250	0.095 ^c ±0.05	0.535 ^d ±0.02	0.040 ^c ±0.01	0.015 ^c ±0.01
251	0.145 ^c ±0.01	0.615 ^c ±0.02	0.060 ^c ±0.00	0.030 ^c ±0.00
252	1.860 ^a ±0.01	2.600 ^a ±0.04	1.550 ^a ±0.01	0.285 ^a ±0.01
253	0.345 ^b ±0.01	1.640 ^b ±0.01	0.270 ^b ±0.06	0.140 ^b ±0.01
LSD	0.09	0.09	0.09	0.02

Values are mean± standard deviation of triplicate determination. Values with different superscript within the same column are significantly different at ($P<0.05$).

Key: 250= 100 % broken rice flour, 00 % African yam bean and 00 % cricket flour

251= 80 % broken rice flour, 00 % African yam bean and 20 % cricket flour

Comment [DM19]: The name of the subsection should be placed above the table with the test results.

Comment [DM20]: Please indicate in the text which table the description relates to.

Comment [DM21]: Please provide the values according to the principle of significant numbers (3 significant places).

252= 80 % broken rice flour, 20 % African yam bean and 00 % cricket flour

253= 100 % broken rice flour, 10 % African yam bean and 10 % cricket flour

3.3 Antinutritional Composition of *Masa*

The anti-nutritional contents of the *Masa* samples showed low levels of alkaloids, tannins, Phytate and oxalate. Sample 252 recorded the highest contents of antinutritional parameters due to the high content of anti-nutrients in African yam bean as presented in Table 5. Alkaloids have been reported to cause gastrointestinal and neurological disorder [31]. The alkaloid content of the 252 sample was significantly ($p < 0.05$) the highest compared to other samples. This was due to the level of African yam bean substitution in the composite *Masa* and the presence of alkaloid in African yam bean [31]. Tannins are known for their ability to precipitate with iron and other metals, thereby reducing their absorption [32]. Tannin is an anti-nutrient that inhibits activity of digestive enzymes [33]. The lower values obtained (0.535-2.600 mg/100g) for tannin is very important because tannic acid above 10 % of total dry weight affects overall nutritional potential of food material. Importantly, tannin can be used in treatment of skin eruption due to their astringent properties [33].

Phytate has strong binding capacity and forms insoluble complexes with multivalent cations, including Ca, Mg, Fe and Zn, and render them biologically unavailable [33]. Phytate ranged from 0.040 mg/100g in the control *Masa* to 1.550 mg/100g in the 252 *Masa* sample.

Oxalates affect the metabolism of magnesium and calcium. It also reacts with proteins to form complexes which have an inhibitory effect in the digestion of peptic. High oxalate diet can increase the risk of renal calcium absorption and has been implicated as a source of kidney stone. Generally, small amounts of oxalate may occur in many vegetables and fruits but do not pose nutritional problems [33]. The levels of these anti-nutrients in all the samples were relatively low, below toxic levels and may not hinder the bioavailability of essential nutrients in the composite *Masas*. Also, the levels observed here are not in considerable levels of inhibitors that may inhibit the absorption of minerals [32]. The results are similar to the findings of [35] but lower than the ones discovered in foods by [36].

Comment [DM22]: The name of the subsection should be placed above the table with the test results.

Comment [DM23]: On what basis was the content of anti-nutritional substances classified, were they compared to any consumption standards? Are these values safe?

Comment [DM24]: Please indicate in the text which table the description relates to.

Samples	Colour	Aroma	Taste	Texture	General Acceptability
250	6.13 ^c	6.67 ^a	7.20 ^{ab}	6.73 ^a	7.73 ^a
251	7.33 ^{ab}	7.07 ^a	6.27 ^c	6.33 ^a	6.67 ^b
252	7.73 ^a	6.33 ^a	7.33 ^a	5.93 ^a	7.73 ^a
253	6.80 ^{bc}	6.60 ^a	6.40 ^{bc}	6.13 ^a	6.67 ^b
LSD	0.68	1.08	0.86	1.07	0.77

Table 6: Sensory Attributes of *Masa*

Values with different superscript within the same column are significantly different at ($P < 0.05$)

Key: 250= 100 % broken rice flour, 00 % African yam bean and 00 % cricket flour

251= 80 % broken rice flour, 00 % African yam bean and 20 % cricket flour

252= 80 % broken rice flour, 20 % African yam bean and 00 % cricket flour

253= 100 % broken rice flour, 10 % African yam bean and 10 % cricket flour

3.4 Sensory Attributes of Masa

Sensory evaluation is usually carried out towards the end of product development or formulation cycle and this is done to assess the reactions of consumers/judges about the product to determine the acceptability of such product. It is an important criterion for assessing quality in the development of new products and for meeting consumer requirements [37]. Colour is an important sensory attribute of any food because of its influence on acceptability. Sensory attributes of various Masa Products are presented in Table 6. The result of the colour of the various Masa samples differ significantly ($p < 0.05$). Sample 252 (7.73) was the highest and the lowest was observed in the control (6.13) Masa sample.

Taste is a sensory parameter that affects the quality and acceptability of food products. No matter how rich or nutritious a food is, if it tastes bad, such food would not be accepted by people. Taste is the primary factor determining the acceptability of any product and has the highest impact in determining the market success of product. Sample 252 (7.33) taste better than other samples and the lowest was recorded in sample 251 (6.33).

General acceptability was determined on the basis of quality scores obtained from evaluation of colour, aroma, taste and texture. There was no significant ($p > 0.05$) different in the Masa samples of 250 and 252 as they recorded same general acceptability. Also, between samples 251 and 253, there was not significant different ($p > 0.05$) in their general acceptability. This results are similar with the findings of [19], [1] and [2].

4. CONCLUSION

The study showed that addition of African yam bean and cricket to broken rice in Masa production greatly increased the protein, fat, ash and fibre contents of Masa. Also, β -carotene, Vitamin C, potassium, calcium, phosphorus and iron contents of broken rice-based Masa increased with African yam bean and cricket addition. The ant-nutrients contents of the Masa products showed significant values at 20 % level of African yam bean substitution. Sensory analysis carried out in this study showed that Masa produced from 100 % broken rice and 20 % African yam bean substitution were generally accepted more than other Masa samples. Though they were most preferred, but the 20 % cricket addition and 10 % African yam bean and 10 % cricket substitution levels were also accepted.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Ayo, J. A., Agu H. and Famoriyo, O. F. Effect of Different Cereals on the Quality of Masa. *Pakistan Journal of Nutrition* 2008; 7 (4): 530-533.
2. Ayo, J. A., Onuoha, G., Agu, H., Ayo, V. A., Avu, E. O, Sosanya, M. and Adeosun, F. Effect of added beneseed paste on the quality of millet-based masa. *African Journal of Food Science and Technology*, 2012; 3(10). 236-243.
3. Dashen, M. M., Edia-Asuke, U.A., Amapu, T. Y., Jidangkat, M. G. and Hassan, B. B. Effect of Fermented Rice Dough on the Organoleptic Quality and Shelflife of Rice "Masa", *UJMR*, 2017; 2(1).
4. Okpala, L. C. and Egwu, P. N. Utilisation of broken rice and cocoyam flour blends in the production of biscuits. *Nigerian Food Journal*, 2015; 33:8-11.
5. Megido, R. C., Alabi, T., Nieuw, C., Blecker, Ch., Danthine, S., Bogaert, J. and Francis F. Optimisation of a cheap and residential small-scale production of edible crickets with local by-

Comment [DM25]: The name of the subsection should be placed above the table with the test results.

Comment [DM26]: Please indicate in the text which table the description relates to.

Comment [DM27]: It comes out 120%, which variant did the authors want?

Comment [DM28]: Please submit a utilitarian application and indicate which type of Masa made is the best according to the authors' research.

- products as an alternative protein-rich human food source in Ratanakiri Province, Cambodia. *J. Sci. Food Agric.*, 2015; 96, 627–632.
6. Ayieko, M. A., Ogola, H. J. and Ayieko, I. A. Introducing rearing crickets (gryllids) at household levels: Adoption, processing and nutritional values. *J. Ins. Food Feed*, 2016; 2(3), 203–211.
 7. Van- Huis, A. Potential of insects as food and feed in assuring food security. *Annu. Rev. Entomol.* 2013; 58, 563–583.
 8. United State Department of Agriculture (USDA). National Genetic Resources Program. *Germplasm Resources Information Network - (GRIN)*. National Germplasm Resources Laboratory, Beltsville, Maryland. URL:2010
 9. Eneche, H. E. (2005). Enrichment of starchy flours with African yam bean Protein Concentrate. *Nigerian Journal of Nutrition and Science*, 2005; 26(2): p. 30-37.
 10. Chavan, R. S. and Chavan, S. R. Sourdough Technology. A Traditional Way for Wholesome Foods: A Review. *Comprehensive Reviews in Food Science and Food Safety* 2011; 10: 78 – 79.
 11. Kukwa, R. E. Okpainya, P. E. and Ikya, J. K. Micronutrients in African Yam Bean-carrot Flours and Acceptability of Its Gruels for Complementary Food. *Asian Food Science Journal*, 2018; 4(2): 1-9
 12. Adamina D., Julia, A., Paulina, C., Justyna, J. and Przemyslaw, K. Quality and Nutritional/Textural Properties of Durum Wheat Pasta Enriched with Cricket Powder. *Foods*, 2019; 8, 46: 1-10.
 13. Nkama, I and Nagappa, G. M. Production and Nutritional Quality of Traditional Nigerian Masa from Mixtures of Rice, Pearl Millet, Cowpea, and Groundnut. *Food and Nutrition Bulletin*, 2015; 19(4):366-373.
 14. AOAC. Official Methods of Analysis of AOAC International. 19th edition. AOAC International, Gaithersburg, Maryland, USA.2012
 15. Ihekoronye, A. I. and Ngoddy, P. O. Integrated Food Science and Technology for the Tropics. London: Macmillan Publ. Ltd. 1985; 265.
 16. Onwuka, G.I. Food Analysis and Instrumentation theory and Practice: Analytical Techniques. 2nd Ed. Surulere., Naphthali prints, 2018; 229-453.
 17. Retapol T. N. and Hooker H. Consumer evaluation and preference heterogeneity for a novel functional food. *Journal of Food Science*, 2006; 71: 41-46.
 18. Lina, V. H. Potential Source of Protein for Human Consumption. Faculty of Natural Resources and Agricultural Sciences, Swedish University of Agricultural Sciences, 2018; 1-33.
 19. Olamide, A. A., Olusola, S. J., Ayo, F. and Blessing, I. Nutritional, Antioxidant and Sensory Properties of Rice-Based Masa Enriched with Grain Amaranth and Carrot Powder. *Annals of Food Science and Technology*, 2018; 19(4), 645-656.
 20. Wang, S., Wakasa, Y., Kawakatsu, T. and Takaiwa, F. Increased lysine content in rice grains by ver-accumulation of BiP in the endosperm. *Bioscience, Biotechnology and Biochemistry* 2010; 74(12): 2529 – 2531.
 21. Abioye, V. F., Olanipekun, B. F. and Omotosho, O.T. Effect of Varieties on the Proximate, Nutritional and Anti-nutritional Composition of Nine Variants of African Yam Bean Seeds (*Sphenostylis Stenocarpa*). *Donnis Journal of Food Science and Technology*, 2015; 1(2) pp. 017-021.
 22. Nkama, I and Nagappa, G. M. Production and Nutritional Quality of Traditional Nigerian Masa from Mixtures of Rice, Pearl Millet, Cowpea, and Groundnut. *Food and Nutrition Bulletin*, 2015; 19(4):366-373.
 23. Wildman, B. E. C. and Medeiros, D. M. Advanced Human Nutrition. Boca Raton, USA: CRC Press. 1999; 61–155.
 24. Schneeman, B. Gastrointestinal Physiology and Functions. *British Journal of Nutrition*, 2002; 88 (2): 159-163.
 25. Ufot, E. I. and Inemesit, E A. Physico-chemical and Sensory Qualities of Functional Bread Produced from Wholemeal Wheat and Unripe Plantain Composite Flours. *Malaysian Orthopaedic Journal of Food Processing and Technology*, 2016; 2(2), 48–53.
 26. Shirika, D., Igyor, M. A. and Gernah, D. I. Nutritional Evaluation of Complementary Food Formulations from Maize, Soybean and Peanut Fortified with *Moringa oleifera* Leaf Powder. *Journal of Food and Nutrition Sciences*, 2015; 6, 494-500.

Comment [DM29]: These are the names, not the names of the authors. Please check the cited items carefully.

27. Carolyne, K., John, N. K., Samuel, I. and Nanna, R. Use of House Cricket to Address Food Security in Kenya: "Nutrient and Chitin Composition of Farmed Crickets as Influenced by Age" *Academic Journals*, 2017; 12(44):3189-3197.
28. Onwueme, I. C. The tropical tuber crops. John Wiley and Sons Ltd, 1978; 118-159.
29. Chidinma, W. A., Jiddari, W. U. and Hassan, S. C. A Student Handbook on Food and Nutrition. De-New Creation Prints Ltd Publishers, Kaduna, Nigeria, 2010; 50-58.
30. Grosvernor, M. B. and Smolin, L. A. Nutrition: From Science to Life. Harcourt College Publishers, New York, USA, 2002; 288-371.
31. Aletor, V. A. Allelochemicals in plant food and feeding stuffs. Part I. Nutritiona biochemical and physiopathological aspects in Animal production. *J. Vet. Hum. Toxicol.* 1993; 34(1): 57-67.
32. Onuoha, C. H., Harry, B. J. and Eze, S. O. Evaluation of Nutrients and Anti-nutritional Factors of Different Species of African Yam Bean (*Sphenostylis stenocarpa*). *European Journal of Basic and Applied Sciences*, 2017; 4(1):1-8.
33. Adesuyi, A. O. and Ipinmoroti, K. O. The Nutritional and Functional Properties of the Seed Flours of three Varieties of *Carica Papaya*. *Current Research in Chemistry* 2011; (3):70-75.
34. Oyeleke, G. O., Isola, A. D., Salam, M. A. and Ajao, F. D. Evaluation of Some Chemical Composition of Pawpaw (*Carica Papaya*) Seeds under Normal Storage Ripening. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 2013; 4, (6), 18-21.
35. Ohizua, E. R., Abiodun, A. A., Micheal, A. I., Olajide, P. S., Adeniyi, T. A., Raphael, O. I., Simeon, O. A., Tolulope, O. O. and Ayorinde, F. Nutrient Composition, Functional, and Pasting Properties of Unripe Cooking Banana, Pigeon Pea, and Sweet Potato Flour Blends. *Journal of Food Science and Nutrition*, 2016; 5:750–762.
36. Ross, P. H. and Martha, K. Estimation of the Oxalate Content of Foods and Daily Oxalate Intake. *Journal of Kidney International*, 2000; Vol. 57, pp. 1662–1667.
37. Dogo, S. B., Doudjo, S., Mohamed, A.Y., and Ernest, K. K. Physico-Chemical, Functional and Sensory Properties of Composite Bread prepared from Wheat and defatted Cashew (*Anacardium occidentale L.*) Kernel Flour. *International Journal of Environmental and Agriculture Research*, 2018; 4(4), Page 88.

Comment [DM30]: These are the names, not the names of the authors. Please check the cited items carefully.