

Original Research Article

G x E interaction studies in single cross maize hybrids for quality traits under different environments

Abstract Sixty eight genotypes of maize (48 hybrids, 16 parents and 4 checks) were evaluated under three different environments to test their stability. Variance due to genotypes, environments and genotype x environment interactions were found significant for oil content, starch content and protein content. Regression approach by Eberhart and Russel 1966 used to investigate the nature of G x E interaction and for identifying genotypes possessing general and specific adaptation. None of the hybrids had stability for all quality traits. For Grain Starch content hybrid EI-1155-1 x EI-2416, for oil content hybrid EI-670-2 x EI-2403 and for Protein content hybrids EI-2187 x EI-2403, EI-536-3 x EI-2403, EI-2176 x EI-561-2, EI-2173 x EI-561-7 had higher mean values, non significant deviation from regression and regression coefficient around unity ($b_i=1$) indicating the stable performance in different environment and wider adaptability.

Keywords: maize; grain quality; protein; oil; starch content, stress, linear regression, regression coefficient

Introduction: Maize (*Zea mays* L.) along with wheat and rice provides at least 30 percent of the food calories to more than 4.5 billion people in 94 developing countries where one-third of children are malnourished. By 2050, the demand for maize in the developing world will be almost double to the current demand. However, an estimated 15 to 20 per cent of yield losses each year reported due to drought and climate changes. Maize is affected by drought at different growth stages in different regions. Germination potential, seedling growth, seedling stand establishment, overall growth and development, pollen development, silk development, anthesis-silking interval, pollination, embryo development, endosperm development and kernel development are the events in the life of maize crop which are seriously hampered by drought stress. (Aslam *et al*, 2015).

The use of genetics to improve drought tolerance and provide yield stability is an important part of the solution to stabilizing global production. That is why the development of maize varieties with enhanced tolerance to drought stress and higher water use efficiency (WUE) has become a high priority goal for

Comment [U1]: Use keywords other than those mentioned in the job title

Comment [U2]: The introduction can be complemented by the insertion of more gaps that the work will fill. Use more current references

Comment [U3]: Carefully check the formatting of the text, there are places with errors

major breeding programs, both in the private and public sectors.(Maazou *et al.*2016). Regression approach is being increasingly used to investigate the nature of G x E interaction and for identifying genotypes possessing general and specific adaption in almost all the crop plants. in present study, we use Eberhart and Russell (1966) model to identify hybrids suitable for different environment for- the protein, oil, and starch compounds of grains.

Materials and Methods:

The experimental material comprised of 12 inbred lines *viz.*, EI-2176, EI-2145, EI-670-2, EI-1155-1, EI-2173, EI-1280-1, EI-2187, EI-2174, EI-586-03, EI-2172, EI-536-3 and EI-2173-56 and 4 testers EI-2403, EI-2127, EI-561-2, EI-2416 their 48 F₁s and 4 checks *viz.*, HM-5, CHM-08-287, Vivek maize hybrid-21 and PMH-3. These 48 F₁s were obtained by crossing 12 inbred with 4 testers in line x tester mating design during late *kharif* 2016.

total 68 genotypes (12 inbred lines, 4 testers, 48 crosses and 4 checks) were sown in a randomized block design with three replications in three different environments during spring 2018 at Plant Breeding and Genetics farm, Rajasthan College of Agriculture, Udaipur (Rajasthan). Each treatment was sown in single row plot of 4 meter length maintaining crop geometry of 60 x 25 cm row to row and plant to plant spacing, respectively. Each environment was separated by 2 m channel for proper water management. Non-experimental rows were planted around the layout to eliminate border effects. The other recommended agronomical practices were used to raise a healthy crop. Starch content of the seeds was estimated by using Anthrone reagent method. Oil content of the seeds was estimated by using Soxhlet's ether extraction method (A. O. A. C., 1965) and Protein content of the seeds was estimated by using Lowry's method. The details of the three environments are presented in Table 1.

Table 1. Details of the environments

| | |
|---------------|--|
| Environment 1 | Normal irrigation |
| Environment 2 | Irrigation withheld at anthesis for 20 days |
| Environment 3 | Irrigation withheld at grain filling stage for 20 days |

Results and Discussion: The pooled analysis of variance (Table 2) revealed significant differences among the hybrids for all the traits thus indicated the existence of -genetic variability. Mean square due to

Comment [U4]: Talk about the statistic to be applied, and you can separate the section into subsections

Comment [U5]: The results and discussion can be improved, since there is much narration of the results and little discussion about what the work really contributed to the medium.

environment, genotype and genotype x environment, were significant for all characters, indicating that all the hybrids interacted considerably well with the environmental conditions. Mean square due to genotypes was significant for all traits under study. Analysis of variance by Eberhart and Russel, 1966 (table 3) revealed that Mean square due to environment (E) plus genotypes x environment (G x E) interaction also found significant for all characters. Mean square due to environment (linear) was significant for grain protein content. The mean square due to G x E linear interaction was significant for all characters. The significant mean square due to pooled deviation for all characters was significant indicated that the genotype differed considerably with respect to their stability.

The stability parameters, such as regression coefficient (b_i) and deviation from regression (S^2_{di}) along with mean performance of genotypes for character under study were computed to assess the stability and suitability of performance over the environment (table 4) and results are narrated as following.

For grain starch content Out of 68 ~~genotype~~ genotypes, 47 genotype exhibited non-significant deviation from regression (S^2_{di}), indicating their predictable behavior for grain starch content. Among parental lines, tester EI-2127 inbred lines EI-1280-1, EI-2187 and EI-EI-2174 showed regression coefficient around unity ($b_i = 1$), whereas tester EI-2416 and inbred lines EI-2172 and EI-2173-56 had above unity ($b_i > 1$) regression coefficient stable performance in different environments as normal as well as water stressed conditions for higher grain starch content, especially.

Non-significant with higher mean value from regression (S^2_{di}) was observed in 36 hybrids. Out of which, hybrids EI-2176 x EI-2403, EI-2145 x EI-2127, EI-2173 x EI-2403, EI-EI-2174 x EI-2403, EI-586-03 x EI-2403, EI than population mean which indicates that inbred line have deviation -2176 x EI-2127, EI-670-2 x EI-2127, EI-1280-1 x EI-2127, EI-2187 x EI-2127, EI-1280-1 x EI-561-2, EI-2187 x EI-561-2, EI-EI-2174 x EI-561-2, EI-2173-56 x EI-561-2, EI-2176 x EI-2416, EI-670-2 x EI-2416, EI-1155-1 x EI-2416, EI-2173 x EI-2416, EI-2187 x EI-2416 and EI-2172 x EI-2416 showed regression coefficient around unity ($b_i = 1$) with higher mean values than population mean indicating the stable performance in different environment for higher grain starch content. Four hybrids viz. EI-2172 x EI-2403, EI-2145 x EI-2127, EI-1155-1 x EI-2127 and EI-536-3 x EI-2127 depicted regression coefficient less than

unity ($b_i < 1$) along with higher mean value than population mean showed hybrids superiority and stability water stressed environments for higher grain starch content. The two hybrids *viz.*, EI-2176 x EI-561-2 and EI-2145 x EI-561-2 exhibited regression coefficient above unity ($b_i > 1$) along with higher mean value than population mean showed hybrids superiority and stability under favorable environments for higher starch content.

Grain oil content:

Out of 68 genotype, twelve genotype showed non-significant deviation from regression (S^2_{di}) indicating their predictable behavior for grain oil content.

Tester EI-2416 and inbred line EI-2173, showed regression coefficient around unity ($b_i = 1$) with higher mean value than population mean indicating stable performance under different environment. Inbred line EI-2176 showed regression coefficient above unity ($b_i > 1$) with higher mean value than population mean indicating stable performance of inbred line under favorable environment.

Non-significant deviation from regression (S^2_{di}) was observed in five hybrids. Out of which, hybrid *viz.*, EI-670-2 x EI-2403, showed regression coefficient around than unity ($b_i = 1$) with higher mean values than population mean showed hybrids superiority and stability under different environments for higher oil content. while hybrid EI-586-03 x EI-561-2 showed regression coefficient less than unity ($b_i < 1$) and one hybrid EI-2145 x EI-2416 exhibited regression coefficient above unity ($b_i > 1$) along with higher mean value than population mean showed hybrids stability under unfavorable and favorable environments for higher oil content.

Grain protein content

Non-significant deviation from regression (S^2_{di}) was observed in 3 parents but only one parent EI-586-03 exhibited higher mean value than population mean with regression coefficient around unity ($b_i = 1$) indicating stable performance under different environment.

Non-significant deviation from regression (S^2_{di}) was observed in fifteen hybrids. Out of them, four hybrids *viz.*, EI-2187 x EI-2403, EI-536-3 x EI-2403, EI-2176 x EI-561-2 and EI-2173 x EI-561-2, showed higher mean value than population mean regression coefficient around unity ($b_i = 1$) indicating stable performance under different environment. One hybrid EI-2145 x EI-2403 showed regression coefficient above unity ($b_i > 1$) with higher mean values than population

mean, While, hybrid EI-586-03 x EI-2416 exhibited regression coefficient less than unity ($b_i < 1$) with higher mean value than population mean, indicating hybrids performance were stable under favorable and unfavorable as moisture deficit environments for protein content, respectively. similar findings for starch content, oil content and protein content were reported by Dadeech and Joshi (2007), Shanthi *et al.* (2011) Baudh Bharti *et al.* (2017) and Patel and Kathiria (2017). Further classification of genotype as per their suitability in different environment is shown in table ~~no.~~ 4.

Table 2 Pooled analysis of variance grain starch content, grain oil content and grain protein content

| SN | Source | df | Grain starch content | Grain Oil content | Grain Protein content |
|----|--------------|-----|----------------------|-------------------|-----------------------|
| 1. | Environment | 2 | 1975.10** | 37.85** | 188.95** |
| 2. | Rep./Env | 6 | 1.03 | 0.01 | 0.01 |
| 3. | Genotype | 67 | 50.60** | 1.68** | 3.61** |
| 4. | G x E | 134 | 7.27** | 0.24** | 0.52** |
| 5. | Pooled Error | 402 | 0.81 | 0.01 | 0.02 |
| 5. | Bartlet | 2 | 2.33 | 1.94 | 0.47 |

*, ** Significant at 5% and 1% respectively

Table 3 Analysis of variance Eberhart and Russel (1966)

| SN | Characters | Genotype [67] | E+(G x E) [136] | E (L) [1] | G x E (L) [67] | Pool dev. [68] | Pool Err [402] |
|----|---------------------------|------------------|--------------------|--------------|-------------------|-------------------|-------------------|
| 1. | Grain starch content (%) | 16.87** | 12.07** | 0.28 | 23.45** | 1.03** | 0.27 |
| 2. | Grain oil content (%) | 0.56** | 0.27** | 0.01 | 0.46** | 0.08** | 0.00 |
| 3. | Grain protein content (%) | 1.20** | 1.10** | 0.03* | 1.97** | 0.26** | 0.01 |

[] Degrees of freedom

*, ** Significant at 5% and 1% respectively

Table 4 Stability parameters for grain starch content (%), grain oil content (%) and grain protein content (%)

| SN | Genotype | Grain starch content (%) | | | Grain oil content (%) | | | Grain protein content (%) | | |
|----|-----------------------|--------------------------|-------|----------|-----------------------|--------|----------|---------------------------|--------|----------|
| | | μ_i | b_i | S^2d_i | μ_i | b_i | S^2d_i | μ_i | b_i | S^2d_i |
| 1 | EI-2403 | 60.91 | 1.89 | 3.580** | 3.96 | 0.09 | 0.078** | 7.54 | 1.05 | 0.021* |
| 2 | EI-2127 | 61.70 | 1.18* | -0.173 | 3.18 | 1.64 | 0.203** | 8.12 | 1.02 | 0.284** |
| 3 | EI-561-2 | 56.85 | 1.13* | -0.236 | 4.08 | 0.38 | 0.007* | 9.27 | 1.03 | 0.031* |
| 4 | EI-2416 | 63.19 | 1.35 | 0.659 | 3.74 | 0.41 | 0.002 | 7.83 | 1.01 | 0.578** |
| 5 | EI-2176 | 58.65 | 1.13* | -0.236 | 3.73 | 1.32 | 0.021** | 8.40 | 1.03 | 0.394** |
| 6 | EI-2145 | 56.12 | 1.13* | -0.236 | 3.92 | 0.63 | 0.069** | 8.55 | 0.76 | 1.230** |
| 7 | EI-670-2 | 54.24 | 0.53 | 1.201* | 3.70 | 0.89 | 0.001 | 9.52 | 0.95 | 0.224** |
| 8 | EI-1155-1 | 64.57 | 0.67 | 0.352 | 3.17 | 1.38 | 0.033** | 7.67 | 0.90* | -0.000 |
| 9 | EI-2173 | 59.14 | 1.13* | -0.236 | 3.97 | 0.77* | -0.001 | 8.08 | 1.06 | 0.112** |
| 10 | EI-1280-01 | 60.00 | 1.13* | -0.236 | 3.73 | 0.97* | -0.000 | 7.63 | 1.16 | 0.082** |
| 11 | EI-2187 | 59.73 | 1.11 | -0.006 | 4.28 | 0.42 | 0.044** | 8.64 | 1.03 | 0.074** |
| 12 | EI-2174 | 61.47 | 0.87 | 0.556 | 3.17 | 0.40 | 0.002 | 7.27 | 1.04 | 0.138** |
| 13 | EI-586-03 | 57.13 | 2.00 | 4.642** | 3.85 | 1.05 | 0.041** | 8.54 | 1.03 | 0.014 |
| 14 | EI-2172 | 60.84 | 1.95 | 4.095** | 3.26 | 0.84 | 0.034** | 7.48 | 1.05** | -0.006 |
| 15 | EI-536-3 | 58.45 | 1.13* | -0.236 | 4.08 | 1.18 | 0.040** | 8.86 | 1.01 | 0.305** |
| 16 | EI-2173-56 | 59.91 | 2.06 | 5.221** | 4.06 | 0.32*+ | -0.002 | 7.75 | 1.04* | 0.000 |
| 17 | EI-2176 x EI-2403 | 63.85 | 1.13* | -0.236 | 4.11 | 1.30 | 0.044** | 8.39 | 0.94 | 0.662** |
| 18 | EI-2145 x EI-2403 | 61.31 | 1.13* | -0.236 | 4.02 | 1.37 | 0.012** | 8.41 | 0.45 | 0.746** |
| 19 | EI-670-2 x EI-2403 | 57.66 | 0.09 | 2.570** | 4.80 | 1.26 | 0.004 | 9.27 | 0.68 | 0.376** |
| 20 | EI-1155-1 x EI-2403 | 61.95 | 0.47 | 1.525** | 3.95 | 0.61 | 0.007* | 7.80 | 1.05 | 0.049** |
| 21 | EI-2173 x EI-2403 | 62.02 | 1.13* | -0.236 | 4.21 | 1.02 | 0.010** | 7.90 | 1.23* | -0.005 |
| 22 | EI-1280-01 x EI-2403 | 62.92 | 0.36 | 2.271** | 3.74 | 1.08 | 0.023** | 7.95 | 1.01* | -0.003 |
| 23 | EI-2187 x EI-2403 | 60.99 | 0.09 | 4.699** | 4.11 | 1.04 | 0.122** | 8.55 | 1.05* | -0.000 |
| 24 | EI-2174 x EI-2403 | 62.14 | 1.13* | -0.236 | 4.03 | 1.73 | 0.064** | 8.03 | 1.01 | 0.339** |
| 25 | EI-586-03 x EI-2403 | 62.45 | 1.13* | -0.236 | 3.88 | 1.01 | 0.065** | 7.72 | 1.04* | 0.001 |
| 26 | EI-2172 x EI-2403 | 63.74 | 0.63 | 0.650 | 3.34 | 1.71 | 0.228** | 7.85 | 1.01 | 0.434** |
| 27 | EI-536-3 x EI-2403 | 57.99 | 1.13* | -0.236 | 3.94 | 1.84 | 0.085** | 8.53 | 1.04* | -0.005 |
| 28 | EI-2173-56 x EI-2403 | 58.88 | 1.13* | -0.236 | 4.82 | 0.97 | 0.055** | 8.50 | 1.01 | 0.478** |
| 29 | EI-2176 x EI-2127 | 59.75 | 1.13* | -0.236 | 4.31 | 1.36 | 0.054** | 8.10 | 1.05* | 0.003 |
| 30 | EI-2145 x EI-2127 | 62.07 | 0.31 | 2.692** | 3.99 | 1.63 | 0.034** | 8.86 | 1.98*+ | -0.001 |
| 31 | EI-670-2 x EI-2127 | 61.73 | 1.13* | -0.236 | 3.86 | 0.91 | 0.023** | 8.12 | 1.02 | 0.174** |
| 32 | EI-1155-1 x EI-2127 | 64.35 | 0.63 | 0.650 | 3.66 | 1.38* | -0.001 | 7.69 | 1.02 | 0.224** |
| 33 | EI-2173 x EI-2127 | 59.57 | 1.13* | -0.236 | 4.45 | 1.02 | 0.402** | 8.94 | 1.05 | 0.047** |
| 34 | EI-1280-01 x EI-2127 | 61.05 | 1.13* | -0.236 | 3.67 | 1.48 | 0.038** | 8.52 | 1.01 | 0.545** |
| 35 | EI-2187 x EI-2127 | 62.55 | 1.13* | -0.236 | 4.44 | 1.20 | 0.294** | 7.61 | 1.02 | 0.250** |
| 36 | EI-2174 x EI-2127 | 62.98 | 0.25 | 3.145** | 3.78 | 1.00 | 0.060** | 7.80 | 1.03 | 0.062** |
| 37 | EI-586-03 x EI-2127 | 57.23 | 1.13* | -0.236 | 3.53 | 1.09 | 0.248** | 9.34 | 1.01 | 0.529** |
| 38 | EI-2172 x EI-2127 | 62.18 | 0.25 | 3.145** | 4.67 | 0.56 | 0.263** | 7.77 | 1.06 | 0.198** |
| 39 | EI-536-3 x EI-2127 | 62.18 | 0.69 | 0.422 | 4.28 | 0.62 | 0.015** | 7.74 | 1.06 | 0.196** |
| 40 | EI-2173-56 x EI-2127 | 62.69 | 1.78 | 2.648** | 4.34 | 1.42 | 0.034** | 6.92 | 1.04* | -0.005 |
| 41 | EI-2176 x EI-561-2 | 59.80 | 1.24 | -0.078 | 4.27 | 0.33 | 0.137** | 8.78 | 1.00 | 0.008 |
| 42 | EI-2145 x EI-561-2 | 61.95 | 1.40 | 0.401 | 4.58 | 1.02 | 0.082** | 8.44 | 1.58 | 0.044** |
| 43 | EI-670-2 x EI-561-2 | 58.98 | 1.13* | -0.236 | 3.61 | 0.50 | 0.027** | 10.13 | 1.02 | 1.316** |
| 44 | EI-1155-1 x EI-561-2 | 61.42 | 1.13* | -0.236 | 4.85 | 0.60 | 0.020** | 7.61 | 1.04 | 0.007 |
| 45 | EI-2173 x EI-561-2 | 58.19 | 1.40 | 0.401 | 4.34 | 1.32 | 0.085** | 8.50 | 1.04* | -0.003 |
| 46 | EI-1280-01 x EI-561-2 | 62.77 | 1.13* | -0.236 | 4.41 | 0.63 | 0.161** | 7.64 | 1.03 | 0.092** |
| 47 | EI-2187 x EI-561-2 | 60.47 | 1.13* | -0.236 | 4.74 | 1.26 | 0.083** | 7.70 | 1.09 | 0.985** |
| 48 | EI-2174 x EI-561-2 | 62.40 | 1.13* | -0.236 | 3.77 | -0.16 | 0.055** | 7.46 | 1.05 | 0.050** |
| 49 | EI-586-03 x EI-561-2 | 58.90 | 1.13* | -0.236 | 3.82 | 0.70 | 0.004 | 8.36 | 1.06 | 0.206** |
| 50 | EI-2172 x EI-561-2 | 62.90 | 0.42 | 1.882** | 3.69 | 0.95 | 0.005 | 8.28 | 1.21 | 0.027* |
| 51 | EI-536-3 x EI-561-2 | 63.98 | 0.58 | 0.909* | 4.77 | 1.58 | 0.300** | 7.73 | 1.01 | 0.445** |
| 52 | EI-2173-56 x EI-561-2 | 63.60 | 1.13* | -0.236 | 4.44 | 1.02 | 0.125** | 7.64 | 1.04** | -0.006 |
| 53 | EI-2176 x EI-2416 | 65.81 | 1.13* | -0.236 | 3.91 | 2.09 | 0.386** | 7.28 | 0.84 | 0.257** |
| 54 | EI-2145 x EI-2416 | 58.80 | 1.13* | -0.236 | 3.90 | 1.57* | 0.000 | 8.75 | 0.59 | 1.297** |
| 55 | EI-670-2 x EI-2416 | 62.05 | 1.13* | -0.236 | 4.25 | 0.24 | 0.005* | 7.84 | 1.01 | 0.375** |
| 56 | EI-1155-1 x EI-2416 | 66.33 | 0.93 | 0.249 | 4.01 | 0.90 | 0.096** | 7.12 | 1.01 | 0.456** |
| 57 | EI-2173 x EI-2416 | 60.92 | 1.13* | -0.236 | 4.31 | 1.29 | 0.141** | 7.95 | 1.00 | 0.794** |

| SN | Genotype | Grain starch content (%) | | | Grain oil content (%) | | | Grain protein content (%) | | |
|----|-----------------------|--------------------------|-------|----------|-----------------------|-------|----------|---------------------------|-------|----------|
| | | μ_i | b_i | S^2d_i | μ_i | b_i | S^2d_i | μ_i | b_i | S^2d_i |
| 58 | EI-1280-01 x EI-2416 | 61.65 | -0.24 | 8.677** | 3.17 | 1.03 | 0.587** | 8.69 | 1.03 | 0.039** |
| 59 | EI-2187 x EI-2416 | 60.71 | 1.13* | -0.236 | 3.96 | 1.61 | 0.129** | 8.28 | 1.02 | 0.299** |
| 60 | EI-2174 x EI-2416 | 58.54 | 1.13* | -0.236 | 4.02 | 1.14 | 0.037** | 8.16 | 0.58 | 0.010 |
| 61 | EI-586-03 x EI-2416 | 65.02 | 1.13* | -0.236 | 3.44 | 0.86 | 0.028** | 8.22 | 0.53 | 0.001 |
| 62 | EI-2172 x EI-2416 | 61.66 | 1.13* | -0.236 | 3.81 | 0.15 | 0.016** | 8.12 | 0.69* | -0.001 |
| 63 | EI-536-3 x EI-2416 | 57.45 | 1.13* | -0.236 | 4.66 | 1.31 | 0.013** | 9.71 | 1.03 | 0.669** |
| 64 | EI-2173-56 x EI-2416 | 62.71 | 0.36 | 2.271** | 4.81 | 1.40 | 0.012** | 7.62 | 1.08 | 0.733** |
| 65 | HM-5 | 62.09 | 1.12* | -0.263 | 3.80 | 0.80 | 0.021** | 8.26 | 1.05 | 0.023* |
| 66 | CHM-08-287 | 61.31 | 1.08 | 0.492 | 4.53 | 1.35* | -0.000 | 8.27 | 1.03 | 0.038** |
| 67 | Vivek Maize Hybrid 21 | 61.51 | 0.38 | 0.459 | 4.49 | 0.92 | 0.034** | 7.06 | 1.04* | -0.001 |
| 68 | Pratap Hybrid Maize-3 | 63.07 | 0.70 | -0.204 | 4.20 | 0.27 | 0.037** | 7.98 | 0.27 | 0.003 |

*, ** Significantly deviating from zero at 5% and 1% respectively
+, ++ Significantly deviating from unity at 5% and 1% respectively

Table 5. Classification of genotype suitable for various environments

| s.no. | Characters | Different environments (bi=1) | Unfavorable environments (bi<1) | Favorable environments (bi >1) |
|-------|---------------------------|---|---|---|
| 1 | Grain Starch content (%) | EI-1155-1 x EI-2416 | EI-2172 x EI-2403, EI-1155-1 x EI-2127 and EI-536-3 x EI-2127 | EI-2176 x EI-2403, EI-2145 x EI-2403, EI-2173 x EI-2403, EI-2174 x EI-2403, EI-586-03 x EI-2403, EI-670-2 x EI-2127, EI-1280-01 x EI-2127, EI-2187 x EI-2127, EI-2145 x EI-561-2, EI-1280-01 x EI-561-2, EI-2187 x EI-561-2, EI-2174 x EI-561-2, EI-586-03 x EI-561-2, EI-2173-56 x EI-561-2, EI-2176 x EI-2416, EI-2145 x EI-2416, EI-670-2 x EI-2416, EI-2187 x EI-2416, EI-586-03 x EI-2416, EI-586-03 x EI-2416 and EI-2172 x EI-2416 |
| 2 | Grain Oil content (%) | EI-670-2 x EI-2403 | EI-586-03 x EI-561-2 | EI-2145 x EI-2416 |
| 3 | Grain Protein content (%) | EI-2187 x EI-2403, EI-536-3 x EI-2403, EI-2176 x EI-561-2, EI-2173 x EI-561-7 | EI-586-03 x EI-2416 | EI-2145 x EI-2403 |

Comment [U6]: Standardize table

Formatted: Font: Bold

Formatted: Font: Bold

Conclusions????

Formatted: Font: (Default) Times New Roman, 12 pt

References:

Comment [U7]: Standardize references!

- Aslam, M., Maqbool, M. A. and Cengiz, R. 2015. Effects of drought on maize. Springer Briefs
- A.O.A.C. 1965. Official methods for oil analysis for Association of Official Agricultural Chemists 10th Ed Washington, D.C.
- Baudh Bharti, Dubey, R. B., Kumar A., Dadheech, A. and Dhobi R. K., 2017. Stability analysis for grain yield and quality parameters in QPM (*Zea mays* L.) inbred line crosses. *Int.J.Curr.Microbiol.App.Sci.* 6(6): 3177-3185
- Clegg K. M.,1956. The application of the anthrone reagent to the estimation of starch in cereals. *J. Sci. Food Agric.*, 7,40-44.
- Dadheech, A. and Joshi, V. N. 2007. Stability analysis for quality traits in maize (*Zea mays* L.) *Agric. Sci. Digest*, 27 (1): 26 – 29.
- Eberhart, S.A. and Russell, W.A. 1966. Stability parameters for comparing varieties. *Crop Sci.*, 6:36-40.
- Lowry, O.H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. 1951. *Journal of Biological Chemistry*, 193: 265 (The original method).
- Maazou, A. R. S., Tu, J. L., Qiu, J. and Liu, Z. Z. 2016. Breeding for drought tolerance in maize (*Zea mays* L.). *American J. of Plant Sci*, 7, 1858- 1870.
- Patel, P. and Kathiria, K. B. 2017. Stability analysis for quality traits in maize (*Zea mays* L.). *Electronic Journal of Plant Breeding*, 7 (3): 634-639.
- Shanthi, P., Satyanarayana, E., Suresh babu, G., and Sai Kumar R. 2011. Studies on exploitation of heterosis for grain yield and quality parameters in quality protein maize (QPM) genotypes (*Zea mays* L.). *Crop Res.*, 41(1, 2 &3): 171-175.