

Monitoring Some Pesticide Residues in Imported Date palm fruits in United Arab Emirates during the Year 2020

Abstract

The occurrence of pesticide residues in representative samples collected from imported dates palm fruits during 2020 to United Arab Emirates (UAE) were investigated to ensure compliance with the standard specifications and requirements by the regulatory and supervisory authorities, maintain health and safety of consumers and improve food safety. An accurate, rapid and reliable method for the simultaneous determination of pesticide multi-residues in 230 samples imported dates by liquid chromatography coupled with tandem mass spectrometry LC-ESI (+)-MS/MS operating in multiple reaction monitoring (MRM) mode and modified quick, easy, cheap, effective, rugged, safe (QuEChERS) method was used. The performance of the analytical method was validated in accordance with EU SANCO guidelines (SANTE/12682/2019) for monitoring pesticide multi-residues to check compliance with existing regulations especially for European Community. Residues level of 343 compounds, were determined in 230 samples. Results indicated that the percentage of samples with residues above the maximum residue levels (MRL) were 4.34% in dates samples. whereas samples with residues within MRL were 7.39% in dates samples. A total of 230 samples 88.26% were free from detectable residues. Out of the 343 pesticides tested, 11 pesticides were found above the limit of detection, according to UAE, Codex and European regulations. **Conclusions**

Keywords: Determination, Dates, Multi residue, pesticides, liquid chromatography, Tandem mass spectrometry (MS/MS), QuEChERS, UAE.

1. Introduction

Dates are the fruit of the date palm tree (*Phoenix dactylifera* L.), which is grown in many tropical regions of the world. The importance of the date palm for human nutrition mainly due to its health benefits of dates such as lowers cholesterol, protein rich, rich in vitamins, improves bone health, strengthens the nervous system, rich in iron, promotes digestion, improves skin, fixes hangovers and assists in weight gain [1]. Dates is considered as principal food in Arab peninsula.

Pesticide residues in food commodities has become a major cause of concern all-over the world. Food safety is crucial and consumers have to be assured that they are not exposed to an unacceptable level of pesticide residues. After the establishment of the world trade order (WTO) and other GATT, SPS, presence of the residues above the permissible level is also a major bottleneck in the international trade of food commodities.

In order to ensure food safety for consumers and protect human health, many organizations and countries around the world have established maximum residue limits (MRLs) for pesticides in food commodities [2, 3, 4]. The MRL is the maximum level of a pesticide residue (expressed in $\text{mg}\cdot\text{kg}^{-1}$) which is legally permitted in or on food or animal feed [3].

Therefore, the pesticide residues in food have been strictly regulated by governments all over the world to determine whether the concentration of pesticides used exceeds their MRLs

At the Ministry of Climate Change and Environment - UAE we are keen to ensure that all foodstuffs and products in the country, both domestically produced and imported, are safe for consumption and to strengthen infrastructure at Quarantine stations to prevent entry of food commodities which have pesticide residues above maximum residue limit (MRL), Testing / Certification of pesticide residue in export / import consignments [4,5].

The maximum permitted levels of pesticide residues in foods are stipulated by regulatory bodies in the UAE. National Laboratories Department, Ministry of climate change and environment had worked Monitoring of Pesticide Residues in imported food commodities through department laboratories which accredited by the British Commission for Accreditation (UKAS) [6].

Consequently, the aim of the present study is to ensure that pesticide residues not exceed the maximum permissible limit of pesticide residues in imported dates to the country, which is considered safe limit for the consumer approved by the local and international specialized agencies

2. Materials and Methods

Chemicals and standard solutions

Certified analytical standards tested pesticides were purchased from Dr Ehrenstorfer GmbH (Germany), with purity between 92.0 and 99.5%, Acetonitrile (Merck, Germany), methanol (LC-MS grade, Scharlab), Formic acid (Honeywell, Germany).

Ready-made QuEChERS kits were purchased from Supelco; Supel™ QuE citrate extraction tube (contains 4.0 g MgSO₄, 1.0 g NaCl, 0.5 g Na Citrate dibasic sesquihydrate, 1.0 g Na Citrate tribasic dehydrate), Supel™ QuE PSA SPE clean up Tube (contains 150.0 mg Supelclean PSA, 900.0 mg MgSO₄.) The solutions were prepared with Ultrapure demineralized water Milli-Q plus system (Merck-Millipore Corporations, USA).

Standard preparation: Individual analytical stock solutions (1000 mg L⁻¹) of each pesticide were prepared in methanol, considering the purity of each pesticide standard. These analytical solutions were diluted in methanol to 100 mg L⁻¹. All solutions were stored in amber flasks at – 18 °C. Afterwards, a mixture with the concentration of 10 mg L⁻¹ containing all pesticides was prepared, that was diluted to 1 mg L⁻¹ and was kept in refrigerator at 4.0 ± 2.0°C (Table 1). Internal standard solution: A solution of Triphenylphosphate (TPP), analytical grade, was used as an internal standard

Apparatus: Analysis of final extracts Agilent 6460 triple quadrupole, on a reversed-phase column and detected by tandem mass spectrometry (MS/MS) using electrospray ionization (ESI). The pesticide residues determined with positive ESI only. The total chromatographic run time was 32 minutes. Injection volume was 2.0 µL and the column temperature was set at 60⁰ C. The Agilent Mass Hunter Workstation software B.04.00 Features was used for data analysis. All pesticides were detected in the multiple reaction monitoring modes (MRM). Each pesticide has

precursor ion there were two product ions determined. One product ion used for quantification and other one was used for qualification, detected pesticides are in Table 1.

Sampling

The pesticide residue random monitoring program aimed to collect samples from all imported agricultural consignments at UAE border ports according to plans based on scientific foundations with the aim of laboratory analysis to assess the extent to contain pesticide residues of pesticides in accordance with the Technical Regulations UAE issued by Emirates Authority for Standardization and Metrology No. «UAE.S MRL 1-2019» regarding maximum limits [7]. Total of 230 date samples were collected from January to December 2020 from imported consignments in all over UAE Emirate's ports.

Samples (230 in total) were taken by the Ministry's inspectors; each sample approximately 1 Kg were placed in sterile polythene bags, in an ice chest box, to avoid contamination, deterioration, labelled, and transferred to the laboratory under appropriate transport conditions within 24 hours for analysis.

The used method was developed at the National Laboratories Department, Ministry of Climate Change and Environment (MOCCA), which is accredited by UKAS (The United Kingdom Accreditation Service) according to ISO 17025:2017 (International Organization for Standardization, 2017) and accreditation was gradually extended, for the analysis of pesticides in several foodstuffs and compliance with the document SANTE/12682/2019[8] “Guidance document on analytical quality control and method validation procedures for pesticide residue analysis in food and feed” which was issued by the European Commission Directorate General for Health and Food Safety, and became effective on January 1, 2020. Samples were labeled and crushed after seeds were discarded; then stored at 4°C until analysis. Blank samples (Pesticide free samples) were acquired from the consumer market used for validation experiments.

Sample preparation

The pesticide residues were analyzed using the method approved by the European Union to determine the level of Pesticides in food products QuEChERS method reported by Anastassiades *et al.*, 2003 [9] by using LCMSMS (liquid chromatography tandem mass spectrometry)

This method is validated to use in National Laboratories Department in the Ministry of Climate Change and Environment in UAE. It employs dispersive solid phase extraction (SPE) followed by chromatographic analysis of the extracts. Liquid Chromatography coupled with tandem Mass Spectrometry was adopted to achieve qualitative and quantitative analysis.

The following sample extraction and clean up steps were conducted: A homogenate date sample (5 g) in a 50-mL PTFE centrifuge tube. was extracted with 10 mL of water, shake then, 10 ml acetonitrile add and shaken for 1 min. Following this step, a mixture of extraction salts (4 g of magnesium sulfate; 1 g of sodium chloride; 1 g of trisodium citrate dihydrate and 0.5 g of disodium hydrogen citrate sesquihydrate) were added to each sample. After shaken vigorously for 1 min, tubes were centrifuged for 10 min at 4000 rpm to separate solid from acetonitrile supernatant which was ready to dispersive Solid Phase Extraction with 150.0 mg Supelclean PSA and 900.0 mg MgSO₄. The final extract was stabilized by 5% formic acid in acetonitrile solution. The sample was then injected into a Liquid chromatograph coupled to tandem mass spectrometry (LC-MS/MS). Recovery compliance with the document SANTE/12682/2019 [8] representative portions of the previously homogenized samples were spiked homogeneously with the appropriate amount of the working standard solution in methanol (0.1 mg kg⁻¹).

Table 1: List of active ingredients tested for their residues

Abamectin	Carbaryl	Demeton S methyl sulfone	Etoazole	Fuberidazole	Metconazole	Phenmedipham	Quinoxifen	Tri-allate
Acenaphthene	Carbendazim	Demeton-S-methyl	Etrifos	Furalaxyl	Methabenzthiazuron	Phenthoate	Quizalofop-ethyl	Triazamate
Acephate	Carbetamide	Desmedipham	Famoxadone	Furathiocarb	Methidithion	Phosalone	Rotenone	Triazophos
Acetamiprid	Carbofuran	Diazinon	Fenamidone	Halfenprox	Methiocarb	Phosmet	Secbumeton	Trichlorfon
Acibenzolar S methyl	Carbofuran-3-Hydroxy	Dichlofluanid	Fenamiphos	Halofenozide	Methiocarb sulfone	Phosphamidon	Siduron	Tricyclazole
Acrinathrin	Carboxin	Dichlorvos	Fenamiphos sulfone	Heptenophos	Methomyl	Picolinafen	Silthiofam	Tridemorph
Alachlor	Chlorantranilprole	Diclobutrazol	Fenarimol	Hexaconazole	Methoprotryne	Picoxystrobin	Simazine	Trifloxystrobin
Alanycarb	Chlorfenapyr	Dicloran	Fenazaquin	Hexaflumuron	Methoxychlor, o,p'-	Piperonylbutoxide	Simetryn	Triflumizole
Aldicarb	Chlorfenvinphos	Dicrotofos (Dicrotophos)	Fenbuconazole	Hexazinone	Methoxychlor, p,p'-	Pirimicarb	Spinetoram	Triflumuron
Aldicarb sulfone	Chlorfluzauron	Dicrotophos	Fenhexamid	Hexythiazox	Methoxyfenozide	Pirimiphos-ethyl	Spinosad	Trifluralin
Aldicarb sulfoxide	Chloridazon	Diethofencarb	Fenitrothion	Hydramethylnon	Metobromuron	Pirimiphos-Methyl	Spiromesifen	Triticonazole
Ametryn	Chlorobenzilate	Difenoconazole	Fenoxycarb	Imazalil	Metribuzin	Prochloraz	Spirotetramat	Uniconazole
Aminocarb	Chlorothalonil	Diflubenzuron	Fenpiclonil	Imidacloprid	Mevinphos	Procymidone	Spiroxamine	Vamidothion
Amitraz	Chlorotoluron	Diflufenician	Fenpropathrin	Indoxacarb	Mexacarbate	Profenofos	Sulfentrazone	Vinclozolin
Atrazine	Chloroxuron	Dimethenamid	Fenpropimorph	Ipconazole	Molinate	Promecarb	Sulfotep	Zoxamide
Azaconazole	Chlorpropham	Dimethoate	Fenpyroximate	Iprobenfos	Monocrotophos	Prometon	Tebuconazole	
Azinphos ethyl	Chlorpyrifos	Dimethomorph	Fenthion	Iprodione	Monolinuron	Prometryne	Tebufenozide	
Azinphos methyl	Chlorpyrifos-methyl	Dimoxystrobin	Fenthion sulfoxide	Iprovalicarb	Monuron	Propachlor	Tebufenpyrad	
Azoxystrobin	Chlorthal-dimethyl	Diniconazole	Fenuron	Isocarbophos	Moxidectin	Propamocarb	Tebutam	
Beflubutamid	Clethodim	Dinotefuran	Fenvalerate	Isofenphos	Myclobutanil	Propanil	Tebuthiuron	
Benalaxyl	Clodinafop-propargylester	Dioxacarb	Fipronil	Isoprocab	Napropamide	Propaquizafop	Tecnazene	
Bendiocarb	Clofentezine	Dioxathion	Flamprop-methyl	Isoproturon	Neburon	Propargite	Teflubenzuron	
Benfuracarb	Clomazone	Diphenylamine	Flonicamid	Isxadifen-ethyl	Nitenpyram	Propazine	Tefluthrin, cis-	
Benomyl	Clothianidin	Diuron	Fluazifop-butyl	Ivermectin	Norflurazon	Propetamphos	Temephos	
Benzoximate	Coumaphos	DMST	Fluazifop-p-butyl	Kresoxim-methyl	Novaluron	Propham	Terbumeton	
Bifenazate	Cyanazine	Doramectin	Fluazinam	Lenacil	Nuarimol	Propiconazole	Terbutryne	
Bifenthrin	Cyazofamid	Emamectin	Flubendimide	Linuron	Ofurace	Propoxur	Tetrachlorvinphos	
Bitertanol	Cycloxydim	Endosulfan -beta isomer	Fludioxonil	Lufenuron	Omethoate	Propyzamide	Tetraconazole	
Boscalid	Cycluron	Endosulfan sulfate	Flufenacet	Malaoxon	Oxadiazon	Proquinazid	Tetradifon	
Bromacil	Cyfluthrin I	Endosulfan-alpha isomer	Flufenoxuron	Malathion	Oxadixyl	Prosulfocarb	Tetramethrin	
Bromopropylate	Cyhalothrin (lambda)	EPN	Fluometuron	Mandipropamid	Oxamyl	Prothiofos	Thiabendazole	
Bromoxynil	Cymoxanil	Epoxiconazole	Fluoxastrobin	Mecarbam	Oxyfluorfen	Pymetrozine	Thiacloprid	
Bromuconazole	Cypermethrin	EPTC	Fluquinconazole	Mefenacet	Paclbutrazol	Pyracarbolid	Thiamectoxam	
Bupirimate	Cyproconazole	Ethiofencarb	Flusilazole	Mefenpyr-diethyl	Paraoxon ethyl	Pyraclostrobin	Thidiazuron	
Buprofezin (Z-isomer)	Cyprodinil	Ethion	Flutolanil	Mepanipyrim	Paraoxon methyl	Pyraflufen-ethyl	Thiobencarb	
Butafenacil	Cyromazine	Ethiprole	Flutriafol	Mepronil	Parathion-methyl	Pyrazophos	Thiodicarb	
Butocarboxim	DDD-p,p'	Ethirimol	Folpet	Mesotrione	Penconazole	Pyridaben	Thiophanate-methyl	
Butoxycarboxim	DDE-p,p'	Ethofenprox	Foramsulfuron	Metaflumizone	Pencycuron	Pyrifenox	Tolclofos-methyl	
Buturon	DDT-o,p'	Ethofumesate	Forchlorfenuron	Metalaxyl	Pencyuron	Pyrimethanil	Tolyfluanid	
Cadusafos	DDT-p,p'	Ethoprophos	Formetanate	Metamitron	Pendimethalin	Pyriproxyfen	Triadimefon	
Cafentrazone ethyl	Deltamethrin	Ethoxyquin	Fosthiazate	Metazachlor	Permethrin	Pyrudaphenthion	Triadimenol	

3.Results and Discussion

To enhance food safety in the UAE and ensure that the highest rated food products are traded only in local markets by monitoring pesticide residues in imported foods. This monitoring present a statistically representative snapshot of the situation of pesticide residues in imported date fruits to UAE, which are the most widely consumed in many Arab Gulf countries and Arab countries. The main purpose of the data presented in this monitoring is to inform citizens who have an interest in food safety on the situation regarding pesticide residue in imported food. In particular, answering the following questions: What actions were taken by the national competent authorities responsible for food control to ensure that pesticide residues in food comply with UAE food standards? How frequently were pesticide residues found in date fruits? Which pesticides were found?In 2020, During this year's monitoring program, carried out to looking for a range of up to 343 pesticides in imported date palm fruits. This program surveyed 230 date samples

Monitoring the presence of pesticides residues

Using the LC chromatography coupled with tandem mass spectrometry LC-ESI (+)-MS/MS method, this study analyzed the presence of 343 pesticide residues (Table 1)using the multi residue methodin imported date to UAE during the year 2020 in 230 samples available from UAE entry points.After optimization of the analytical conditions, a total of 343 pesticides were detected with good sensitivity by the multi-residue analysis method. Following parameters were investigated according to the criteria for validation parameters described by the European Commission (SANTE/12682/2019), recovery, linearity and precision of peak areas. Good linearity was obtained with correlation coefficients of regression ($R^2 > 0.99$), the recovery results ranged from 70% to 120% and the precision based on the relative standard deviations (RSD) ranged from 2% to 20%, under replicated conditions; these values are considered acceptable according to the requirements of the European Commission (2019). No residues were detected in the blank date palm fruits samples at the LOD of method.

The European Commission regulation sets maximum levels of certain pollutants in food products. This regulation is often updated, and regardless of the specific limits of general

foodstuffs, there are a number of specific limits for pollutants for specific products, including dates. The most common requirements for date pollutants relate to pesticide residues[10].

Herein the results showed that 95.65% of the samples (220in total) were in conformity with the specifications, the percentage of samples free of pesticide residues was 88.26%, while the percentage of samples that contain pesticide residues exceeded the permissible limits (MRL)were4.34% (10 samples) of the total number of samples (Table 2) andconsidered as non-compliant with CODEX and EU legislation on pesticide residues MRLs [11]

Table (2):The percentage of samples that compatible and exceed the permissible limits during the year 2020

year/quarter	Free from pesticide residues limits	Within pesticide residues limits	Exceed pesticides residues limits
Quarter 1 2020	80.82	10.95	8.21
Quarter 2 2020	82.35	17.64	0.00
Quarter 3 2020	88.63	6.81	4.54
Quarter 4 2020	93.59	3.84	2.56
Total	88.26	7.39	4.34

It was also shown by following up on the implementation of the monitoring that the percentage of samples that exceeded the permissible limits decreased, as the percentage in the first quarter of the year was 8.21%, while in the fourth quarter of the year it amounted to 2.56%,

On the contrary, no MRL exceedance was reported in second quarter for pesticides that were found to exceed the legal limit in first quarter e. g. Carbendazim, Deltamethrin, Ethion, Fipronil, Diflubenzuron, Indoxacarb and Triflumuron.

In overall quantification rate recorded in fourth quarter,2020 was slightly lower than the one in first quarter, 2020 and this bringing the total percentage of samples exceeding the limits allowed in the program 4.34% (Table 2). For pesticides with MRL exceedances during2020 year a significant decrease of MRL exceedance rate was observed within fourth quarter 2020.

Nevertheless, continuous consumption of food products even with moderate pesticide contamination may have negative consequences on human health in the long term[12].

Several countries have relied on applications of the broad spectrum of synthetic pesticides, such as Amitraz, Fenpropythrane, Finperochemate, Finazakuen, Propargate, Ttradyphon, and

Hexithiasoxy[13]and many studies have detected **M**onitoring pesticide residues in dates marketed, in Saudi Arabia within 2018 [14] Pesticide residues were detected in 36 (18%)within the MRL allowed but 15 samples (7.5%) exceeded the maximum residue levels, in Tunisia monitor pesticide residues in dates sampled from Tunisian oasis during the agricultural season 2017-2018 and to check compliance with existing regulations especially for European Community[15].

MRLs are based on good agricultural practices and examined to ensure that acceptable daily intake or acute reference dose is not exceeded. In the light of these results, action has been taken on samples containing MRL overrun in accordance with approved standards requiring the importer to re-export or destroy product to maintaining food safety in imported date palm fruits

Residues measured above the MRL

Given the maximum residue limits (MRL) of pesticides in date palms, a total of 10 samples with residues exceeding the legal limit (above MRL) of one or several pesticides were detected as shown in table 3 during this monitoring program

Table (3): list of detected pesticides above the MRL

Sample no	Pesticide Detected	Residue Detected (mg/kg)	MRL* (mg/kg)	EU Pesticides Database**
1	Carbendazim	1.016	0.1	Reg. (EU) No 559/2011
	Spirodiclofen	0.13	0.02	Reg. (EU) 2016/1902
2	Deltamethrin	0.166	0.01	Reg. (EU) 2018/832
	Ethion	0.134	0.01	Reg. (EU) No 310/2011
	Fipronil	0.037	0.005	Reg. (EU) 2019/1792
4	Diflubenzuron	0.64	0.01	Reg. (EU) 2019/91
	Indoxacarb	0.01	0.02	Reg. (EU) 2015/845
5	Deltamethrin	0.04	0.01	Reg. (EU) 2018/832
6	Triflumuron	0.1	0.01	Reg. (EU) 2018/1516
7	Indoxicarb	0.29	0.02	Reg. (EU) 2015/845
8	Imidacloprid	0.477	0.05	Reg. (EU) No 491/2014
9	Acetamiprid	0.12	0.01	Reg. (EU) 2019/88
	Pyriproxyfen	0.14	0.05	Reg. (EU) 2020/856
10	Acetamiprid	0.489	0.01	Reg. (EU) 2019/88

*MRL maximum residue level

** EU Pesticides Database

In this study residues exceeding the legal limits were related to 11 different pesticides in date samples as shown in above table were Acetamiprid, Carbendazim, Deltamethrin, Diflubenzuron, Ethion, Fipronil, Imidacloprid, Indoxacarb, Pyriproxyfen, Spirodiclofen and

Triflumuron, accounted 47.8% (11/23) and out of 11 pesticides that exceeded the MRLs, banned pesticides were detected in 3 (27.27%) pesticides were banned for use in agriculture (Carbendazim, Ethion, Fipronil). The most frequently detected pesticides exceeded MRLs were Acetamiprid, Deltamethrin and Indoxacarb. The fact that 4.34% of samples exceeding the MRLs and contained 3 banned pesticides, suggested that banned pesticides might still be frequently used in date in exported country. These results reflect that a certain proportion of farmers may not follow Good Agricultural Practices (GAP).

Residues of more than one pesticide (Multiple residues)

In this study, some samples contained only one pesticide residue, but 40% (4 out of 10) of exceeded the permissible limits samples had multiple residues from more than one insecticide present in the same sample while 17.64% (3 out of 17) within MRL. More than one residue has been detected in date samples, and in recent years, indicating the mixed use of pesticides due to pest resistance, multiple pesticides have been widely applied.

Table (4): Distribution of pesticides residues above MRL in samples monitored

Pesticides	Maximum value detected (mg/kg)	Status of Pesticides	Pesticides	Maximum value detected (mg/kg)	Status of Pesticides	Pesticides	Maximum value detected (mg/kg)	Status of Pesticides
Deltamethrin		Allowed	Imidacloprid		Restricted	Carbendazim		Banned
Acetamiprid		Allowed	Spirodiclofen		Restricted	Ethion		Banned
Triflumuron		Allowed				Fipronil		Banned
Indoxacarb		Allowed						
Diflubenzuron		Allowed						
Pyriproxyfen		Allowed						

*According to List of Registered Pesticides in the Ministry (MOCCA) Last update 13 August 2021

Pesticide targets in many countries as well as the World Organization for the Prohibition of Pesticides more toxic to humans, as well as pesticides that remain for the longest period in the environment and protect public health by setting limits on pesticide residues in food. From a regulatory point of view, pesticides are among the most regulated chemical products in the UAE. All pesticides that have a significant threat to human health and the environment are banned in the UAE.

Of the pesticide residues detected above EU residue limits (MRLs), according to the Ministry's List of Pesticides (MOCCA), a total of 11 pesticides are incompatible with regulations

exceeding the legal limit, including 6 pesticides permitted in the UAE, two restricted pesticides and 3 unapproved substances (banned pesticides) (Table 4).

In UAE Carbendazim, Ethion and Fipronil are not allowed to be used, it banned compounds (Table 4) according to Ministerial Decree No. (36) of 2018 regarding banned and restricted pesticides in the United Arab Emirates.

Although WHO classified fipronil and Ethion by hazard as moderately hazardous (Class II) [16]. Fipronil has been associated with human Possible Human Carcinogen according to EPA classified.

Carbendazim is classified as pesticides unlikely to present acute hazard in normal use is also detected in some of the samples. Fipronil (0.037 mg kg^{-1}) to be detected above the upper limit of the maximum (MRLs level) allowed residue set by Reg. (EU) 2019/1792 (0.005 mg kg^{-1}) for date palm fruits, these results give an indication of possible misuses of non-approved active substances, should follow-up on these national monitoring program and take corrective measures where appropriate

4. Conclusion

Pesticide Residue Monitoring Program is a compliance program used to monitor the level of chemical residues of pesticides in imported palm fruits in the UAE during 2020 to ensure that they do not exceed MRL limits using liquid mass spectrometry (MS/MS) techniques because improper use of pesticides leads to professionalism, environmental and food security risks, and monitoring pesticide residues helps to assess the risks of consumer exposure to such residues and ensure a high level of consumer protection.

Although pesticides are used to control pests and diseases, it is therefore recommended that the pesticide residue control program should continue in imported dates in the UAE while maintaining the analytical scope as widely as possible, taking into account the fact that the residues of unappointed active substances in the 2020 control program are low. On the other hand, there is control of all pesticide residues in all imported food products, and the high observed rate of sample compliance with the regulation that does not exceed their MRL in palm fruit dates of good agricultural practices (GAP) follows

5.

6. COMPETING INTERESTS DISCLAIMER:

7.

8. Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

9.

10.

11.

12.

13.

14. References

[1] Vayalil P (2012) Date fruits (*Phoenix dactylifera* Linn): an emerging medicinal food. *Crit. Rev. Food Sci. Nutr.* 52, 249–271.

[2] United States Department of Agriculture Maximum Residue Limit Database. Foreign Agricultural Service Department. [(accessed on 18 December 2016)]; Available online: <http://www.fas.usda.gov/maximum-residue-limits-mrl-database..>

[3] Codex Alimentarius Commission Pesticide Residues in Food and Feed. Plant Production and Protection Division. [(accessed on 4 January 2017)]; Available online: <http://www.fao.org/fao-who-codex-alimentarius/standards/pestres/en>.

[4] Nasra M. Abd El- Mageed, Ideisan I. Abu- Abdoun, Abdulla S. Janaan (2020) Monitoring of Pesticide Residues in Imported Fruits and Vegetables in United Arab Emirates during 2019. *International Research Journal of Pure and Applied Chemistry*, Page 239-260. DOI: 10.9734/irjpac/2020/v21i2330322. Published: 31 December 2020.

[5] Nasra M Abd El-Mageed, Ideisan I Abu-Abdoun, Abdulla S Janaan (2020)

“Monitoring of Pesticide Residues in Imported Fruits and Vegetables in United Arab Emirates during 2019”. *Journal International Research Journal of Pure and Applied Chemistry* PP239-260 (2020).

[6] European Commission Health and Food Safety, Regulation. [(accessed on 10 October 2016)]; Available online: http://ec.europa.eu/food/plant/pesticides/max_residue_levels/eu_rules/index_en.htm.

- [7] Emirates Authority for Standards & Metrology (ESMA) UAE.S MRL1 :2019. Maximum Residue Limits (MRLs) for Pesticides in agricultural and food products.
- [8] Guidance document on analytical quality control and method validation procedures for pesticides residues analysis in food and feed ANTE/12682/2019
- [9] Anastassiades M, Lehotay SJ, Stajnbaher D, Schenck F (2003) Fast and easy multiresidues method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *J AOAC Int.* 86:412–431
- [10][Entering the European market for dates | CBI](#)
- [11] EU Pesticides Database (2020) Pesticide residues and maximum residue levels (mg/kg).
- [12]Chronic neurological sequelae of acute organophosphate pesticide poisoning. Savage EP, Keefe TJ, Mounce LM, Heaton RK, Lewis JA, Burcar PJ *Arch Environ Health.* 1988 Jan-Feb; 43(1):38-45
- [13] Fernando PC. Pesticides, environment, and food safety. *Food and Energy Security.* 2017; 6(2):48-60.
- [14] Osama I Abdallah, Saleh S Alamer, Abdulmajeed M Alrasheed (2018) Monitoring pesticide residues in dates marketed in Al-Qassim, Saudi Arabia using a QuEChERS methodology and liquid chromatography-tandem mass spectrometry. *Biomed Chromatogr*, 32(6): e4199.
- [15] Sabrine Attia, Sahar Zougari, HajerSahraoui, RafikaAloui, SabrineNsir, WiemHached, Ghada Nsir and KaoutharGrissaLebdi. Pesticide residues surveillance of date palm (*Phoenix dactylifera*) in the south of Tunisia *Journal of Entomology and Zoology Studies* 2019; 7(2): 1085-1088.
- [16]The WHO Recommended Classification of Pesticides by Hazard and Guidelines to Classification 2019.