

Original Research Article

INFLUENCE OF THE CONSUMPTION OF FOUR DIETS ON THE BIOLOGICAL BLOOD PARAMETERS OF JAPANESE QUAIL (*Coturnix japonica*) FARMED IN CÔTE D'IVOIRE.

ABSTRACT

To ensure that newly formulated food (A1, A2 and A3) based on local products and commercial love (AT), ingested to quail reared in captivity for 9 weeks, did not affect their health biological, biochemical blood analyzes were carried out on them at the end of the experiment. The results obtained show that the quails fed with A3 and AT foods have the highest respective crude protein levels (55.67 ± 1.70 and 55 ± 0.82 g/l). In addition, the blood Ca/P ratio which represents the index of a good metabolism of these electrolytes, between 1 and 2, is in accordance with the recommendations in force. In addition, the average urea and creatinine levels of all the quails studied are identical to those of the control quails. Also the highest uricemia in this study is that obtained in quails fed with commercial feed AT (66 ± 0.82 mg/l). Quails fed with commercial food AT have the highest cholesterol level (2.24 ± 0.04 g/l). However, quails fed with feed A3 are those with the lowest level of triglycerides (6.80 ± 0.02 g/l). Thus, consumption of the newly formulated feeds A1, A2 and A3 by quails did not affect their renal, hepatic, pancreatic and heart health like that of quails fed the commercial feed AT.

Keywords: Food, quail, biological blood parameters, health, organs.

INTRODUCTION

Coturniculture is a recent activity. It has attracted the attention of specialists as a new way of diversifying poultry farming, offering consumers new tastes and improving meat production in order to meet the growing demand for animal protein [1]. However, it is worth pointing out that currently, there are no commercial feed formulas for quails in livestock feed industries in Côte d'Ivoire, where coturnic farmers generally use feed intended for chickens, where the high selling price of quail on the market [2, 3]. The art of rearing and adapting animals to their dietary needs is one of the most significant crucial human activities. Today, the main objective of the research addressed by this activity consists in developing techniques allowing

not only the optimal expression of the performance of animal species subjected to breeding but also and above all guaranteeing their sanitary quality [4]. This study aims to assess the impact of the consumption of four foods including three (03) newly formulated foods (A1, A2 and A3) based on local products then a commercial food or control food (AT) and ingested at Japanese quail (*Coturnix japonica*) raised in captivity for 9 weeks in Côte d'Ivoire on their biological health.

I- MATERIALS AND METHODS

I-1- Materials

I-1-1- Animal material

The animal material used for experimentation in this work is exclusively composed of one-day-old Japanese quail (*Coturnix japonica*).

The animal material used as animal protein sources are fish (*Sardinella maderensis*) and garden snail (*Achatina fulica*).

I-1-2-Plant material

The plant material used in this work consists on the one hand of yellow maize grains (*Zea mays*) and on the other hand of néré seeds and pulp (*Parkia biglobosa*) harvested at physiological maturity.

I-2-Methods

I-2-1-Receipt of the quails

Eighty (80) one-day-old quails were used for breeding in a farm in the village of Akoué Santai located in the commune of Bingerville (Ivory Coast). These quails were labelled, weighed and divided into four (4) groups of twenty (20) quails. Finally, each group of animals was housed in a compartment according to the four (4) diets.

I-2-2-Formulation of diets

Three (3) types of food were formulated:

- Food A1: The animal protein powder is that of fish (*Sardinella maderensis*) (100%);
- Food A2: The animal protein powder is that of the snail (*Achatina fulica*) (100%);
- Food A3: The animal protein powder is that of a powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; W/W).

For each of these formulations, the main carbohydrate constituent is exclusively yellow corn (*Zea mays*) powder. Then, were added to these foods, the same quantities of *Parkia biglobosa* powder (grains and pulp) as well as the other usual inputs as detailed in the table below.

Table 1: Centesimal composition of formulated foods A1, A2 and A3

Quantity per 100 kg of feed

Ingredients	Launch stage			Growth		
Corn	56	56	56	58	58	58
Nere pulp flour	3	3	3	3	3	3
Nere seed meal	20,8	20,8	20,8	19	19	19
Fishmeal	15	00	7,5	14,5	00	7,5
snail meal	00	15	7,5	00	14,5	7,5
Shell	2	2	2	2,2	2,2	2,2
Red oil	2	2	2	2	2	2
Vitamin complex	0.5	0.5	0.5	0.7	0.7	0.7
Salt	0.3	0.3	0.3	0.3	0,3	0.3
lysine	0.25	0.25	0.25	0.2	0.2	0,2
methionine	0.15	0.15	0.15	0.1	0.1	0,1

I-2-3-Breeding procedure

From the first day of reception, the day-old quails (young quails) were fed according to the following method:

- The 1st group received food A1 ;
- The 2nd group received food A2 ;

- The 3rd group received food A3 ;
- The 4th group received the commercial food AT (control food).

Every day at 5 p.m. the chicks received a very precise quantity of food (10g / day / head) during the first week and (15g / day / head) during the second week. Water containing vitamin (VMD-Amin special) and dewormer (Norfloxacin 20%) were also served to them. Twenty-four (24) hours later, the refusals were collected and weighed before renewing the food. Every 2 days, at 5 p.m., the quails were weighed. The start-up period lasted 2 weeks, then the chicks were transferred to the cages reserved for the growth phase which lasted about 7 weeks. The activities carried out during this phase were identical to those of the previous phase. During this phase the quantities of food received per quail per day varied from 17 to 25g from the third to the seventh week. At the end of each rearing period, three quails from each batch were sacrificed, their viscera and feathers were extracted and then weighed, as well as their carcasses. All the weighings were carried out using an electronic balance (ACCULAB Sartorius group-VICON 0.01g, CHINA).

I-2-4- Blood biochemical analyzes

I-2-4-1- Blood samples and serum preparations

At the end of the nine (9) weeks of experimentation, three (3) quails from each batch were fasted for a period of sixteen (16) hours. After that, a volume of five (5) ml of blood was taken per quail subjected to fasting and put in EDTA (Ethylenediaminetetraacetic acid) tubes. This blood was then centrifuged at 3000 rpm for 5 min in a centrifuge (SIGMA laborzentrifugen 3-16P, GERMANY). The serum (Should be plasma here !!! if blood collection was in EDTA tubes) obtained was stored in wells at -20°C for future biochemical analyzes using a Coulter ACT diff 2 (GEMANY) type automaton.

I-2-4-2- Dosage of serum proteins

The soluble serum proteins were assayed using the Biuret colorimetric method described by [5].

I-2-4-3- Dosage of serum glucose

Quail serum glucose was assayed using the colorimetric enzymatic method of [6]. The amount of serum glucose was calculated according to the following mathematical formula:

$$QG \left(\frac{g}{l} \right) = \frac{DO \text{ sample}}{DO \text{ standard}} \times 100$$

100 = standard concentration (mg/dL)

DO sample: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with glucose (0.5 g/L) ;

QG: quantity of glucose (g/L).

I-2-4-4-Determination of serum urea

Quail serum urea assay was performed using the method previously described by [7] with some modifications. Ten (10) μl of quail blood serum and ten (10) μl of urea (6.66 mmol/L) (standard) were diluted in each case in 1 mL of reagent (urease solution). Then, the two reaction media were homogenized by manual stirring for 1 min at ambient temperature (28° C.), then incubated at 37° C in a water bath for 10 min. The color intensities of these reaction media were determined using a spectrophotometer (MS-V5100 visible, CHINA) at 590 nm against a control containing no urea. The urea rate was calculated from the following mathematical equation:

$$\text{UREA } \left(\frac{\text{g}}{\text{l}} \right) = \frac{\text{DO test}}{\text{DO standard}}$$

DO test: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with urea (6.66 mmol/L).

I-2-4-5-Dosage of uric acid

Uric acid was measured using the enzymatic method of [8].

$$\text{URIC ACID } \left(\frac{\text{mg}}{100\text{ml}} \right) = \frac{\text{DO test}}{\text{DO standard}} \times 6$$

6 = standard concentration (mg/dL)

DO test: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with uric acid (6 mL /100 mL).

I-2-4-6-Determination of creatinine

In fact, creatinine in quail blood serum was measured according to the method of Hare [9].

The absorbance corresponding to each reaction medium was determined from the following mathematical relationship:

DO = DO 1min –DO 30s.

DO 1min = density of the test 1 min after incubation of the reaction medium at 37°C in a water bath;

DO 30s = density of the test 30s after incubation of the reaction medium at 37°C in a water bath.

The creatinine level was obtained from the following equation:

$$\text{Creatinine level} \left(\frac{\text{mg}}{\text{dl}} \right) = \frac{\text{DO test}}{\text{DO standard}} \times 2$$

2 = standard concentration (mg/dl)

DO 30 s: absorbance obtained after 30 seconds of incubation;

DO 1 min: absorbance obtained after 1 min of incubation.

I-2-4-7-Dosage of total cholesterol

Quail blood serum cholesterol was determined according to the colorimetric method of [6] using cholesterase. The absorbance of each reaction medium (DO) was calculated according to the following mathematical formula:

DO = DO 30 sec - DO 1 min

DO: absorbance of the test or of the standard;

DO 30 s: absorbance obtained after 30 seconds of incubation;

DO 1 min: absorbance obtained after 1 min of incubation.

The quail blood serum cholesterol level was obtained from the following mathematical expression:

$$\text{Cholesterol level} \left(\frac{\text{mg}}{\text{dl}} \right) = \frac{(\text{DO sample})}{(\text{DO standard})} \times 200$$

200 = standard concentration (mg/dL)

DO sample: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with cholesterol (5.17mmol/L).

I-2-4-8-Dosage of triglycerides

Quail blood serum triglycerides were measured according to the colorimetric method of [10] using glycerophosphate oxidase. The absorbance corresponding to each reaction medium was calculated from the following equation:

DO = DO 30s – DO 1min

DO: absorbance of the test or of the standard;

DO 30s: absorbance obtained after 30 seconds of incubation;

DO 1min: absorbance obtained after 1 min of incubation.

The quail blood serum triglyceride level was obtained from the following mathematical reaction:

$$\text{Triglyceride level} \left(\frac{\text{mg}}{\text{dl}} \right) = \frac{\text{DO sample}}{\text{DO standard}} \times 200$$

200 = standard concentration (mg/dL);

DO sample: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with triglyceride (2.28 mmol/L).

I-2-4-9-Calcium dosage

Quail blood serum calcium was measured according to the colorimetric method of [11] [Stern & Lewis \(1957\)](#). The calcium level was calculated from the mathematical relationship :

$$\text{Calcium level } \left(\frac{\text{mg}}{\text{dl}}\right) = \frac{\text{DO sample}}{\text{DO standard}} \times n$$

n = standard concentration (mg/dl);

DO sample: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with calcium (2.5 mmol/L).

I-2-4-10-Dosage of phosphorus

Quail blood serum phosphorus was measured according to the colorimetric method of [12].

The phosphorus rate was calculated from the following mathematical relationship:

$$\text{Phosphorus } \left(\frac{\text{mg}}{\text{dl}}\right) = \frac{\text{DO sample}}{\text{DO standard}} \times 50$$

50: standard concentration (mg/dL);

DO sample: absorbance obtained with quail blood serum;

DO standard: absorbance obtained with phosphorus (1.61 mmol/L).

II- RESULTS AND DISCUSSION

II-1-Results

II-1-1-Protein and triglyceride levels

The blood serum protein levels of quail fed with newly formulated feed (A1, A2, A3) and commercial feed (AT) at the end of rearing are respectively 49 ± 0.51 ; 43 ± 1.01 ; 55.67 ± 1.70 and 55 ± 0.82 g/l. Those of the blood sera of quails fed with A3 and AT foods are statistically ($P \geq 0.05$) identical. They are statistically higher than the blood serum protein levels of quails fed the newly formulated feeds A1 and A2 (Figure 1).

The blood serum triglyceride levels of quail fed with newly formulated feed (A1, A2, A3) and commercial feed (AT) at the end of rearing are respectively 13 ± 0.21 ; 9 ± 1.11 ; 6.8 ± 1.09 and 11.9 ± 0.02 g/l. The descending order of blood serum triglyceride levels of quails fed with newly formulated growth feed A1, A2, A3 and commercial feed (AT) is as follows: triglyceride level of growth feed A1 > triglyceride level of commercial growth food AT > triglyceride level of growth food A2 > triglyceride level of growth food A3 (Figure 1).




Figure 1: Protein and triglyceride levels of blood serum of quails fed with newly formulated feed and commercial feed

The same letters assigned to the means mean that they are not different at the 5% threshold.

- A1: food whose major animal protein source is only *Sardinella maderensis* fish powder (100%);
- A2: food whose main animal protein source is only the powder of the *Achatina fulica* snail (100%);
- A3: food whose major animal protein source is powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; w/w);
- AT: Commercial food (control food).

II-1-2-Phosphorus and calcium levels

The phosphorus levels of the blood serums of the quails fed with the newly formulated feeds A1, A2, A3 and the commercial feed AT at the end of rearing are respectively 0.051 ± 0.04 ; 0.051 ± 0.10 ; 0.050 ± 0.01 and 0.046 ± 0.12 g/l. Those of the blood sera of the quails fed with the newly formulated food A1, A2 and A3 are statistically ($P \geq 0.05$) identical to that of the quails fed with the commercial food AT (**Figure 2**).

The calcium levels of the blood serums of the quails fed with the newly formulated feeds A1, A2, A3 and the commercial feed AT at the end of rearing are respectively 0.094 ± 0.01 ; 0.095 ± 0.02 ; 0.097 ± 0.01 and 0.093 ± 0.03 g/l. All these calcium levels are statistically ($P \geq 0.05$) identical to each other (**Figure 2**).




Figure 2: Phosphorus and calcium levels in blood serum of quails fed with newly formulated feed and commercial feed

The same letters assigned to the means mean that they are not different at the 5% threshold.

- A1: food whose major animal protein source is only *Sardinella maderensis* fish powder (100%);
- A2: food whose main animal protein source is only the powder of the *Achatina fulica* snail (100%);
- A3: food whose major animal protein source is powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; w/w);
- AT: Commercial food (control food).

II-1-3- Ca/P ratios

The Ca/P ratios of the blood sera of the quails fed with the newly formulated growth feeds A1, A2, A3 and the commercial feed AT at the end of rearing are respectively 1.82 ± 0.04 ; 1.86 ± 0.12 ; 1.94 ± 0.19 and 2.03 ± 1.03 . They are statistically ($P \geq 0.05$) all identical (Figure 3).




Figure 3: Ca/P ratios of blood serum of quail fed with newly formulated feed and commercial feed.

The same letters assigned to the means mean that they are not different at the 5% threshold.

- A1: food whose major animal protein source is only *Sardinella maderensis* fish powder (100%);
- A2: food whose main animal protein source is only the powder of the *Achatina fulica* snail (100%);
- A3: food whose major animal protein source is powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; w/w);
- AT: Commercial food (control food).

II-1-4-Glycemic and cholesterol levels

The blood serum glycaemia levels of quails fed with newly formulated feed (A1, A2, A3) and commercial feed AT at the end of rearing are respectively 2.04 ± 1.07 ; 1.97 ± 1.10 ; 2.06 ± 1.09 and 1.48 ± 0.40 g/l. Those of the blood sera of the quails fed with the newly formulated foods A1, A2 and A3 are statistically ($P \geq 0.05$) the same. They are higher than that of the blood serum of quails fed with commercial growth food AT (Figure 4).

The blood serum cholesterol levels of quails fed with the newly formulated feed (A1, A2, A3) and the commercial feed AT at the end of rearing are respectively 1.87 ± 0.07 ; 1.85 ± 0.10 ; 1.83 ± 0.09 and 2.24 ± 0.02 g/l. Those of the blood sera of the quails fed with the newly formulated foods A1, A2 and A3 are statistically ($P \geq 0.05$) identical. They are statistically lower than those of the blood sera of quails fed with commercial food AT (Figure 4).




Figure 4: Blood serum glucose and cholesterol levels of quail fed the newly formulated feed and the commercial feed

The same letters assigned to the means mean that they are not different at the 5% threshold.

- A1: food whose major animal protein source is only *Sardinella maderensis* fish powder (100%);
- A2: food whose main animal protein source is only the powder of the *Achatina fulica* snail (100%);
- A3: food whose major animal protein source is powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; w/w);
- AT: Commercial food (control food).

II-1-5-Urea, uric acid and creatinine levels

The blood serum urea levels of quail fed with newly formulated growth feed A1, A2, A3 and commercial feed AT at the end of rearing are respectively 0.18 ± 0.03 ; 0.2 ± 0.12 ; 0.17 ± 0.03 and 0.16 ± 0.05 g/l. They are statistically ($P \geq 0.05$) all identical (Figure 5).

The uric acid levels of the blood serum of the quails fed with the newly formulated growth feeds A1, A2, A3 and the commercial feed AT at the end of rearing are respectively 0.05 ± 0.01 ; 0.05 ± 0.00 ; 0.05 ± 0.00 and 0.07 ± 0.00 g/l. They are statistically ($P \geq 0.05$) identical to each other (Figure 5).

The creatinine levels of the blood serums of the quails fed with the newly formulated growth feeds A1, A2, A3 and the commercial feed AT at the end of rearing are respectively 0.03 ± 0.01 ; 0.03 ± 0.00 ; 0.03 ± 0.00 and 0.03 ± 0.01 g/l. They are statistically ($P \geq 0.05$) identical (Figure 5).




Figure 5: Levels of urea, uric acid and creatinine in the blood serum of quails fed with the newly formulated feed and the commercial feed

The same letters assigned to the means mean that they are not different at the 5% threshold.

- A1: food whose major animal protein source is only *Sardinella maderensis* fish powder (100%);
- A2: food whose main animal protein source is only the powder of the *Achatina fulica* snail (100%);
- A3: food whose major animal protein source is powder composed of snail (*Achatina fulica*) and fish (*Sardinella maderensis*) (50/50; w/w);
- AT: Commercial food (control food).

II-2-Discussion

Blood biochemical parameters refer to substances synthesized by the body whose concentrations are relatively constant. However, the serum biochemical parameters of some birds could vary depending on several factors, including diet, age, breed, sex, physiological state and husbandry management [13, 14]. Their variations make it possible to assess the functional state of the organism [15, 16]. Consequently, an excess or a deficit of production of a substance synthesized by an organ is indicative of a dysfunction of this one. It is both a means of indirect exploration of the regulatory organs of metabolism and a means of assessing nutrient metabolism [17]. Proteins that are polymers of amino acids include albumin, globulins and fibrinogen. They are important in that they exert a beneficial effect on health through the maintenance of osmotic pressure, the transport of molecules, the purification of plasma, the strengthening of the immune system and the coagulation of blood [18, 19]. In this study, quails fed A3 and AT feeds show the highest respective crude protein levels (55.67 ± 1.70 and 55 ± 0.82 g/l). These high levels of serum proteins obtained in these

quails could be justified by the high levels of proteins contained in the A3 and AT feeds consumed by these birds. Our results corroborate those of [20]. Indeed, these authors, in their work on breeding systems, mentioned that proteinemia is higher in the conventional breeding system. Additionally, [5] then [21] reported that the level of total serum protein depends on the protein content of the diet. Thus, a high protein diet would lead to an increase in serum total protein content beyond the recommended ranges. In this work, the serum protein levels obtained could reassure us as to the good health of the quails studied.

Plasma electrolytes (calcium and phosphorus) play an important role in the growth of bones and teeth. They are involved in maintaining the body's hydrodynamic balance [22]. In this study, quails fed the newly formulated diets showed higher levels of these electrolytes compared to quails fed the commercial diet. In addition, the blood Ca/P ratio which represents the index of a good metabolism of these electrolytes, between 1 and 2, is in agreement with the recommendations of [22].

Urea and creatinine are metabolites resulting from the catabolism of total serum proteins and creatine synthesized by muscles and the brain. Their determination makes it possible to evaluate the effect of the consumption of food formulations on the glomerular function of the kidneys of subjects [23]. The average levels of urea and creatinine of all the quails studied are identical to those of the control quails; which could indicate normal kidney function in the quails studied. This result is supported by the determination of the blood uric acid of the quails. Indeed, this substance is the main end product of nitrogen metabolism in birds. Its secretion takes place mainly in the liver [24], and in the renal tubules [25]. The determination of plasma uric acid concentrations is widely used in birds for the purpose of assessing renal function. The highest serum uric acid level in this study is that obtained in quail fed with commercial food AT (66 ± 0.82 mg/l). This further substantiates that the newly formulated foods would not negatively affect kidney function.

Quails fed the newly formulated feeds had increased blood glucose levels compared to those fed the control feed. This increase could be due to the incorporation of **nere pulp** (?? Or write scientific name also) into newly formulated feeds. Indeed, **néré pulp** is said to be very rich in carbohydrates (60-85%) [26, 27].

The determination of cholesterol and triglyceride levels also made it possible to evaluate the effect of food consumption on the health status of the quails. Indeed, cholesterol and triglycerides are all fatty substances found in blood, bile and brain tissue [28]. They serve as precursors to the formation of bile acids, steroids and vitamin D [28]. It appears from the statistical analyzes that there is no significant difference in the level of cholesterol levels of

Formatted: Font color: Red

quails fed with the newly formulated feeds A1, A2 and A3 at the 5% threshold. Quails fed with commercial food AT have the highest cholesterol level (2.24 ± 0.04 g/l). Quails fed with feed A3 are those with the lowest level of triglycerides (6.80 ± 0.02 g/l). Consumption of the newly formulated feeds A1, A2 and A3 by quails would be beneficial as it could protect the animals against many metabolic and coronary diseases such as diabetes mellitus, nephropathy, biliary obstruction and various metabolic abnormalities due to endocrine disorders[24].

CONCLUSION

The objective of this study is to ensure that newly formulated feeds made from local products and ingested to quail raised in captivity for 9 weeks did not affect their biological health. The biochemical blood analyzes carried out at the end of the experiment revealed to us that the quails fed with the newly formulated feed have good renal, hepatic, pancreatic and cardiac health like that of the quails fed with the commercial feed AT.

REFERENCES

1. **Ukashatu S., Bello A., Umaru M. A., Onu J. E., Shehu S. A., Mahmuda A. & Saidu B.A.(2014).** Study of some serum biochemical values of Japanese quails (*Coturnix c. japonica*) fed graded levels of energy diets in Northwestern Nigeria. *Scientific Journal of Microbiology*, **3**: 1-10.
2. **Vali N. (2006).** Growth, feed consumption and carcass composition of *Coturnix japonica*, *Coturnix ypsilophorus* and their reciprocal crosses. *Asian Journal of Poultry Science*, **4**: 132-137.
3. **Djitie K.F., Kana J.R., Ngoula F., Nana N.F.C. & Tegua A. (2015).** Effect of crude protein level on growth and carcass in finishing phase quail (*Coturnix sp*) in the Cameroon Highlands. *Livestock Research for Rural Development*, **27**: 1-10.
4. **Boukhelifa A. (1993).** Study of poultry production parameters in the broiler and laying sector. Technico-economic impact on the development of poultry farming in Algeria:

the case of organic production factors (hatching egg, day-old chick flesh and started hen. Master thesis National Institute of Agronomy El Harrach.

5. **Gornall A.G., Bardawill C.J. & David M.M. (1949).** Détermination of serum protéins by means of the biuret reaction. *Journal Biology Chemistry*, **177**: 66-751.
6. **Trinder P. (1969).** Determination of Glucose in Blood Using Glucose Oxidase with an Alternative Oxygen Acceptor. *Annals of Clinical Biochemistry*, **6**: 24-27.
7. **Takemoto S., Tomonaga S., Funaba M. & Matsui T. (2017).** Effect of long-distance transportation on serum metabolic profiles of steer calves. *Animal Science Journal*, **88**: 1970–1978. doi: 10.1111/asj.12870.
8. **Fossati P. & Prencipe L. (1980).** Use of 3,5-dichloro-2-hydroxybenzensulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. *Clinical Chemistry*, **2**: 227-231.
9. **Hare R.S. (1950).** Endogenous creatinine in serum and urine. *Proceedings of Society Experimental Biology and Medecine*, **74**: 148.
10. **Fossati P. & Prencipe L. (1982).** Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clinical Chemistry*, **28**: 2077-2080.
11. **Stern J. & Lewis H. P. W. (1957).** The colorimetric estimation of calcium in serum with o-cresolphthalein complexone. *Clinical Chimistry Acta*, **6**: 576-580.
12. **Baginski E.S., Fao P.P. & Zak B. (1967).** Détermination of phosphate: study of labile organic phosphate interference. *Clinica Chimical Acta*, **1**: 155-158.
13. **Addass PA, David DL, Edward A., Zira K. E. & Midau A. 2012.** Effect of Age, Sex and Management System on Some haematological parameters of intensively and semi-intensively kept chicken in Mubi, Adamawa State, Nigeria. *Iranian Journal of Applied Animal Science*, **2** (3): 277-282.
14. **Koukoura K. K., Brah M., Kantati Y. T. & Tchacondo T. (2021).** Evaluation de l'effet trophique de l'huile de neem (*Azadirachta indica*) chez les poulets de chair de

souche Sasso. *International Journal of Biological and Chemical Sciences*, **15**(6): 2402-2416.

- 15. Melillo A. (2007).** Rabbit clinical pathology. *Journal of Exotic Pet Medicine*, **16**: 135-145.
- 16. Jenkins J.R. (2008).** Rabbit diagnostic testing. *Journal of Exotic Pet Medicine*, **17**: 4-15.
- 17. Bouafou K.G.M. (2007).** Etude de la production d'asticots à partir d'ordures ménagères biologiques et de la valeur nutritionnelle de la farine d'asticots séchés (f.a.s) chez le rat en croissance. Thèse de doctorat de l'Université de Cocody-Abidjan (Côte d'Ivoire), 195 p.
- 18. Koller A. & Kaplin L.A.(1989).** Total serum protein: Theory, analysis and Correlation, 2nd edition St. Louis: The C.V. Mosby Compagny, *Clinical Chemistry*, **1989**: 1057-1060.
- 19. Piotrowska A., Burlikowska K. & Szymeczko R. (2011):** Changes in blood chemistry in broiler chickens during the fattening period. *Folia biologica* (Krakow), **59** (3-4): 183-187.
- 20. Mingoas K.J.P., Awah-Ndukum J., Mampom B.J., Mfopit M.Y. & Zoli P. A. (2017).** Effets du système d'élevage sur les performances zootechniques et les paramètres sanguins et biochimiques chez les poulets de chair en zone péri-urbaine de Ngaoundéré, Cameroun. *Journal of Animal & Plant Sciences*, **32** (1): 5079-5094.
- 21. Atchadé G. S. T., Segbotangni E. F., Mensah S. E. P., Houndonougbo M. F., Attakpa S. E. & Crysostom C. A. A. M. (2019).** Les grains de céréale dans l'alimentation des poulets au Bénin : digestibilité métabolique et paramètres biochimiques sériques induits. *Afrique SCIENCE*, **15** (5):25 – 38.

22. Koinarski V., Angelov G. & Lalev M. (2001). Haematologic and blood electrolytes changes in broiler chickens experimentally invaded with *Eimeria tenella*. *Bulgarian Journal of Veterinary Medicine*, **4**: 237-242.
23. Wyss M. & Kaddurah-Daouk R. (2000): Creatine and creatinine metabolism. *Physiological reviews*, **80**: 1107-1213.
24. Lumeij J.T. (1987): A contribution to clinical investigative methods for birds, with special reference to the racing pigeon (*Columba livia domestica*). Utrecht, 186 p.
25. Chou S.T. (1972). Relative importance of liver and kidney in synthesis of uric acid in chickens. *Canadian Journal of Physiology and Pharmacology*, **50**: 936-939.
26. Bonkougou E. (1987). Monographie du néré, *Parkia biglobosa* (Jacq) Benth, espèce agroforestière à usage multiples IRBETI CNRST - Ouagadougou (Burkina Faso), 45 p.
27. Touré M. (2018). Le néré, un arbre du patrimoine de la Haute Guinée. *Belgeo, Revue belge de géographie*, **2** : 1-25.
28. Talebi A. (2006). Biochemical parameters in broiler chickens vaccinated against ND, IB and IBD. *International Journal of Poultry Science*, **5**: 1151-1155.