

### EFFECT OF MACRONUTRIENTS COMBINATION WITH PLANT SPACING ON THE GROWTH AND YIELD OF BLACK CUMIN (*Nigella sativa* L.)

#### Abstract

An experiment was conducted at the Horticulture Farm of Sher-e-Bangla Agricultural University, Dhaka during October 2019 to March 2020 to study the effect of macronutrients combination and plant spacing on growth and yield of black cumin (*Nigella sativa* L.). Black cumin variety BARI Kalozira 1 was used as planting material in this study. The experiment consisted of two factors: Factor-A: macro nutrient combinations (4 levels): T<sub>1</sub>= N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>2</sub>= N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>3</sub>= N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup>; Factor-B: plant spacing (3 levels): S<sub>1</sub>= 20 cm × 10 cm, S<sub>2</sub>= 20 cm × 15 cm and S<sub>3</sub>= 20 cm × 20 cm. The experiment was laid out in a randomized complete block design (factorial) with three (3) replications. Data on different growth, yield contributing and yield parameter of black cumin were recorded and significant variation was observed from different treatments. In case of nutrient combinations the tallest plant (54.86 cm) was observed from T<sub>3</sub> (N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup>) treatment. The maximum primary branch plant<sup>-1</sup> (8.62) and secondary branch plant<sup>-1</sup> (12.18), flower plant<sup>-1</sup> (22.20), capsules plant<sup>-1</sup> (19.69) and 1000 seed weight (2.99 g) was observed from T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment. The highest seed yield ha<sup>-1</sup> (1.18 t) was observed from T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment. In case of plant spacing the tallest plant (48.34 cm) was observed from S<sub>1</sub> (20 cm × 10 cm) treatment. The maximum primary branch plant<sup>-1</sup> (7.80) and secondary branch plant<sup>-1</sup> (11.23) flower plant<sup>-1</sup> (20.26), capsules plant<sup>-1</sup> (18.81), 1000 seed weight (2.99 g) were observed from S<sub>3</sub> (20 cm × 20 cm) treatment. The highest seed yield ha<sup>-1</sup> (1.09 t) was recorded from S<sub>1</sub> (20 cm × 10 cm) treatment. It can be concluded that, sowing of black cumin providing 90 kg N, 50 kg P and 40 kg K nutrient combination with 20 cm × 10 cm plant spacing was recorded to be more suitable practice for getting higher amount and quality of seed yield of black cumin.

**Keyword: macronutrients; plant spacing; growth; yield; black cumin.**

#### 1.Introduction:

Black cumin (*Nigella sativa* L.) is well known as a spice crop in Bangladesh as well as in the world. It is commonly known as 'Kalozira' belongs to family Ranunculaceae and is cultivated in the winter season. It is believed to have originated in western southeast Asia. It is widely cultivated in Southwest Asia, South Europe, Syria, Egypt, Pakistan, India, Iran, Japan, China and Turkey[1]. In Bangladesh, it is grown well in Faridpur, Sariatpur, Madaripur, Pabna, Sirajganj, Jessore, Kusthtia and Natore districts. The flowers are delicate, and usually coloured pale blue and white with 5 to 10 petals. The fruit is a large and inflated capsule composed of 3 to 7 united follicles, each containing throughout Black cumin is numerous seeds. The seed is used as a spice [2]. Black Cumin has a long history of use as food flavors, perfumes and medicinal values. Essential oil has been used for bringing smell to some medicines, for sterilizing of surgical operation fiber, for producing of some veterinary and agricultural medicines and plastic [3]. The seed contain 30-35 % of oil which has several uses for pharmaceutical and food industries [4]. The medicinal value of the spice is immense and numerous workers appreciated its unique, varied and powerful pharmacological traits. The popularity of the plant was highly enhanced by the ideological belief in the herb as a cure for multiple diseases likes anti-tumour, anti-diabetic, cardioprotective, gastroprotective, antibacterial and antifungal activities were immensely appreciated. On Bangladesh black cumin was cultivated on 31 hectares of land and average yield 0.8-1.0 ton per hectares[5].

Fertilizer application to the plant greatly effects their growth, production plant constituents. Nitrogen (N) has the largest effect on plant physiology and is probably the most important limiting nutrient for crop growth. Nitrogen strongly stimulates growth expansion of the plant canopy and ultimately increases yield contributing characters and gross yield. Availability of nitrogen is of most important for growing plants as it is a major and indispensable constituent of protein and nucleic acid molecules [6]. An adequate supply of nitrogen is associated with vigorous vegetative growth and more efficient use of available inputs finally lead to higher productivity. Phosphorus (P) is essential for the

general health of the plant and root development and more stem strength. It improves flower formation and makes seed production more uniform. It also improves seed quality and resistant to plant disease ([plz provide supportive reference](#)). The early supply of phosphorous to the crop is influenced by soil phosphorous and phosphorous application affect phosphorous phyto availability and root growth. Roots absorb phosphorous ions from the soil solution [7]. Potassium (K) fertilizer is another essential component to reduce the severity of disease of black cumin plants. Application of K ha along with N ha<sup>-1</sup> decreased plant death due to disease and also increased yield in black cumin [8].

Plant spacing plays an important role in growth and yield. Optimum plant spacing ensures the plant to grow properly with their aerial and underground parts by utilizing more sunlight and soil nutrients [9]. High plant spacing make difficult for various intercultural operations. In a densely populated crop, the interplant competition is very high for essential nutrients, light and air, which usually results in mutual shading, lodging and thus favors more vegetative growth than grain yield. Black cumin lacks research recommendations on optimum plant density and row spacing that could lead to the poor yield of the crop. Hence, it is vital to identify the suitable seed rates and row spacing for black cumin production. Therefore, this study was initiated with the objective of finding out the optimum seed rate and inter- row spacing for better growth and yield attributes of black cumin.

## 2. MATERIALS AND METHODS

### 2.1 Experimental site and experimental framework

The experiment was carried out at Horticultural Farm in Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh. It is located at 23° 41' N latitude and 90° 22' E longitude at an altitude of 8.6 meters above the sea level. The land belongs to Agro-ecological zone of Modhupur Tract, AEZ-28. The selected land was medium high land with adequate irrigation facilities. The physical and chemical properties of soil of the experimental site was sandy loam in texture having pH 5.47 - 5.63. Organic matter content was very low (0.83).

### 2.2 Seed

In this experiment black cumin variety of BARI Kalozira 1 was used as a planting material. BARI Kalozira 1 was developed by Bangladesh Agricultural Research Institute (BARI) in 2009. The seed was collected from the Regional Spice Research Centre, BARI, Joydebpur, Gazipur.

### 2.3 Treatments

The experiment consisted of two factor as: Factor A: Fertilizer (4 levels) T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup> and Factor B: spacing (3 levels) S<sub>1</sub> = 20 cm×10 cm, S<sub>2</sub> = 20 cm×15 cm and S<sub>3</sub> = 20 cm×20 cm. The two factorial experiment was laid out in Randomized Complete Block Design (RCBD) with three replications to minimize the soil heterogeneous effects. The experiment was divided into three blocks and consisted of 36 plots. Each unit plot in from of raised bed was 1.44 m<sup>2</sup> (1.2m x 1.2m) in size. Altogether there were 36 unit plots in experiment and required 51.84 m<sup>2</sup> land. Row to row distance was 20 cm and plant to plant distance at S<sub>1</sub> S<sub>2</sub> S<sub>3</sub> plot were 10 cm, 15 cm and 20 cm respectively. The treatments were randomly assigned to each of the block. Each unit plot had 6 rows and each with 6, 9 and 12 plants. So there were 36, 54 and 72 plants at S<sub>1</sub> S<sub>2</sub> and S<sub>3</sub> plot respectively

### 2.4 Fertilizer management

Chart 1: Following doses of manures and fertilizers were recommended for the black cumin production.

Manure and Fertilizer		Dose
Cow dung		10 t/ha
Kitchen compost		5 t/ha
N	T <sub>1</sub>	45 kg/ha
	T <sub>2</sub>	90 kg/ha
	T <sub>3</sub>	135 kg/ha

P <sub>2</sub> O <sub>5</sub>	T <sub>1</sub>	25 kg/ha
	T <sub>2</sub>	50 kg/ha
	T <sub>3</sub>	75 kg/ha
K <sub>2</sub> O	T <sub>1</sub>	20 kg/ha
	T <sub>2</sub>	40 kg/ha
	T <sub>3</sub>	60 kg/ha
Gypsum		112 kg/ha
Boron		24 kg/ha

The above doses of fertilizers were converted into manure and fertilizer mixed per treatment of the experiment and supplied by each type of manure and fertilizer except control. After conversion the dose of each manure used in the experiment was as below.

Chart 2: Each manure used in the experiment

Treatments	Cowdung (kg/ha)	Kitchen compost (kg/ha)	Urea (g/ha)	TSP (g/ha)	MOP (g/ha)	Gypsum (g/ha)	Boron (g/ha)
T <sub>0</sub>	1.44	0.72	-	-	-	16.13	3.5
T <sub>1</sub>	1.44	0.72	14.11	18	5.76	16.13	3.5
T <sub>2</sub>	1.44	0.72	28.22	36	11.52	16.13	3.5
T <sub>3</sub>	1.44	0.72	42.19	54	17.28	16.13	3.5

Note:

T<sub>0</sub> = Control, T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup>

The calculated amounts of cowdung, kitchen compost, triple super phosphate, muriate of potash, gypsum, boron and half of the urea were applied before seed sowing. Rest half of the Urea was applied in two equal splits at 25 and 50 days after seed sowing.


## 2.5 Data analysis

The recorded data on different parameters were statistically analyzed using Statistic 10 software. The significance of the difference among the treatments means was estimated by the least significant difference test (LSD) at 5% level of probability.

## 3. Result and discussion


### 3.1 Plant height (cm)

Different macro nutrients (N, P, K) had significant influence on plant height of black cumin at different growth stages (Fig. 1). At 135 DAS, the highest plant height (54.86 cm) was obtained from  $T_3$  ( $N_{135}P_{75}K_{60}$ ) treatment. Similarly, the lowest plant height (32.89 cm) was recorded from the  $T_0$  (control) treatment. Hence it may be inferred that the increase in plant height may be due to the favorable influence and balanced absorption of nutrients, increased role of photosynthesis, reduced transpiration and stimulation of root system, increase cell division, cell enlargement and metabolic processes. It is also observed that plant height increased with macronutrient doses [10][11]. Different spacing had significant variation on plant height of black cumin at different growth stages (Fig. 2). Plant height increased with decreased plant spacing. At 135 DAS, the highest plant height (48.34 cm) was achieved from  $S_1$  (20 cm x 10 cm) treatment. Again, the lowest plant height (45.72cm) was observed from  $S_3$  (20 cm x 20 cm) treatment. The treatment  $S_1$  was statistically identical to  $S_2$  which gave plant height of 18.13 cm at 45 DAS. The variation in plant height as influenced by spacing was perhaps due to proper utilization of nutrient, moisture and light(Fig.2).[12] was observed that plant height was increased by decreasing plant spacing an antagonistic relationship was found between vegetative growth and plant spacing.[13] reported the smallest inter-row spacing (20 cm) produced the highest average plant heights while the lowest values were obtained at the largest inter-row spacing (40 cm) respectively. These findings on plant height were in accordance with[14][15][16]. Combined effect of different nutrients and spacing on plant height of black cumin was statistically significant at different days after sowing (DAS). At 135 DAS, the highest plant height (56.36 cm) was obtained from  $T_2S_1$  ( $N_{90}P_{50}K_{40}$  kg ha<sup>-1</sup> and 20 cm x 10 cm) treatment



combination, which was statistically similar to  $T_2S_2$  ( $N_{90}P_{50}K_{40}$  kg ha<sup>-1</sup> and 20cmx15cm) treatment combination and the lowest plant height (32.20 cm) was observed from  $T_0S_3$  (control and 20cmx20 cm) treatment combination which was statistically identical to  $T_2S_1$ .

**Fig. 1. Effect of nutrients combination on plant height of black cumin (*Nigella sativa* L.)**(LSD value= 0.4479, 1.1080 and 0.5814 at 45, 90 and 135 DAS respectively), Note:  $T_0 = N_0P_0K_0$  kgha<sup>-1</sup>(control),  $T_1 = N_{45}P_{25}K_{20}$  kgha<sup>-1</sup>,  $T_2 = N_{90}P_{50}K_{40}$  kgha<sup>-1</sup>,  $T_3 = N_{135}P_{75}K_{60}$  kgha<sup>-1</sup>



**Fig. 2. Effect of plant spacing on plant height of black cumin (*Nigella sativa* L.)**(LSD value= 0.3879, 0.9596 and 0.5035 at 45, 90 and 135 DAS respectively) Note: S<sub>1</sub> = 20 cm x 10 cm, S<sub>2</sub> = 20 cm x 15 cm and S<sub>3</sub> = 20 cm x 20 cm.

**Table 1. Combined effect of nutrients combination and plant spacing on plant height at 45 DAS, 90 DAS and 135 DAS of black cumin (*Nigella sativa* L.)**

Treatments	Plant height (cm)		
	45 DAS	90 DAS	135 DAS
T <sub>0</sub> S <sub>1</sub>	10.49 h	20.71 fg	33.87 g
T <sub>0</sub> S <sub>2</sub>	11.55 g	21.75 f	32.60 h
T <sub>0</sub> S <sub>3</sub>	10.30 h	19.42 g	32.20 h
T <sub>1</sub> S <sub>1</sub>	14.20 e	32.27 d	48.33 e
T <sub>1</sub> S <sub>2</sub>	14.33 e	27.47 e	47.67 e
T <sub>1</sub> S <sub>3</sub>	13.40 f	26.53 e	46.53 F
T <sub>2</sub> S <sub>1</sub>	25.27 a	50.27 a	56.36 a
T <sub>2</sub> S <sub>2</sub>	24.62 ab	48.96 a	55.63 ab
T <sub>2</sub> S <sub>3</sub>	20.57 d	39.13 c	52.60 c
T <sub>3</sub> S <sub>1</sub>	23.91 b	42.27 b	54.81 b
T <sub>3</sub> S <sub>2</sub>	22.00 c	41.67 b	52.53 cd
T <sub>3</sub> S <sub>3</sub>	20.60 d	41.87 b	51.53 d
<b>LSD(0.05)</b>	0.7758	1.9191	1.0070
<b>CV%</b>	2.60	3.30	1.26

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup> and S<sub>1</sub> = 20 cm x 10 cm, S<sub>2</sub> = 20 cm x 15 cm, S<sub>3</sub> = 20 cm x 20 cm

### 3.2 Number of primary and secondary branches plant<sup>-1</sup>

There was a significant variation due to effect of nutrient in the number of primary and secondary branches plant<sup>-1</sup>. The maximum number of primary branches plant<sup>-1</sup> (8.62) and secondary branches plant<sup>-1</sup> (12.18) was observed at T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment while controlled treatment showed comparatively lower (5.50) and (8.60) number of primary and secondary branches plant<sup>-1</sup>. Respectively (Table 2). From the results of the present study indicated that optimum levels of macro nutrients combination might have induced better growing condition, perhaps due to supply of adequate plant nutrients which ultimately led to the production of more primary and secondary branches per plant.

[15][17]found that the number of primary and secondary branch plant<sup>-1</sup> of black cumin was influenced significantly under different level of fertilizer.

Significant variation was found due to the effect of spacing on number of primary and secondary branches plant<sup>-1</sup>. It was observed that the lowest number of primary and secondary branches plant<sup>-1</sup> (7.09) and (10.35) was recorded from S<sub>1</sub> (20 cm × 10 cm) treatment and highest (7.80) and (11.23) from S<sub>2</sub> (20 cm × 20 cm) treatment respectively (Table 3). Generally lower number of plants was provided more nutrients compared to higher population with same nutrient status in the soil that was provided and caused more number of primary branches per plant from higher plant spacing. These findings are in agreement with those of [18]. [19] attributed the increments in vegetative characteristics to less competition among plants for the environmental conditions necessary for building up more metabolites and producing more lateral branches at wider spaces.


Number of primary and secondary branches plant<sup>-1</sup> showed significant variation among the treatments due to the combined effect of macronutrients and plant spacing. Highest primary and secondary branches plant<sup>-1</sup> (8.80) and (12.47) was found from the T<sub>2</sub>S<sub>3</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>and 20cm × 20cm) treatment combination. On the other hand, lowest value for primary and secondary branches plant<sup>-1</sup> (4.96) and (8.20) was recorded in T<sub>0</sub>S<sub>1</sub> (control and 20cm × 10 cm) treatment combination (Table 4). Generally more nutrients and higher spacing plant get more food and space for vigorous growth and produce more primary branches compared to lower nutrients and spacing.

### 3.3 Number of flowers and capsules per plant


The number of flowers plant<sup>-1</sup>and capsule plant<sup>-1</sup> was significantly affected by different levels of nutrient combinations (Figure 3 and Table 2). Number of flowers and capsule plant<sup>-1</sup> increased with the increase level of nutrients. The highest number of flowers and capsule plant<sup>-1</sup> (22.20) and (19.81) was recorded at T<sub>2</sub>(N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kgha<sup>-1</sup>)treatment. The lowest number of flowers plant<sup>-1</sup> was recorded from T<sub>0</sub> (control) treatment (17.56) and (16.17). From the results of the present study indicated that number of flowers plant<sup>-1</sup> increased with the increase in nutrient doses. These findings are in agreement with those of [11].

Significant variations were clearly evident in case of number of flowers plant<sup>-1</sup> and capsule plant<sup>-1</sup> with different plant spacing (Figure 4 and Table 3). The highest number of flowers and capsule plant<sup>-1</sup> (20.26) and (18.81) resulted from S<sub>3</sub>(20 cm × 20 cm) treatment and the lowest (19.90) and (17.62) was obtained from S<sub>1</sub>(20 cm × 10 cm) treatment which was statistically identical (19.96) to S<sub>2</sub> (20 cm × 15 cm) treatment. From the results of the present study indicated that increase in plant density decreased number of flowers plant<sup>-1</sup>.

Combined effect of different macronutrient and spacing was statistically significant in respect of number of flowers plant<sup>-1</sup> (Table 4). The highest number of flower and capsule plant<sup>-1</sup> (22.87) and (20.80) obtained from T<sub>2</sub>S<sub>3</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>and 20 cm × 20 cm) treatment combination. The lowest number of flower and capsule plant<sup>-1</sup> (17.20) and (15.33) obtained from T<sub>0</sub>S<sub>1</sub> (control and 20 cm×10 cm) treatment combination respectively which was statistically identical (17.35) to T<sub>0</sub>S<sub>2</sub> (control and 20 cm×15 cm) treatment combination.



**Fig. 3. Effect of nutrients combination on number of flowers plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)** (LSD value= 0.2007) Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kgha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kgha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kgha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kgha<sup>-1</sup>



**Fig. 4. Effect of plant spacing on number of flowers plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)** (LSD value= 0.1738) S<sub>1</sub> = 20 cm x 10 cm, S<sub>2</sub> = 20 cm x 15 cm, S<sub>3</sub> = 20 cm x 20 cm

**Table 2. Effect of nutrients combination on number of primary branches plant<sup>-1</sup>, secondary branches plant<sup>-1</sup> and capsules plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)**

Treatments	Number of Primary branches plant <sup>-1</sup>	Number of Secondary branches plant <sup>-1</sup>	Number of capsules plant <sup>-1</sup>
T <sub>0</sub>	5.50 d	8.60 d	16.17 c
T <sub>1</sub>	7.42 c	10.98 c	17.04 b
T <sub>2</sub>	8.62 a	12.18a	19.81 a
T <sub>3</sub>	8.09 b	11.51 b	19.69 a
<b>LSD(0.05)</b>	0.2077	0.1622	0.2773
<b>CV%</b>	2.87	1.53	2.56

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kgha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kgha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kgha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kgha<sup>-1</sup>

**Table 3. Effect of plant spacing on number of primary branches plant<sup>-1</sup>, secondary branches plant<sup>-1</sup> and capsules plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)**

Treatments	Number of Primary branches plant <sup>-1</sup>	Number of Secondary branches plant <sup>-1</sup>	Number of capsules plant <sup>-1</sup>
S <sub>1</sub>	7.09 c	10.35 c	17.62 c
S <sub>2</sub>	7.34 b	10.87 b	18.10 b
S <sub>3</sub>	7.80 a	11.23 a	18.81 a
<b>LSD(0.05)</b>	0.1799	0.1404	0.2402

CV%

2.87

1.53

2.56

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: S<sub>1</sub> = 20 cm × 10 cm, S<sub>2</sub> = 20 cm × 15 cm, S<sub>3</sub> = 20 cm × 20 cm

**Table 4. Combined effect nutrients combination and plant spacing on primary branches plant<sup>-1</sup>, secondary branches plant<sup>-1</sup>, number of flowers plant<sup>-1</sup>, number of capsules plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)**

Treatments	Number of primary branches plant <sup>-1</sup>	Number of secondary branches plant <sup>-1</sup>	Number of flowers plant <sup>-1</sup>	Number of capsules plant <sup>-1</sup>
T <sub>0</sub> S <sub>1</sub>	4.96 g	8.20 h	17.20 g	15.33 f
T <sub>0</sub> S <sub>2</sub>	5.22 g	8.40 h	17.35 g	16.00 e
T <sub>0</sub> S <sub>3</sub>	6.33 f	9.20 g	18.13 f	17.19 d
T <sub>1</sub> S <sub>1</sub>	7.20 e	10.67 f	19.20 e	17.47 d
T <sub>1</sub> S <sub>2</sub>	7.67 d	11.07 e	18.87 e	16.27 e
T <sub>1</sub> S <sub>3</sub>	7.40 de	11.20 e	19.20 e	17.40 d
T <sub>2</sub> S <sub>1</sub>	8.47 abc	11.80 cd	21.53 c	18.22 c
T <sub>2</sub> S <sub>2</sub>	8.60 ab	12.29 ab	22.20 b	20.40 a
T <sub>2</sub> S <sub>3</sub>	8.80 a	12.47 a	22.87 a	20.80 a
T <sub>3</sub> S <sub>1</sub>	7.73 d	10.73 f	21.67 c	19.47 b
T <sub>3</sub> S <sub>2</sub>	8.13 c	11.73 d	21.40 c	19.74 b
T <sub>3</sub> S <sub>3</sub>	8.40 bc	12.06 bc	20.87 d	19.86 b
<b>LSD(0.05)</b>	0.3598	0.2809	0.3476	0.484
<b>CV%</b>	2.87	1.53	2.02	2.56

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability.

Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup> and S<sub>1</sub> = 20 cm × 10 cm, S<sub>2</sub> = 20 cm × 15 cm, S<sub>3</sub> = 20 cm × 20 cm

### 3.4 1000 seed weight (g)

Different levels of NPK fertilizer had significant effect on 1000 seed weight of black cumin (Table 5). Application of macronutrients at different levels significantly increased the 1000 seed which produced maximum seed weight (2.99 g) at T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment where T<sub>0</sub> (control) treatment gave the lowest 1000 seed weight (2.80 g). [20] stated that there were no statistical differences among the different nitrogen doses in black cumin. In different studies, thousand seed weight of black cumin was reported as 3.50 g.

1000 seed weight of black cumin was significantly influenced by different level of spacing (Table 6). It was observed that higher spacing gave maximum yield. The maximum 1000 seed weight (2.96 g) was recorded from S<sub>3</sub> (20 cm × 20 cm) treatment where the lowest 1000 seed weight (2.82 g) was recorded from S<sub>1</sub> (20 cm × 10 cm) treatment. Similar findings were also obtained by [11][21].


Combined effect of different levels of macronutrient and spacing proved significant differences on 1000 seed weight of black cumin (Table 7). Results revealed that the highest 1000 seed weight (3.06 g) was obtained from T<sub>2</sub>S<sub>3</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> with 20 cm × 20 cm) treatment combination which was statistically similar to T<sub>2</sub>S<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> and 20 cm × 15 cm) treatment combination. The lowest 1000 seed weight (2.75 g) was recorded from T<sub>0</sub>S<sub>1</sub> (control and 20 cm × 10 cm) treatment combination which was statistically similar to T<sub>0</sub>S<sub>2</sub> (control and 20 cm × 15 cm) and T<sub>1</sub>S<sub>1</sub> (N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup> and 20 cm × 10 cm) treatment combination. Rest of the treatment combination performed intermediate results in terms of 1000 seed weight compared to all other treatments. [22][23] also found significant effect of fertilizer levels on thousand seed weight of black cumin.

### 3.5 Yield per plant (g)


Yield per plant was significantly influenced due to different levels of macronutrient (Figure 5). Yield was increased with increasing plant nutrients. Results showed that the maximum yield per plant (3.36 g) was recorded from T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment and the lowest yield of per plant (2.51 g) was recorded from T<sub>0</sub>(control)treatment. From the above results, it was noted that NPK when used the nutrients become available to plants and much seed formation was occurred. The available soil nutrients supported proper vegetative growth with more protoplasm in the cells in comparison to less available nutrient in black cumin. The results found from the findings of [24][25] were similar with the present study.

Yield plant<sup>-1</sup> was significantly influenced by different level of spacing (Figure 6). It was observed that higher spacing gave maximum yield. The maximum yield plant<sup>-1</sup> (3.36 g) was recorded from S<sub>3</sub> (20 cm × 20 cm) treatment where the lowest yield plant<sup>-1</sup> (2.51 g) was recorded from S<sub>1</sub> (20 cm × 10 cm) treatment. It might be due to the fact that in treatment S<sub>3</sub> (20 cm × 20 cm) treatment received adequate plant nutrients, no inter competition among plants, favorable growing atmosphere which contributed yield plant<sup>-1</sup> (g). The result achieved from the present study was conformity with the findings of [26][21] who observed higher yield plant<sup>-1</sup> from wider spacing.

Combined effect of different levels of macronutrient and spacing proved significant differences on yield plant<sup>-1</sup> of black cumin (Table 7). Results revealed that the highest yield plant<sup>-1</sup> (4.34 g) was obtained from T<sub>2</sub>S<sub>3</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> and 20 cm × 20 cm) treatment combination. The lowest yield plant<sup>-1</sup> (1.87 g) was recorded from T<sub>0</sub>S<sub>1</sub> (control and 20 cm × 10 cm) treatment combination. Rest of the treatment combination performed intermediate results in terms of yield plant<sup>-1</sup> compared to all other treatments.



**Fig. 5. Effect of nutrients combination on yield plant<sup>-1</sup> of black cumin (*Nigella sativa* L.)**(LSD) value= 0.1211)Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kgha<sup>-1</sup>(control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kgha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kgha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kgha<sup>-1</sup>



**Fig. 6. Effect of plant spacing on yield plant<sup>-1</sup> of black cumin (*Nigella sativa* L.) (LSD value= 0.1049). Note: S<sub>1</sub> = 20 cm × 10 cm, S<sub>2</sub> = 20 cm × 15 cm, S<sub>3</sub> = 20 cm × 20 cm**

### 3.6 Yield per plot (g) and ha (t)

Yield per plot of black cumin was significantly affected by different levels of macronutrient (Table 5). Higher application of plant nutrients gave higher seed yield per plot. Results specified that the highest yield per plot and ha of black cumin (170.40 g) and (1.18 t) was recorded from T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment where the lowest yield per plot and ha of black cumin (127.14 g) and (0.88 t) was recorded from T<sub>0</sub> (control) treatment. The results obtained from the present study were similar with the findings of [27][11].

Different levels of spacing had significant effect on yield per plot of black cumin (Table 6). It was found that the highest yield per plot of black cumin (156.66 g) and (1.09 t) was recorded from S<sub>1</sub> (20 cm × 10 cm) treatment where the lowest yield per plot of black cumin (140.04 g) and (0.97 t) was recorded from S<sub>3</sub> (20 cm × 20 cm) treatment. The obtained results represented that maximum yield contributing characters was best with higher spacing but in case of yield closer spacing gave maximum yield and this result might be due to cause of higher plant population from closer spacing. [26] studied the effect of row spacing on seed yield and yield components of black cumin and found similar results on seed yield per plot with the present study.

Yield per plot of black cumin was significantly affected by combined effect of different levels of macronutrient and spacing (Table 7). It was observed that the highest yield per plot (178.32 g) and yield per ha (1.24 t) was obtained from T<sub>2</sub>S<sub>1</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> and 20 cm × 10 cm) treatment combination which is statistically identical to T<sub>2</sub>S<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> and 20 cm × 15 cm) combination. Results also revealed that the lowest yield per plot of black cumin (118.68 g) and yield per ha (0.82 t) was recorded from T<sub>0</sub>S<sub>3</sub> (control and 20 cm × 20 cm) treatment combination, which was statistically similar to T<sub>0</sub>S<sub>2</sub> (control and 20 cm × 15 cm) treatment combination. The results obtained from all other treatment combination gave intermediate results compared to highest and lowest results. Higher number of plant population need higher amount of nutrients. Under the present study, closer spacing with higher nutrient doses gave the higher yield and this type of achievement might be due to higher plant population.

### 3.7 Dry matter (g)

Significant influence was noted on dry matter affected by different macro nutrients (N, P, K) (Table 5). The highest dry matter at harvest (11.29 g) was obtained from the T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment where the

lowest dry matter at harvest (7.90 g) was obtained from the T<sub>0</sub> (control) treatment. These findings are also supported by [28] dry matter weight plant<sup>-1</sup> was increased gradually with the higher level of applied fertilizer.

Significant variations were clearly evident in case of dry matter with different plat spacing (Table 6). The highest dry matter resulted from S<sub>3</sub> (20 cm × 20 cm) treatment (18.81) and the lowest (17.62) was obtained from S<sub>1</sub> (20 cm × 10 cm) treatment. From the results of the present study indicated that increase in plant density dry matter.

Combined effect of different levels of macronutrient and spacing proved significant differences on dry matter weight of black cumin (Table 7). Results revealed that the highest dry matter weight (12.13 g) was obtained from T<sub>2</sub>S<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> and 20 cm × 15 cm) treatment combination. The lowest dry matter weight (7.64 g) was recorded from T<sub>0</sub>S<sub>1</sub> (control and 20 cm × 10 cm) treatment combination which was statistically similar to T<sub>0</sub>S<sub>2</sub> treatment combination. Rest of the treatment combination performed intermediate results in terms of dry matter weight compared to all other treatments.

**Table 5. Effect of nutrients combination on 1000 seed weight, yield plot<sup>-1</sup>, yield ha<sup>-1</sup> and dry matter of black cumin (*Nigella sativa* L.)**

Treatments	1000 seed weight (g)	Yield plot <sup>-1</sup> (g)	Yield ha <sup>-1</sup> (t)	Dry matter (g)
T <sub>0</sub>	2.80 c	127.14 d	0.88 d	7.90 d
T <sub>1</sub>	2.90 b	144.52 c	1.00 c	8.93 c
T <sub>2</sub>	2.99 a	170.40 a	1.18 a	11.29 a
T <sub>3</sub>	2.89 b	154.30 b	1.07 b	10.70 b
<b>LSD(0.05)</b>	0.0347	6.2480	0.0434	0.2528
<b>CV%</b>	0.91	4.29	4.29	2.66

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup>

**Table 6. Effect of plant spacing on 1000 seed weight, yield plot<sup>-1</sup>, yield ha<sup>-1</sup>, and dry matter of black cumin (*Nigella sativa* L.)**

Treatments	1000 seed weight (g)	Yield plot <sup>-1</sup> (g)	Yield ha <sup>-1</sup> (t)	Dry matter (g)
S <sub>1</sub>	2.82 c	156.66 a	1.09 a	9.30 c
S <sub>2</sub>	2.91 b	150.57 b	1.05 b	10.03 a
S <sub>3</sub>	2.96 a	140.04 c	0.97 c	9.79 b
<b>LSD(0.05)</b>	0.0272	5.4109	0.0376	0.2189
<b>CV%</b>	0.91	4.29	4.29	2.66

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: S<sub>1</sub> = 20 cm × 10 cm, S<sub>2</sub> = 20 cm × 15 cm, S<sub>3</sub> = 20 cm × 20 cm, NS= Not significant

**Table 7. Combined effect nutrients combination and plant spacing on 1000 seed weight, yield plant<sup>-1</sup>, yield plot<sup>-1</sup>, yield ha<sup>-1</sup>, and dry matter of black cumin (*Nigella sativa* L.)**

Treatments	1000 seed weight (g)	Yield per plant (g)	Yield per plot (g)	Yield per ha (t)	Dry matter (g)
T <sub>0</sub> S <sub>1</sub>	2.75 h	1.87 j	134.40 fg	0.93 fg	7.64 h
T <sub>0</sub> S <sub>2</sub>	2.81 fgh	2.38 gh	128.34 gh	0.89 gh	7.97 gh
T <sub>0</sub> S <sub>3</sub>	2.85 efg	3.30 d	118.68 h	0.82 h	8.08 g
T <sub>1</sub> S <sub>1</sub>	2.79 gh	2.12 i	152.40 bcd	1.06 bcd	8.553 f
T <sub>1</sub> S <sub>2</sub>	2.93 cd	2.66 ef	143.64 def	0.99 def	9.25 e
T <sub>1</sub> S <sub>3</sub>	2.98 bc	3.82 c	137.52 efg	0.96 efg	8.99 e
T <sub>2</sub> S <sub>1</sub>	2.90 cde	2.48 fg	178.32 a	1.24 a	10.14 d
T <sub>2</sub> S <sub>2</sub>	3.02 ab	3.27 d	176.76 a	1.23 a	12.13 a
T <sub>2</sub> S <sub>3</sub>	3.06 a	4.34 a	156.12 bc	1.08 bc	11.61 b
T <sub>3</sub> S <sub>1</sub>	2.84 efg	2.24 hi	161.52 b	1.12 b	10.85 c
T <sub>3</sub> S <sub>2</sub>	2.88 def	2.84 e	153.54 bcd	1.07 bcd	10.76 c
T <sub>3</sub> S <sub>3</sub>	2.96 bc	4.11 b	147.84 cde	1.03 cde	10.50cd
<b>LSD(0.05)</b>	0.0786	0.2097	10.822	0.0752	0.4379
<b>CV%</b>	0.91	4.20	4.29	4.29	2.66

In a column means having similar letter (s) are statistically similar and those having dissimilar letter (s) differ significantly by LSD at 0.05 levels of probability. Note: T<sub>0</sub> = N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> kg ha<sup>-1</sup> (control), T<sub>1</sub> = N<sub>45</sub>P<sub>25</sub>K<sub>20</sub> kg ha<sup>-1</sup>, T<sub>2</sub> = N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>, T<sub>3</sub> = N<sub>135</sub>P<sub>75</sub>K<sub>60</sub> kg ha<sup>-1</sup> and S<sub>1</sub> = 20 cm x 10 cm, S<sub>2</sub> = 20 cm x 15 cm, S<sub>3</sub> = 20 cm x 20 cm. NS= Not significant

### Conclusion:

Considering the above result of this experiment; the following conclusion and recommendations can be drawn:

- In the experiment T<sub>2</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup>) treatment was more effective than without nutrient T<sub>0</sub> (control)
- The spacing S<sub>3</sub> (20 cm x 20 cm) gave higher seed yield per plant but the spacing S<sub>1</sub> (20 cm x 10 cm) gave maximum yield per hectare
- During the investigation, the best treatment combination was obtained from T<sub>2</sub>S<sub>1</sub> (N<sub>90</sub>P<sub>50</sub>K<sub>40</sub> kg ha<sup>-1</sup> with 20 cm x 10 cm ) having yield potentiality of 1.24 t/ha

### References

1. Shewaye L. (2011). Antifungal substances from essential oils. MSc Thesis, Addis Ababa University, Ethiopia.
2. Abadi, B.H.M., Ganjali H.R. and Mobasser H.R. (2015). Effect of mycorrhiza and phosphorous fertilizer on some characteristics of black cumin. *Biological Forum*.7(1): 1115-1120.
3. Simon, J.E., Chadwick A.F. and Crack L.E. (1984). Herbs: An indexed bibliography. 1971- 1980. The scientific literature on selected herbs, and aromatic and medicinal plants of the temperate zone. Arcon Book press, Hamden CT, p. 770.
4. Ustun, G., Kent, L., Cekin, N. and Civelekoglu, H. (1990). Investigation of the technological properties of *Nigella sativa* L. *Seed Oil*. *JAOCS*, **67**(12): 71- 86,
5. BBS (Bangladesh Bureau of Statistics). (2018). Agricultural Statistics Yearbook-2018.
6. Troug, E. (1973). Mineral nutrition in relation to autogeneity of plants. In: Nutrition of plants. Oxford and IBH publishers, New Delhi, pp. 345.
7. Barber, S.A. (1984). Soil nutrient bioavailability: A mechanistic approach. John Wiley & Sons Inc., New York, NY. p. 398.

8. Champawat R.S. and Pathak, V.N. (1982). Role of nitrogen, phosphorous and potassium fertilizers and organic amendments in cumin (*Cuminum cyminum* L.) with incites by *Fusarium oxysporum* fsp. *cumin* *Indian J. Agric. Sci.* **58**(9): 728-730.
9. Miah, M.H.N., Karim, M.A., Rahman, M.S. and Islam, M.S. (1990). Performance of Nizersail mutants under different spacing. *Bangladesh J. Train. and Dev.* **3**(2): 31-34.
10. Shaalan, M.N. (2005). Influence of bio-fertilization and chicken manure on growth, yield and seeds quality of *Nigella sativa*, L. plants. *Egypt. J. Agric. Res.* **83**(2): 811-828.
11. Tuncturk, R., Tuncturk, M. and Ciftci, V. (2012). The effects of varying nitrogen doses on yield and some yield components of black cumin (*Nigella Sativa* L.). Yuzuncu Yil University, Faculty of Agriculture, Department of Field Crops, Van, Turkey, *Advances in Environmental Biology*, **6**(2): 855-858.
12. Gamal, T.M., Mohamed M.G., Gamal, A.A. and Sabah A.M. (2012). Comparative study on black cumin (*Nigella sativa*, l.) plants, grown under different plant spacing and fertilization treatments. *Assiut J. Agric. Sci.* **43**(6): 56-70.
13. Mengistu, F.G., Wegayehu, G., Ali, D.F.A., & Fufa, D.T.N. (2021). The influence of seed rate and inter-row spacing on seed yield and yield attributes of black cumin in arsi highlands, *Ethiopia. J. Biol. Agric. and Healthcare*. **11**(1): 254-263.
14. Toncer, O. and Kizil, S. (2004). Effect of seed rate on agronomic and technologies characters of *Nigella sativa* L. *Int. J. Agric. and Bio.* **6**(3): 529-532.
15. Tuncturk, M., Ekin, Z. and Turkozu, D. (2005). Response of black cumin (*Nigella sativa* L.) to different seed rates growth, yield components and essential oil content. *J. Agron.* **4**(3): 216-219.
16. Koli, S.A. (2013). Effect of variety and plant spacing on seed yield and yield attributes of black cumin (*Nigella sativa* L.). MS Thesis, Sher-e-Bangla Agricultural University, Dhaka.
17. Roussis, I., Travlos, I., Bilalis, D. and Kakabouki, I. (2017). Influence of Seed rates and Fertilization on Yield and Yield Components of *Nigella sativa* L. Cultivated under Mediterranean Semi-Arid conditions. *AgroLife Scientific J.* **6**(1): 218-223.
18. Fekadu G.M., Gizaw W., Demis F., Ali, A., Tsagaye, D. and Fufa, N. (2021). Influence of seed rate and inter-row spacing on seed yield and yield attributes of black cumin. *J. Biol, Agric. and Healthcare*. **11**: 215-221.
19. Kandeel, Y.R., Nofal, E.A., Menesi, F.A., Reda, K.A., Taher M.N. and Zaki Z.T. (2001): Effect of some cultural practices on growth and chemical composition of *Foeniculum vulgare*, Mill. Proc. 5 th Arabian Hort. Conf., March 24-28, Ismailia, Egypt, 61-72.
20. Ozguven, M, and Sekeroglu, N. (2007). Agricultural practices for high yield and quality of black cumin (*Nigella sativa* L.) cultivated in Turkey. *Acta Hort.* **756**: 329-337.
21. Giridhar, K., Reddy, G. S., Kumari, S. S., Kumari, A. L., & Sivasankar, A. (2017). Influence of sowing window and plant density on growth, phenology, yield and quality of *Nigella sativa* L. in coastal humid tropic. *Int. J. Curr. Microbiol. App. Sci.* **6**(9): 499-512.
22. Rana, S., Singh, P.P., Naruka, I.S. & Rathore, S.S. (2012). Effect of nitrogen and phosphorus on growth yield and quality of black cumin (*Nigella sativa* L.). *Int. J. Seed Spic.* **2**(2): 5-8.
23. Kaheni A., Ramazani, S.H.R., Ganjali, H.R. and Mobaser, H.R. (2013). Effect of nitrogen fertilizer and plant density on yield and its components in Cumin (*Cuminum cyminum* L.) in South Khorasan Province. *Int. J. Agric. Crop Sci.* **6**(5): 248-251.
24. Datta S. (2004). Black Cumin: Potentials and constraints in indian agriculture: A review. Cooch Behar, West Bangal, India, p. 34
25. Weiss E.A. (2002). Spice Crops. CABI Publishing, London, UK, 411.
26. Kizil, Suleyman, & Toncer, O. (2005). Effect of raw spacing on seed yield, yield components, fatty oil and essential oil of *Nigella sativa* L. *Crop Res.* **30**(1): 107 -112.
27. Valadabadi S.A. and Aliabadi, H.F. (2011). Investigation of bio-fertilizers influence on quantity and quality characteristics in *Nigella sativa* L. *J. Hort. and Forestry*, **3**(3): 88-92.
28. Ali, M.M.K., Hasan, M.A. and Islam, M.R. (2015). Influence of fertilizer levels on the growth and yield of black cumin (*Nigella sativa* L.). *The Agriculturists.* **13**(2): 97-104.