

## Original Research Article

### Effect of *H. Pylori* on fibrinogen level among Sudanese patients at Khartoum state

Formatted: Font: Italic

#### ABSTRACT

**Background:** *Helicobacter pylori* (*H. pylori*) is a gram-negative, spiral shaped pathogenic bacterium that specifically colonizes the gastric epithelium causing chronic gastritis, peptic ulcer disease, and/or gastric malignancy.

**Aim:** To assess the effect of *H. pylori* on fibrinogen level of Sudanese patients with diabetes ~~millets-mellitus~~ (D.M) living in Khartoum state.

**Subjects and Methods:** ~~This was ease~~ It is a case control study conducted among Sudanese diabetic patients during the period from 2018 to 2021. A total of 181 diabetics patients are involved in this study. Citrated blood and stool samples were collected from each participant. All stool samples were tested for the presence of *H. pylori* antigen by using commercially available *H.pylori* antigen detection card. Based on ~~the result~~ the result examination for *H.pylori* the study population are sub grouped in two groups one group consisted of 124 of cases with Positive *H. pylori* antigen and another group had 57 members of controls with negative *H. pylori* infection. Both groups were assessed for the fibrinogen level. Data was gathered and analyzed by using SPSS version 20.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

**Results:** Our results revealed that positive cases have slightly higher values of fibrinogen levels 326.29± 89.99 mg/dl compared to 309.47± 90.82 mg/dl for control group. However, the statistical analysis indicated d that the difference was remain with *P*-value of 0.246. Among the positive cases group ~~gender concerned~~ the fibrinogen mean concentration ~~mean~~ in the plasma of males s 336.2±98.5 mg/dl was higher than that in females s -311.1±73.4 mg/dl.

Comment [S1]: What do you mean by remain?

Formatted: Font: Italic

**Conclusion:** They were insignificant association between fibrinogen level and presence of *H.pylori* infection among Sudanese diabetic patients. Ggender concerned the fibrinogen concentration in the plasma in *H. pylori* among males showed higher levels than females.

## Key words

Diabetes mellitus, *H. pylori* infection, Fibrinogen level

Formatted: Font: Italic

## Introduction:

*Helicobacter pylori* is a gram-negative bacterium that colonizes the stomach and causes persistent infection. The infection is typically acquired in the first few years of life<sup>(1,2)</sup>. The associated risk factors of *H. pylori* infection include living in crowded households, low socioeconomic conditions and infected family members<sup>(3,4)</sup>. The infection is common worldwide with highest prevalence rates reaching 80–90% in developing countries and underprivileged communities, while a much lower prevalence of 20–50% is recorded in developed countries<sup>(5,6,7)</sup>.

Formatted: Font: Italic

Formatted: Font: Italic

Associations of *H. pylori* infection with DM incidence have been reported. Recent meta-analyses showed a significant 1.7 to 2-fold higher prevalence of *H. pylori* infection in persons with T2DM vs. non-diabetic individuals<sup>(8)</sup>. In some of the studies that reported a positive association between *H. pylori* infection and DM<sup>(9)</sup>, the association became on-statistically significant after adjustment for potential confounders such as age and socioeconomic status. Other studies reported no significant association between *H. pylori* and DM,<sup>(10)</sup> or a significant association only in persons with BMI>25.<sup>(11)</sup> Several studies did not control adequately for socioeconomic status and for traditional risk factors of DM, such as obesity and physical inactivity<sup>(12)</sup>.

Comment [S2]: Make it italic all over the manuscript.

Furthermore, most of the evidence is based on small-scale hospital-based case-control studies, in which the source population, selection of control population and representativeness of the sample were not fully described.<sup>(13)</sup> For these reasons, inference and generalizability of findings from such studies should be done with caution. On the other hand, recent well-designed studies show convincing evidence of the potential involvement of *H. pylori* infection in the occurrence of DM, and possibly in IGT.<sup>(14)</sup>

Diabetic patients have higher cardiovascular morbidity than non-diabetic subjects. Several studies have shown that homeostatic factor especially hyperfibrinogenemia is implicated as a source of atherosclerosis and its complications. Studies have reported that fibrinogen levels were higher in diabetics than in controls. Very few studies have been done regarding the association of *H. pylori* with fibrinogen in type 2 diabetes mellitus<sup>(15)</sup>.

~~In~~ [This](#) review aimed to evaluate the possible relationship between *H. pylori* and T2DM according to epidemiological surveys of 70 studies retrieved from databases, including Scopus, PubMed, and Google Scholar about the relationship between *H. pylori* and T2DM, and discuss the reported background mechanisms of this correlation. According to the results of their study, the different studies have shown that *H. pylori* is more prevalent in Type 2 diabetic patients than healthy individuals or nondiabetic patients. The reason is development of *H. pylori* infection-induced inflammation and production of inflammatory cytokines as well as different hormonal imbalance by this bacterium, which are associated with diabetes mellitus. On the other hand, by tracing anti-*H. pylori* antibodies in patients with diabetes mellitus and occurrence of symptoms such as digestive problems in >75% of these patients, it can be concluded that there is a relationship between this bacterium and T2DM. Considering the evidence, it is crucially important that the probability of infection with *H. pylori* is evaluated in patients with T2DM so that medical process of the patient is followed with higher cautious.<sup>(16)</sup>

#### **Material and Method:**

##### **Study setting and population:**

This ~~was~~ case control study [was](#) conducted among Sudanese diabetic patients during the period from 2018 to 2021. A total of 181 diabetics patients are involved in this study. Citrated blood, stool samples were collected from each participant. All stool samples were tested for the presence of *H. pylori* antigen by using commercially available *H.pylor* antigen detection card. Based on the result examination for *H.pylori* the study population are sub grouped in two groups one group 124 of cases with Positive *H.pylori* antigen and another group 57 of controls with negative *H. pylori* infection. Both groups were assessed for the fibrinogen level.

##### **Detection of *H. pylori* antigen in stool sample**

The stool samples were evaluated by the card test according to the manufacturer's protocol. A single red band appearing across the central window in the site marked with the control line was considered negative. A red band appearing in the site marked with the result line and in the site marked with the control line was considered positive. A total absence of the control band, regardless of the appearance of the result site was considered invalid.

### **Estimation of fibrinogen level**

Several steps have been followed according to manufacture protocol to assess the fibrinogen level in the blood samples of cases and controls. The citrated platelet poor plasma (PPP) of control pool plasma was used and the following dilutions in buffer were made: Dilution: 1:5 1:10. Then in plastic plain tube 90ul of buffer + 10ul of control plasma was added to obtain a 1:5 dilution. Then each of a second and a third plastic plain tubes 50 ul of buffer was added. In the second tube 50 ul from the first tube was added to 50 ul buffer to obtain a dilution of 1:10. Alternatively, a calibration curve which was provided with each kit. The same calibration curve could be used when using the same lot of reagents and performing a daily quality control. The PPP samples were diluted 1:10 with Imidazol buffer solution. Then the diluted samples were assayed for fibrinogen assay using the automated coagulometer and the clotting time of each sample was blotted on the log-log paper and the corresponding concentration was gotten from the curve. Reference values : 200 - 400 mg/dl.

### **Quality control**

Pathological and normal control plasma were used to assure the accuracy of the result.

### **Data collection and analysis**

The cases and controls demographic data as well as laboratory test data were obtained and recorded. Data was analyzed by using computer software package for social science (SPSS). Independent T test was used to compare between cases and controls in the level of plasma fibrinogen. The probability value  $<0.05$  was considered to indicate a statistically significant value

### **Ethical consecrations**

All individuals signed informed consent prior to their enrolment in the study. Also, the study was planned according to the ethical guidelines following the Declaration of Ethics Committee of Karary University of Medical Sciences approved it.

## RESULTS

This study ~~was~~ involved a total of 181 participants, about one third of them 58 (32.0% ) are aged 53 – 63 years, while only 6 (3.3%) of them are aged less than 20 years old (Table 1). More than half ~~around~~ -103 (56.9%) of the study participants ~~are-were~~ males while ~~the rests~~ 78 (43.1%) ~~are were~~ females (Table 2). Our results revealed that cases have slightly higher values of fibrinogen levels  $326.29 \pm 89.99$  mg/dl compared to  $309.47 \pm 90.82$  mg/dl for control group. However, the statistical analysis indicate that the difference was remain with P-value of 0.246 (Table 3). Our study results showed that among cases group the mean of male's fibrinogen level  $336.2 \pm 98.5$  mg/dl was lower than females' level  $311.1 \pm 73.4$ . While among control group the mean of male's fibrinogen level  $306.2 \pm 76.1$  mg/dl was lower than females' level  $312.6 \pm 104.3$  mg/dl (Table 4). The current study results showed that there were insignificant association between cases and controls age groups and the level of fibrinogen (Tables 5,6).

## Discussion

In the present study they were insignificant different in fibrinogen level among case and control which found fibrinogen level was 326.2 and 309.4 mg/dl in control group which agreement with Yusuf et al.<sup>(17)</sup> which found the median serum fibrinogen level was 434 mg/dl in *H. Pylori* positive patients and 486 mg/dl in H Pylori negative patients, with no significant difference between the two groups,  $p = 0.78$ . In the study conducted by Longo et al.,<sup>(18)</sup> which found Fibrinogen level was  $471.5 \pm 23.6$  in case and  $297.2 \pm 14.7$  in control and the P.value was (0.0001).

H. pylori infection could be linked to the early stages of coronary atherosclerosis rather than advanced coronary atherosclerosis.<sup>(19)</sup> Stroke is commoner than coronary heart disease in Africans with severe and uncontrolled hypertension. Progression of carotid lesions already associated with H. pylori infection may explain the onset of carotid plaques and stroke<sup>(20)</sup> in this screening population. However, carotid atherosclerosis is not related to H. pylori infection in the United Kingdom,<sup>(21)</sup> whereas HP-seropositivity is associated with carotid plaques and cerebrovascular and cardiovascular events in Italy.<sup>(21)</sup> Our data clearly demonstrate that severity

Formatted: Font: Italic

of *H. pylori* infection was significantly associated with the most important traditional risk factors for CVD, such as diabetes mellitus, arterial hypertension, high levels of serum fibrinogen and total cholesterol, and low HDL-cholesterol. In these HP seropositive, male sex was the only independent predictor of both cerebrovascular and coronary heart diseases. *H. pylori* infection showed an extremely broad spectrum of disease outcomes. These findings lend support to the notion that chronic *H. pylori* infection with exacerbation of inflammation (elevated fibrinogen), male sex (smoking) and dyslipidemia may contribute to early onset of atherosclerosis<sup>(22,23)</sup> in those Africans facing demographic transition. Severity of Seropositivity in this infected group may reflect virulent strains bearing the cytotoxin-associated protein (CagA), with changes in inflammatory markers and higher risk of myocardial infarction.<sup>(24,25)</sup>

As in previous studies, we did not observe any influence of *H. pylori* status on the white blood cell or platelet counts,<sup>(26,27)</sup> but the mean plasma fibrinogen level was reduced by approximately 5% after *H. pylori* eradication therapy. By multivariate analysis, eradication treatment, regardless of its effect on *H. pylori* infection, was the only independent predictor of a reduction in the fibrinogen level. Conceivably, this is due to the effect of antibiotic treatment on the total pathogen burden or to the intrinsic anti-inflammatory properties of macrolide antibiotics.<sup>(28)</sup> Previous studies in patients with ischaemic heart disease have shown similar decreases in plasma fibrinogen after treatment of *H. pylori* infection,<sup>(29)</sup> although there is also evidence against any effect of *H. pylori* eradication treatment on fibrinogen.<sup>(27)</sup>

## Conclusion

From our findings we concluded that, there was insignificant association between fibrinogen level and *H. pylori* infection. Also, there was insignificant association between fibrinogen level and gender among [positive](#) cases group. Further studies on the existing modalities which lower fibrinogen and finding newer treatment measures that can lower the fibrinogen levels without adverse effects, should be done.

## References

1. Sipponen P, Maaroos HI. Chronic gastritis. *Scand J Gastroenterol*. 2015;50(6):657-667. doi:10.3109/00365521.2015.1019918
2. Mera RM, Bravo LE, Camargo MC, et al. Dynamics of *Helicobacter pylori* infection as a determinant of progression of gastric precancerous lesions: 16-year follow-up of an eradication trial. *Gut*. 2018;67(7):1239-1246.
3. Sherman PM, Lin FY. Extradigestive manifestation of *Helicobacter pylori* infection in children and adolescents. *Can J Gastroenterol*. 2005;19(7):421-424. doi:10.1155/2005/971974
4. Gasbarrini G, Racco S, Franceschi F, et al. Manifestazioni extragastriche dell'infezione da *Helicobacter pylori* [*Helicobacter pylori* infection: from gastric to systemic disease]. *Recenti Prog Med*. 2010;101(1):27-33.
5. Mou WL, Feng MY, Hu LH. Eradication of *Helicobacter Pylori* Infections and GERD: A systematic review and meta-analysis. *Turk J Gastroenterol*. 2020;31(12):853-859. doi:10.5152/tjg.2020.19699.
6. White JR, Winter JA, Robinson K. Differential inflammatory response to *Helicobacter pylori* infection: etiology and clinical outcomes. *J Inflamm Res*. 2015;8:137-147. Published 2015 Aug 13. doi:10.2147/JIR.S64888
7. Chen XZ, Schöttker B, Castro FA, et al. Association of *helicobacter pylori* infection and chronic atrophic gastritis with risk of colonic, pancreatic and gastric cancer: A ten-year follow-up of the ESTHER cohort study. *Oncotarget*. 2016;7(13):17182-17193. doi:10.18632/oncotarget.7946
8. Gao C. Molecular pathological epidemiology in *Helicobacter pylori* infection and risk of chronic atrophic gastritis. *Journal of Gastroenterology and Hepatology Research*. 2017; 6(3):2354-2357.
9. Epplein M, Zheng W, Xiang YB, et al. Prospective study of *Helicobacter pylori* biomarkers for gastric cancer risk among Chinese men. *Cancer Epidemiol Biomarkers Prev*. 2012;21(12):2185-2192. doi:10.1158/1055-9965.EPI-12-0792-T
10. Epplein M, Zheng W, Li H, et al. Diet, *Helicobacter pylori* strain-specific infection, and gastric cancer risk among Chinese men. *Nutr Cancer*. 2014;66(4):550-557. doi:10.1080/01635581.2014.894096

11. Franceschi F, Gasbarrini A, Polyzos SA, Kountouras J. Extragastric Diseases and Helicobacter pylori. *Helicobacter*. 2015;20 Suppl 1:40-46. doi:10.1111/hel.12256.
12. Lai CY, Yang TY, Lin CL, Kao CH. Helicobacter pylori infection and the risk of acute coronary syndrome: a nationwide retrospective cohort study. *Eur J Clin Microbiol Infect Dis*. 2015;34(1):69-74. doi:10.1007/s10096-014-2207-7.
13. Liu J, Wang F, Shi SL. Helicobacter pylori infection increase the risk of myocardialinfarction: a meta-analysis of 26 studies involving more than 20,000 participants. *Helicobacter* 2015; 20(3): 176–83.
14. Shmueli H, Wattad M, Solodky A, Yahav J, Samra Z, Zafrir N. Association of Helicobacter pylori with coronary artery disease and myocardial infarction assessed by myocardial perfusion imaging. *Isr Med Assoc J* 2014; 16(6): 341–6.
15. Xu Y, Wang Q, Liu Y, Cui R, Zhao Y. Is Helicobacter pylori infection a critical risk factor for vascular dementia?. *Int J Neurosci*. 2016;126(10):899-903. doi:10.3109/00207454.2015.1081387.
16. Hosseinasab Nodoushan SA, Nabavi A. The Interaction of *Helicobacter pylori* Infection and Type 2 Diabetes Mellitus. *Adv Biomed Res*. 2019;8:15. Published 2019 Feb 27. doi:10.4103/abr.abr\_37\_18.
17. Yusuf SW, Mishra RM. Effect of Helicobacter pylori infection on fibrinogen level in elderly patients with ischaemic heart disease. *Acta Cardiol*. 2002;57(5):317-322. doi:10.2143/AC.57.5.2005446
18. Longo-Mbenza B, Nsenga JN, Mokondjimobe E, et al. Helicobacter pylori infection is identified as a cardiovascular risk factor in Central Africans. *Vasc Health Risk Manag*. 2012;6:455-461. doi:10.2147/VHRM.S28680
19. Park MJ, Choi SH, Kim D, et al. Association between Helicobacter pylori Seropositivity and the Coronary Artery Calcium Score in a Screening Population. *Gut Liver*. 2011;5(3):321-327. doi:10.5009/gnl.2011.5.3.321
20. Corrado E, Rizzo M, Tantillo R, et al. Markers of inflammation and infection influence the outcome of patients with baseline asymptomatic carotid lesions: a 5-year follow-up study. *Stroke*. 2006;37(2):482-486. doi:10.1161/01.STR.0000198813.56398.14



21. Mayr M, Kiechl S, Tsimikas S, et al. Oxidized low-density lipoprotein autoantibodies, chronic infections, and carotid atherosclerosis in a population-based study. *J Am Coll Cardiol*. 2006;47(12):2436–2443.
22. Spence JD, Norris J. Infection, inflammation, and atherosclerosis. *Stroke*. 2003;34:333–334.
23. Marra M, Bonfigli AR, Bonazzi P, et al. Asymptomatic *Helicobacter pylori* infection increases asymmetric dimethylarginine levels in healthy subjects. *Helicobacter*. 2005;10(6):609–614.
24. Elkind MS, Luna JM, Moon YP, et al. Infectious burden and carotid plaque thickness: the northern Manhattan study. *Stroke*. 2010;41(3): e117–e122.
25. Suzuki H, Matsuzaki J, Hibi T. Lifestyle-related diseases and *H. pylori*. *Nihon Rinsho*. 2009;67(12):2366–2371.
26. Parente F, Imbesi V, Cucino C, et al. *Helicobacter pylori* CagA seropositivity does not influence inflammatory parameters, lipid concentrations and hemostatic factors in healthy individuals. *J Intern Med* 2000; 247: 213–7.
27. Schweeger I, Fitscha P, Sinzinger P. Successful eradication of *Helicobacter pylori* as determined by <sup>13</sup>C-urea breath test does not alter fibrinogen and acute phase response markers. *Thromb Res* 2000; 97: 411–20.
28. Ianaro A, Ialenti A, Maffia P, et al. Anti-inflammatory activity of macrolide antibiotics. *J Pharmacol Exp Ther* 2000; 292: 156–63.
29. Koenig W. Fibrin(ogen) in cardiovascular disease: an update. *Thromb Haemost*. 2003;89(4):601-609.

**Table (1): Distributions of age group among study of case populations:**

Age group	Frequency	Percent
less than 20 years	6	3.3
20-30	10	5.5
31-41	22	12.2
42-52	47	26.0
53-63	58	32.0
64-74	31	17.1
75-85	7	3.9
Total	181	100.0

**Table (2): Distributions of gender among study of case populations:**

Gender	Frequency	Percent
Female	78	43.1
Male	103	56.9
Total	181	100.0

**Table (3): the mean and Std of Glucose and fibrinogen among study of case and control populations:**

Parameter	Case (N=124) (mean $\pm$ Std)	Control (N=57) (mean $\pm$ Std)	P. value
Fibrinogen	326.29 $\pm$ 89.99	309.47 $\pm$ 90.82	0.246

The table shows the mean  $\pm$  SD (mini - max) and probability (P). Independent T-test was used for comparison. P value  $\leq$  0.05 was considered significant.

**Table (4): Cases and controls fibrinogen level according to their gender**

<b>Participants</b>	<b>Gender</b>	<b>Fibrinogen level mg/dl</b>
Case	Female	311.1±73.4
	Male	336.2±98.5
Control	Female	312.6 ±104.3
	Male	306.2±76.1

**Table (5): association between age group and fibrinogen level among H. pylori positive result**

<b>Age group</b>	<b>Fibrinogen</b>
less than 20 years	390.0±36.3
20-30	375.4± 88.9
31-41	329.8±119.5
42-52	332.9±76.9
53-63	321.6±87.5
64-74	293.9± 96.1
75-85	326.2±89.9

**Table (6): association between age group and, fibrinogen level among H. pylori negative result**

<b>Age group</b>	<b>Fibrinogen</b>
<b>less than 20 years</b>	231.0±21.0
<b>20-30</b>	356.0±80.2
<b>31-41</b>	354.8±109.1
<b>42-52</b>	280.1±101.0
<b>53-63</b>	327.8±80.6
<b>64-74</b>	298.4±100.4
<b>75-85</b>	241.0±72.3