

## Implication of Nephropathy in Sickle Cell Anaemia

### ABSTRACT

**Aim:** This study sought to investigate the effect of sickle cell anemia on kidney using endogenous biomarkers.

**Methodology:** This study is a case-control study which enrolled 60 sickle cell anemia patients (30 males and 30 females) in a stable state at the Obafemi Awolowo University Teaching Hospital in Ile-Ife, Osun State, Nigeria, as well as 60 healthy controls (30 males and 30 females). Each participant had five millilitres (5 mL) of venous blood taken and dispensed into a lithium heparin sample vial. Renal biomarkers were assessed using established procedures utilizing the blood sample. Urinalysis was also conducted for 41 sickle cell patients and 11 control subjects.

**Results:** According to the findings of this investigative study, renal indices, except uric acid were lower in sickle cell anemia patients than in control subjects and are related to glomerular hyperfiltration in sickle cell nephropathy. It is noteworthy that the effect of sickle cell anemia on renal parameters was stronger in male patients than in females. Furthermore, the identical plasma creatinine levels could be attributed to the sickle cell participants' similarities in muscle mass. Protein was found in the urine of 21.95% of sickle cell patients.

**Conclusion:** The results of this study confirmed that kidney abnormality could be implicated in sickle anemia.

**Keywords:** *Nephropathy; sickle cell anemia; steady state; proteinuria*

### 1. INTRODUCTION

Sickle cell anemia is a hereditary disorder in which the oxygen-carrying protein "hemoglobin" becomes defective. This

defect makes the red blood cells sickle when exposed to a relatively deoxygenated state inside the vasculature [1] As a result of this single action, sickle cell anemia is

portrayed as a life-threatening illness and one of the most common hereditary diseases worldwide [2,3]. Sickle cell disease is more prevalent in Sub-Saharan Africa [1]. This is probably due to the rate of malaria transmission among the people in these areas. It was proposed that hemoglobinopathies, especially the mutated gene that induces sickle cells may protect against malaria's severe life-threatening effects [3-5]. Sickle cell anemia affects multiple organs and has acute and chronic consequences [6]. An example of these multiple-organ systems in the body is the kidney. The kidneys are a pair of bean-shaped organs present in all vertebrates [7]. The kidneys play a unique role in the elimination of wastes (majorly creatinine, uric acid, and urea) and extra fluids (electrolyte/acid-base balances) from the body, and deterioration in its functions is known as nephropathy [3]. Sickle cell nephropathy (SCN) is prevalent in sickle cell disease and is characterized by medullary hypoperfusion, renal vasoconstriction, and kidney and/or cortical hyperperfusion [8]. The hallmark of this disease is the presence of sickled red blood cells in the renal medullary vessels which leads to a disturbance in tubular function, glomerular filtration, blood pressure regulation, and water and electrolyte metabolism [6]. The aftermath result of sickle cell nephropathy may be a progressive end-stage kidney

disease with an elevated risk of morbidity and mortality [7,9] if it is not quickly detected and properly managed. Plasma creatinine, urea, and uric acids (to a lesser extent) are part of the commonest and generally recognized parameters often used in the assessment of GFR-kidney functions and as such, their elevated level is an indicator of impaired renal function [10-13]. In this study, we evaluated the impact of sickle cell anemia on the kidney using endogenous markers such as creatinine, uric acid, and urea as well as urine parameters.

## **2. METHODOLOGY**

### **2.1 Research Design**

This study carried out in the Obafemi Awolowo University Teaching Hospital in Osun State, Nigeria, is a cross sectional research on adult SCA patients in the steady state. The study was authorized by the Hospital's Ethical and Scientific Committee. Adult Nigerians of both genders who volunteered and given informed written consent were used as research subjects. Patients who came to the hematology clinic were enrolled in the order in which they arrived. People without sickle cell anemia served as the control group.

The study included 60 SCA patients (30 males and 30 females) and 60 control volunteers (30 males and 30 females). Each participant had five millilitres (5 mL) of

venous blood collected and distributed into a lithium heparin sample container for the measurement of liver parameters.

## 2.2 Inclusion criteria for Patients

SCA patients 16 years of age and older in steady state (a period of stable clinical condition occurring at least one week before or three weeks after a VOC or three months after a haemolytic crisis requiring a blood transfusion).

## 2.3 Inclusion Criteria for Control Subjects

Healthy individuals with Hb A from the Ile-Ife community who were 16 years of age and older.

## 2.4 Exclusion Criteria for Controls

People who are taking any drugs as well as those who smoke or drink too much alcohol (14 units per week for females and 21 units per week for males) were excluded from the study.

## 2.5 Exclusion Criteria for Patients

- ✓ Any additional medical conditions, such as hypertension or diabetes mellitus.
- ✓ Patients with sickle cell anemia who smoke or drink excessively (14 units per week for females and 21 units per week for males). 12 pint of beer (approx. 300 mL) equals 1 unit of

alcohol (8-10 g): 25 mL distilled spirit  
1 glass sherry, 1 glass wine

- ✓ Patients in crisis or those who have had a blood transfusion within the last three months.

Comment [1]: New topic

## 2.6 Ethical Considerations

Before being enrolled in the study, all individuals gave their informed consent. Before beginning the study, the Ethical and Research Committee of OAUTHC, Ile-Ife, was consulted and ethical approval was granted.

## 2.7 Determination of Renal Indices

Creatinine concentration was determined using Jaffe reaction described by Toora and Rejagopal [14]. Urea and uric acid concentrations were determined using a Randox Commercial Kit based on the methods of Fesus et al. [15].

## 2.8 Urinalysis

Urinalysis was conducted for 41 sickle cell patients and 11 Control subjects using dipsticks (Combi 9)


## 2.9 Statistical Analysis

The mean and standard deviation are used to express the results. One-way Analysis of Variance (ANOVA) and Tukey's test were used to determine the degree of homogeneity among the groups. P values

less than 0.05 were considered statistically significant in all analyses performed with Graph Pad Prism Software Version 8.00.


## 2. RESULTS

The effect of sickle cell disease on the concentrations of creatinine, uric acid and urea in both male and female patients are presented in Figs. 1- 3 respectively.




**Figure 1: Effect of Sickle Cell Disease on the Concentration of Creatinine in Male and Female Patients**

Results are presented as mean  $\pm$  standard deviation with n = 30



**Figure 2: Effect of Sickle Cell Disease on the Concentration of Uric Acid in Male and Female Patients**

Results are presented as mean  $\pm$  standard deviation with n = 30



**Figure 3: Effect of Sickle Cell Disease on the Concentration of Urea in Male and Female Patients**

Results are presented as mean  $\pm$  standard deviation with n = 30

**Table 1: Urinalysis of Sickle Cell Patients and Control Subjects**

Parameter	Sickle Cell Patients (n = 41)		Control Subjects (n = 11)	
	Positive	Negative	Positive	Negative
Protein	9	32	3	8
Nitrite	3	38	0	11

Blood	5	36	0	11
Bilirubin	4	37	0	11
Urobilinogen	12	29	1	10
Leukocytes	26	15	2	9

#### 4. DISCUSSION

There is a range of abnormalities that could arise in the kidney of a sickle cell patient. These abnormalities vary from hematuria, proteinuria, hypertrophy, hypertension, decreased renal blood flow, acidification defect (acidosis), and urinary concentration defect (UCD) [8,16]. Biochemical indices (such as creatinine, uric acid, and urea) are of great importance in the assessment of kidney functions and abnormalities, particularly the estimated glomerular filtration rate and tubular renal function.

Creatinine is a non-toxic metabolic waste product of creatine and creatine phosphate that is produced during muscular activity (muscle wear and tear) and excreted mainly by the kidney in healthy individuals [17]. Unlike urea which is being reabsorbed, creatinine undergoes filtration and secretion. Hence, it is a better indicator of renal function and estimator of glomerular function. Urea on the other hand is a toxic end product of protein metabolism. It is partially reabsorbed by the renal tubules and could be influenced by protein diets [7,9]. Urea and uric acid undergo filtration,

secretion, and reabsorption, therefore, these biomarkers are often used together with other parameters (like creatinine and inulin) to assess renal function. Uric acid is a toxic metabolic waste product of purines. Diets high in purines can influence the level of uric acid in the body [18], thus leading to hyperuricemia. Hyperuricemia can settle as crystals (urate) in the joints and then develop into painful gout. When these crystals also settle in the kidneys, they form kidney stones that lead to kidney diseases. For sickle cell individuals, hyperuricemia is linked with an increase in erythrocyte turnover [18]. Creatinine, urea, and uric acid are all nitrogenous products of protein metabolism, and their elevated levels in the body are called Azotemia [7,19].

The concentrations of plasma creatinine and uric acid were statistically significant ( $p < 0.05$ ) in patients with sickle cell disease when compared to control subjects, as shown in Figs. 1 and 2 respectively. It was speculated that a decline in glomerular filtration rate is associated with a rise in plasma urea, creatinine, and uric acid concentrations [20]. However, in this current

study, we experienced an increase in the concentration of uric acids only.

There was a significant decrease in plasma creatinine concentration in both male and female sickle cell patients compared to healthy individuals with HbA. The decrease in the concentration of plasma creatinine and urea is similar to the previous reports from studies conducted in pediatrics and may be attributed to glomerular hyperfiltration [11,21]. In these studies [11,21], glomerular hyperfiltration is common and has been reported as an early manifestation (one of the pathogenic mechanisms) of renal dysfunction among sickle cell children. Another study by Scheinman [13] explained the changing profile of kidney dysfunction in sickle cell disease with respect to age and showed a higher frequency of hyperfiltration in pediatrics. However, other studies [22-24] have shown the possibility of glomerular hyperfiltration in young adult sickle cell patients. Aygun et al. [25] also claimed that the incidence of hyperfiltration increases with age and tends to decrease in the second decade of life with the progression of sickle cell nephropathy and the development of chronic kidney disease. Glomerular hyperfiltration, which is also known as glomerular hypertrophy is defined as an elevated filtration fraction above the mean GFR of healthy individuals

( $18.7 \pm 3.2\%$ ) and is suggested to be triggered by the secretion of both prostaglandins and nitric oxide, following an ischemia damage [26,27].

Furthermore, sickle cell patients are fond of having smaller stature compared to their healthy hemoglobin (HbA) counterparts [11,21] and the rate of creatinine production in the body daily is relatively constant and related to the body composition (muscle mass, age, and gender) [28]. This may also explain the reasons for the lower concentrations of plasma creatinine in sickle cell subjects despite being adults. Also, creatinine elimination is constant and is usually higher in males compared to females partly due to the difference in body mass [29], and the higher frequency of hyperfiltration in males [30]. However, for our study in sickle cell patients, we observed similar levels of plasma creatinine.

In this investigative study, there were no significant changes in the urea concentrations of sickle cell patients when compared to the healthy individuals with Hb A in the normal control. This is consistent with the reports of Al-Naama et al. [31]. Urea is produced by the urea cycle enzymes present in the liver [32]. A slight decrease in urea levels was observed in this study and may be correlated to liver dysfunction, hyperfiltration, and other influencing factors such as diets and

hormones. Indeed, Uche et al. [33] recently reported that sickle cell anemia contributes to liver abnormality

The result of this present study shows a significant increase in the concentration of plasma uric acid in both male and female sickle cell patients compared to the subjects with healthy Hb A. This result is also in agreement with reports from previous studies conducted in both children and adults with SCD [21,34].

Several studies have reported and suggested SCN among pediatrics (7 months-15 years) and the need for early discovery and intervention. Only a few have reported SCN in young adults. This investigative study was conducted on teenagers and adolescents (16 years and above) to show the impact of sickle cell anemia on the kidney, hence the need for periodic renal assessments and comprehensive care.

The results of the urinalysis of this study are presented in table 1. Protein was only found in the urine of 21.95% of the sickle cell patients. Proteinuria is usually an early manifestation of sickle cell nephropathy [35]. In disease state involving the proximal tubule, there is reduced reabsorption of low molecular weight proteins, and consequently increased spillage of these proteins into the urine [36]. This increase of

the urinary concentration of low molecular weight proteins is a sensitive marker of tubular dysfunction, and therefore, tubulointerstitial damage [35]. This result is similar to the findings of Nsiah et al. [37] who reported the diagnostic implications of urinalysis of subjects from a Ghanaian sickle cell clinic. It is interesting to know that patients with proteinuria also had elevated creatinine and urea levels in their blood which is an indication of nephropathy.

## 5. CONCLUSION

Sickle cell nephropathy (SCN) is prevalent in sickle cell anemia and it is associated with severe clinical complications. Sickle cell nephropathy, if not detected early and managed well, can progress to chronic kidney disease and perhaps increase the risk factors for morbidity and mortality. Biochemical markers such as creatinine, urea, and uric acid are of great clinical significance in the early discovery and monitoring of kidney functions and dysfunction in SCD. Further investigations are needed to ascertain the reason for the difference in urea and uric acid levels in male and female sickle cell patients.

## REFERENCE

1. Alabi OJ, Adegboyega FN, Olawoyin DS, Babatunde OA. Functional Foods: Promising Therapeutics for Nigerian Children with Sickle Cell Diseases.

- Heliyon. 2022. <https://doi.org/10.1016/j.heliyon.2022.e09630>.
2. Kjellander C, Sennström MK, Stiller V, Ågren A. Sick cell anemia causes varied symptoms and high morbidity. Serious prognosis in the most common genetic disease in the world. *Lakartidningen*, 2015;112, DCPM.
  3. Olaniran KO, Eneanya ND, Nigwekar SU, Vela-Parada XF, Achebe MM, Sharma A, Thadhani RI. Sick Cell Nephropathy in the Pediatric Population. *Blood purification*, 2019;47(1-3), 205–213. <https://doi.org/10.1159/000494581>
  4. Taylor SM, Parobek CM, Fairhurst RM. Haemoglobinopathies and the clinical epidemiology of malaria: a systematic review and meta-analysis. *Lancet Infect Dis*. 2012;12(6):457–68.
  5. Taylor SM, Cerami C, Fairhurst RM. Hemoglobinopathies: slicing the Gordian knot of Plasmodium falciparum malaria pathogenesis. *PLoS Pathog*. 2013;9(5):e1003327.
  6. Farooq S, Testai FD. Neurologic Complications of Sick Cell Disease. *Current neurology and neuroscience reports*, 2019;19(4), 17. <https://doi.org/10.1007/s11910-019-0932-0>
  7. Wallace MA. Anatomy and physiology of the kidney. *AORN journal*. 1998;68(5), 800–824. [https://doi.org/10.1016/s0001-2092\(06\)62377-6](https://doi.org/10.1016/s0001-2092(06)62377-6)
  8. Nath KA, Hebbel RP. Sick cell disease: renal manifestations and mechanisms. *Nature reviews. Nephrology*, 2015;11(3), 161–171. <https://doi.org/10.1038/nrneph.2015.8>
  9. Wang H, Ran J, Jiang T, Urea. In: Yang, B., Sands, J. (eds) *Urea Transporters. Subcellular Biochemistry*, vol 73. Springer, Dordrecht. 2014; [https://doi.org/10.1007/978-94-017-9343-8\\_2](https://doi.org/10.1007/978-94-017-9343-8_2)
  10. Airaodion AI, Ogbuagu EO. Effect of *Cyperus esculentus* L. (tiger nut) milk on hepatic and renal indices of Wistar rat. *Asian Journal of Research in Nephrology*. 2020;3(2):10-16.
  11. Olawale OO, Adekanmbi AF, Sonuga AA, Sonuga OO, Akodu SO, Ogundeyi MM. Assessment of Renal Function Status in Steady-State Sick Cell Anaemic Children Using Urine Human Neutrophil Gelatinase-Associated Lipocalin and Albumin:Creatinine Ratio. *Medical principles and practice: international journal of the Kuwait University, Health Science Centre*, 2021;30(6), 557–562. <https://doi.org/10.1159/000518458>
  12. van Veldhuisen DJ, Ruilope LM, Maisel, AS, Damman K. Biomarkers of renal injury and function: diagnostic, prognostic and therapeutic implications in heart failure. *European Heart Journal*,

- 2016;37(33), 2577–2585.  
<https://doi.org/10.1093/eurheartj/ehv588>
13. Scheinman JI. Sick cell disease and the kidney. *Nature clinical practice. Nephrology* 2009;5(2), 78–88.  
<https://doi.org/10.1038/ncpneph1008>
14. Toora BD, Rejagopal G. Measurement of creatinine concentration by Jaffe's Reaction. *Indian Journal of Experimental Biology*. 2002;40(3):352–354.
15. Fesus PD, Pressac M, Braconnier F, Aymard P. Automated determination of urinary Na, K, inorganic phosphate, urea, and uric acid without sample dilution, with the RA-XT. *Clin Chem*. 1983;35:481-483.
16. Abdulrahman IS. The Kidney in Sick Cell Disease: Pathophysiology and Clinical Review. 2004.
17. Airaodion AI, Alabi OJ, Ogbuagu EO, Atiba FA, Olawoyin DS. Nephro and hepatotoxic effect of air-freshener in Wistar rats. *Asian Journal of Research in Nephrology*. 2020;3(2):1-9.
18. Diamond HS, Meisel AD, Holden D. The natural history of urate overproduction in sickle cell anemia. *Annals of internal medicine*, 1979;90(5), 752–757.  
<https://doi.org/10.7326/0003-4819-90-5-752>
19. Tyagi A, Aeddula NR. Azotemia. In *StatPearls*. StatPearls Publishing. 2022.
20. Airaodion AI, Megwas AU, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Nephro- and hepato-toxicity of common household insecticides used in Nigeria. *International Research Journal of Gastroenterology and Hepatology*. 2020;3(2):21-28.
21. Aloni MN, Ngiyulu RM, Gini-Ehungu JL, Nsibu CN, Ekila MB, Lepira FB, Nseka NM. Renal function in children suffering from sickle cell disease: challenge of early detection in highly resource-scarce settings. *PloS one*, 2014;9(5), e96561.  
<https://doi.org/10.1371/journal.pone.0096561>
22. Helal I, Fick-Brosnahan GM, Reed-Gitomer B, Schrier RW. Glomerular hyperfiltration: definitions, mechanisms and clinical implications. *Nature reviews. Nephrology*, 2012;8(5), 293–300.  
<https://doi.org/10.1038/nrneph.2012.19>
23. Schmitt F, Martinez F, Brillet G, Giatras I, Choukroun G, Girot R, Bachir D, Galacteros F, Lacour B, Grünfeld JP. Early glomerular dysfunction in patients with sickle cell anemia. *American journal of kidney diseases : the official journal of the National Kidney Foundation*, 1998;32(2), 208–214.  
<https://doi.org/10.1053/ajkd.1998.v32.p009708603>
24. Haymann JP, Stankovic K, Levy P, Avellino V, Tharaux PL, Letavernier E, Grateau G, Baud L, Girot R, Lionnet F. Glomerular hyperfiltration in adult sickle cell anemia: a frequent hemolysis associated feature. *Clinical journal of the American Society of Nephrology*. 2010;5(5), 756–761.  
<https://doi.org/10.2215/CJN.08511109>

25. Aygun B, Mortier NA, Smeltzer MP, Hankins JS, Ware RE. Glomerular hyperfiltration and albuminuria in children with sickle cell anemia. *Pediatric nephrology (Berlin, Germany)*, 2011;26(8), 1285–1290. <https://doi.org/10.1007/s00467-011-1857-2>
26. Becker AM (2011) Sickle cell nephropathy: challenging the conventional wisdom. *Pediatric Nephrol* 26:2099–2109
27. Ataga KI, Orringer EP. Renal abnormalities in sickle cell disease. *American journal of hematology*, 2000;63(4), 205–211. [https://doi.org/10.1002/\(sici\)1096-8652\(200004\)63:4<205::aid-ajh8>3.0.co;2-8](https://doi.org/10.1002/(sici)1096-8652(200004)63:4<205::aid-ajh8>3.0.co;2-8)
28. Airaodion AI, Ene AC, Ogbuagu EO, Okoroukwu VN, Ekenjoku JA, Ogbuagu U. Biochemical changes associated with consumption (by rats) of “garri” processed by traditional and instant mechanical methods. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2019;2(4): 1-11.
29. Nitsch D. Is there a difference in metabolic burden between men and women?. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association - European Renal Association*, 2014;29(6), 1110–1112. <https://doi.org/10.1093/ndt/gft518>
30. Saraf SL, Shah BN, Zhang X, Han J, Tayo BO, Abbasi T, Ostrower A, Guzman E, Molokie RE, Gowhari M, Hassan J, Jain S, Cooper RS, Machado RF, Lash JP, Gordeuk VR. APOL1,  $\alpha$ -thalassemia, and BCL11A variants as a genetic risk profile for progression of chronic kidney disease in sickle cell anemia. *Haematologica*, 2017;102(1), e1–e6. <https://doi.org/10.3324/haematol.2016.154153>
31. Al-Naama LM, al-Sadoon, EA, al-Sadoon TA. Levels of uric acid, urea and creatinine in Iraqi children with sickle cell disease. *JPMA. The Journal of the Pakistan Medical Association*, 2000;50(3), 98–102.
32. Ogbuagu EO, Airaodion AI, Ogbuagu U, Nweke IN, Unekwe PC. Nephrotoxicity of ethanol extract of *Xylopiya aethiopica* fruit in Wistar rats. *International Journal of Advances in Nephrology Research*. 2021;4(1):1-16.
33. Uche CL, Isaiah AO, Ezirim EO, Airaodion AI. Sickle cell anemia contributes to liver abnormality. *International Journal of Research and Reports in Hematology*. 2022; 5(2):122-134.
34. Uche CL, Akinola NO. Biochemical Parameters in the Steady State of Sickle Cell Anaemia: Females Seem to Have Advantage. *Journal of Medical and Dental Science Research*. 2017; 4(1): 62-66.

35. D'Amico G, Bazzi C. Pathophysiology of proteinuria. *Kidney International*. 2003;63, 809-825.
36. Marsenic O, Coulooures KG, Wiley JM. Proteinuria in children with sickle cell disease. *Nephrol Dialys Transpl*. 2008;23, 715-720.
37. Nsiah K, Dzogbefia VP, Ansong D, Osei-Akoto A. The Diagnostic Implications of Spot Urinalysis of Subjects from a Ghanaian Sickle Cell Clinic. *Journal of Internal Medicine Research*. 2014;2(2):15-

UNDER PEER REVIEW