

# Management of Mungbean Yellow Mosaic Virus (MYMV) disease using chemical insecticides and bio-pesticides

## Abstract

Field trials ~~was~~ ~~were~~ conducted during 2018-19 and 2019-20 crop seasons to evaluate the efficacy of insecticides and bio-pesticides for the management of Mungbean Yellow Mosaic Virus (MYMV) disease naturally spread by vector whitefly (*Bemisia tabaci*). The mean of two-year data showed ~~that the~~ lowest disease incidence and ~~the~~ least number of whiteflies were recorded in Imidacloprid (17.8% SL) treated plots. Fifteen days after ~~the~~ 1<sup>st</sup> spray and 2<sup>nd</sup> spray of the Imidacloprid (17.8% SL), recorded 17.09% & 18.15% disease incidence, respectively, and also reported least number of whiteflies being 3.10 five days after 1<sup>st</sup> spray and 3.32 five days after 2<sup>nd</sup> spray which consequently resulted in higher yield (9.10 q/ha). In bio-pesticides, *Beauveria bassiana* @0.5% and Neem Oil @ 3ml/lit, were found to be moderately effective in managing the disease incidence and the population of whiteflies, 22.82 and 25.11 per cent disease incidence recorded at 15 days after first spray and 25.32 and 26.73 per cent at 15 days after second spray.

**Keywords:** Imidacloprid, *Beauveria bassiana*, Bio-pesticide, Neem and Mungbean

## Introduction

Mungbean or green gram (*Vigna radiata* L.) is important pulse crop in India. It is cultivated in tropical and sub-tropical regions of the world. In India it is cultivated throughout the year in an area 4.07 mha and its production is 1.90 mt ~~in 0000 year~~ [9]. The highest acreage under the crop and its production are in the States of Rajasthan, MP Maharashtra ~~and~~ followed by Bihar. The major constraints in the production of ~~mungbean~~ ~~Mungbam~~ is biotic stress factors like ~~infection with pathogenic f~~ ~~ungusi~~, bacteria and viruses. Among which virus is causing extensive losses. Mungbean yellow mosaic virus (MYMV) is the major restraint in the production of the crop. The disease is spread in nature through the whitefly vector, *Bemisia tabaci* in a persistent (~~circulative~~) manner, and by inoculation with sap [2]. Nymphs of *Bemisia tabaci* acquires the virus from diseased leaves. Higher incidences occur in regions where the temperature ranges between 31 to 35°C with a relative humidity of around 70 percent [12]. These conditions favour disease development and multiplication of the whitefly vector, *Bemisia tabaci* [2,7,8]. In the present work, in order to restrict the natural spread of MYMV disease through the whitefly vector newer insecticides and biopesticides were evaluated under prevailing agro-climatic conditions of Jharkhand state.

**Comment [pL1]:** Please consider splitting this sentence into several sentences because it's too long and confusing.

**Comment [pL2]:** Please include more relevant references including the role of the insecticides in the control of the viruses and vector insects!!

**Comment [pL3]:** Where references from 1 to 8? Aren't numbered references style should start from the number one?

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

## Material and Method

Field trials were laid out during *Kharif/Automne*, 2018-19 and 2019-20 crop seasons in Randomized Block Design. Nine chemical insecticides and bio-pesticides as treatments were replicated three times. The Mungbean variants were sown in plots of 3×1.5 m<sup>2</sup> with spacing of 30×10cm<sup>2</sup>. For seed priming treatment seeds were pre-soaked in the Pesticide/Bio-agent/Chemical before sowing. The treatments were as follows: 1. Dupont Tm Rekord TM(Acetamiprid) @ 1.5 ml/lit, 2. A-One (Imidacloprid) @ 0.25 ml/lit, 3. Rogor Plus (Dimethoate) @ 1.5 ml/l, 4. Success (Spinosad) @ 0.25%, 5. Maxtara (Thiomethoxam) @ 0.5g/lit, 6. Daman (*Beauveria bassiana*) @ 0.5 %, 7. Vertifire-L (*Verticillium lecani*) @ 0.25ml/l, 8. Neem Baan (Neem Oil) @ 3ml/lit, 9. Control.

**Comment [pL4]:** Please list treatments in a Table including chemical name, commercial name and field application rate!!

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Comment [pL5]:** Based on what?

Spraying of insecticides was undertaken at intervals of 25 and 40 days after sowing. The per cent yellow mosaic virus disease incidence was recorded by counting the number of plants infected with the disease at 15 Days-days after 1<sup>st</sup> spray subsequently 15 Days-days after the 2<sup>nd</sup> spray and the total number of plants in the plot and converted to per cent incidence. The samples were taken to the laboratory and using stereo zoom microscope the live population of nymphs were counted. Data of whitefly population was recorded before and after spraying and taken at 20 DAS, 30 DAS followed by 45 DAS, further disease incidence was recorded at 15 days after 1<sup>st</sup> spray and 15 days after 2<sup>nd</sup> spray. The data thus obtained was subjected to ANOVA after using proper transformations.

**Comment [pL6]:** Abbreviations must be written in full when first appear!

$$\text{Percent disease incidence} = \frac{\text{Number of infected plants in a row}}{\text{Total number of plants in a row}} \times 100$$

### Where is the statistical analysis section?

**Formatted:** Indent: First line: 0 cm

### Result and discussion:

**Comment [pL7]:** Please write short sentences to clarify the effects described!!

### Effect of chemical and bio-pesticide on the incidence of MYMV and grain yield of *V. radiata* during, 2018-19 and 2019-20

Results obtained have been presented in Tables 1 and 2, and also depicted in Figs 1 and 2. During *Kharif*, 2018-19, two sprays of A-One (Imidacloprid 17.8% SL) on 25 and 40 days after sowing (DAS) were recorded highly efficacious and best for the management of the disease. Foliar spray application of A-One (Imidacloprid) 17.8 SL @ 0.5/lit (T<sub>2</sub>) on 25 DAS and 40 DAS recorded higher grain yield (9.04q/ha), lowest disease incidence of MYMV (18.70%) and highest disease reduction over control (61.41%). This treatment was followed by Success (Spinosad) 2.5% SC @ 0.5ml/lit (T<sub>4</sub>) which recorded disease incidence of MYMV of 20.88%, disease reduction over control of 56.91%; and grain yield of 8.37q/ha, table 1, fig. 1. During *Kharif*, 2019-20, foliar sprays of A-One (Imidacloprid) 17.8 SL @ 0.5/lit (T<sub>2</sub>) on 25 DAS and 40 DAS recorded notable higher grain yield (9.16q/ha) with the lowest disease incidence of MYMV (17.60%) and highest disease reduction over control (62.10%). This is followed by Success (Spinosad) 2.5% SC @ 0.5ml/lit (T<sub>4</sub>) disease incidence of MYMV (19.48%), disease reduction over control (58.06%) and grain yield (8.80q/ha). Mean of two-year data showed lowest disease incidence and least number of whiteflies was recorded in A-One (Imidacloprid) treated plots. Fifteen days after 1<sup>st</sup> spray and 2<sup>nd</sup> spray recorded 17.09%

**Formatted:** Highlight

**Comment [pL8]:** Please correct name throughout the manuscript

&18.15% disease incidence respectively, and also reported least number of whiteflies being 3.10 five days after 1<sup>st</sup> spray and 3.32 five days after 2<sup>nd</sup> spray which consequently resulted in higher yield (9.10 q/ha), table 2, fig. 2. Effectiveness of A-One (Imidacloprid 17.8 % SL) for the management of whitefly and MYMV disease, as it is having nicotinic acetylcholine receptor agonists. Similar results have been reported by other workers under different agro-climatic situations (Hossain et al. [5]; Akram & Naimuddin [1]; Jayappa et al. [3]; Islam et al. [6]; Reang et al. [10]).

During *Kharif* 2018-19, in bio-pesticides Vertifire-L1% WP @ 1.5 g/lit (T<sub>7</sub>) sprayed plot disease incidence of MYMV of 26.41%, disease reduction over control (45.50%) and grain yield (6.15/ha) followed by Daman 2 % WP@ 1.5g/lit(T<sub>6</sub>) sprayed plot disease incidence of MYMV of 27.60%, disease reduction over control of 43.04% and grain yield of 6.03q/ha, table 1, fig. 1. During *Kharif* 2019-20, Vertifire-L 1% WP @ 1.5 g/lit (T<sub>7</sub>) sprayed plot disease incidence of MYMV of 23.82%, disease reduction over control of 48.71% and grain yield of 7.46q/ha followed by Daman 2 % WP@ 1.5g/lit(T<sub>6</sub>) sprayed plot disease incidence of MYMV (25.94%), disease reduction over control (44.15%) and grain yield (6.72q/ha). Two sprays of Daman (*Beauveria bassiana*) and Neem Baan (Neem Oil) were found to be moderately effective in managing the disease and whiteflies population recording, 22.82 and 25.11 per cent incidence at 15 days after first spray and 25.32 and 26.73 per cent at 15 days after second spray. Whitefly counts of 4.19 and 6.13 per plant at 5 days after first spray and 4.32 and 6.41 whiteflies per plant at 5 days after second spray, table 2, fig. 2. Efficacies of bio-pesticides in the management of whitefly –borne MYMV, it acts as insect antifeeding and affect fecundity of insects, the azadirachtin-based product can also act as an insect growth regulator by blocking the synthesis and release of moulting hormones leading to incomplete moulting of immature insects. Similar results have also been reported by El Shafie [4], Sharma et al. [11], Sujatha & Bharpoda [13], Reang et al. [10] reported that among the five plant origin products with different doses tested as spray application, Neem Baan (Neem Oil) @ 0.2% showed lowest incidence (10.49%) and severity (9.58%).

## Conclusion

Comprehensive analysis of two year's data showed that combination of seed priming/ treatment with A-One (Imidacloprid) 17.8 SL @ 3ml/kg and two foliar sprays of A-One (Imidacloprid) 17.8 SL @ 0.5/lit (T<sub>1</sub>) sprayed on 25 DAS and 40 DAS recorded lowest disease incidence of MYMV 15.37% and 17.90%, respectively. Two sprays of Daman (*Beauveria bassiana*) (2% WP@ 1.5 g/L) and Azadirachtin oil (0.2%) were efficacious in the management of whitefly vector population build-up.

**Table-1: Population of whitefly, *B. tabaci* after the application of in Chemicals-chemical and bio-pesticides on-in management of transmitted MYMV in *V. radiata* during 2018-19 & 2019- 20**

Treatment		No. of Vector (DAS)								
		20			30			45		
		2018-19	2019-20	Mean	2018-19	2019-20	Mean	2018-19	2019-20	Mean
1	Dupont Tm Rekord TM @0.5g/lit	21.69 (4.71)	20.20 (4.54)	20.94	4.15 (2.15)	3.73 (2.06)	3.94	4.57 (2.25)	4.17 (2.14)	4.37
2	A-One @0.5ml/lit	18.46 (4.35)	17.80 (4.27)	18.13	3.27 (1.94)	2.93 (1.85)	3.10	3.55 (2.01)	3.10 (1.89)	3.32
3	Rogor Plus @ 1.5ml/lit	20.99 (4.63)	19.10 (4.40)	20.04	3.88 (2.09)	3.54 (2.01)	3.71	4.11 (2.15)	4.04 (2.13)	4.07
4	Success @0.5 ml/lit	19.21 (4.43)	18.51 (4.36)	18.86	3.31 (1.95)	3.18 (1.91)	3.24	3.75 (2.06)	3.35 (1.96)	3.55
5	Maxtara @ 0.5g/lit	22.44 (4.77)	22.78 (4.81)	22.61	4.97 (2.32)	4.31 (2.19)	4.64	5.27 (2.40)	4.67 (2.27)	4.97
6	Daman @ 1.5g/lit	23.15 (4.86)	23.81 (4.93)	23.48	4.33 (2.19)	4.06 (2.13)	4.19	4.53 (2.24)	4.11 (2.15)	4.32
7	Vertifire-L @ 1.5 g/lit	25.59 (5.09)	24.59 (5.00)	25.09	6.77 (2.69)	6.09 (2.57)	6.43	6.99 (2.73)	6.59 (2.66)	6.79
8	Neem Baan @ 0.2%	24.77 (5.02)	23.77 (4.92)	24.27	6.31 (2.61)	5.95 (2.54)	6.13	6.81 (2.70)	6.01 (2.55)	6.41
9	Control	27.77 (5.30)	28.10 (5.35)	27.93	7.38 (2.80)	7.76 (2.87)	7.57	7.32 (2.78)	7.89 (2.90)	7.60
	SEm±	0.21	0.20	0.205	0.12	0.08	0.1	0.10	0.09	0.09
	CD 5%	0.64	0.61	0.62	0.35	3.73	2.04	0.31	0.28	0.29
	CV %	7.65	7.39	7.52	8.74	2.93	5.83	7.47	7.13	7.3

\*The Trade and Chemical Names of Chemicals are used in the Trial is Dupont<sup>™</sup> Rekord TM (Acetamiprid 20% SP), Maxtara (Thamethoxam 25% WG), A-One (Imidacloprid 17.8% SL), Rogor Plus (Dimethoate 30% EC), Success (spinosad 2.5% SC), Daman (Beauveria bassiana 2% WP, CFUI 2X10<sup>8</sup>), Vertifire-L (Verticillium lecani 1% WP, CFUI 1X10<sup>8</sup>), Neem Baan (Azadiractin 0.15%W/W 1500 ppm).

Comment [pL9]: Please ±SD to the mean value!!

Comment [pL10]: What these numbers refer to??

Formatted: Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Formatted: Font: 11 pt, Highlight

Formatted: Highlight

Comment [pL11]: These infos are in the table, please consider deleting. But you need to add statistical analysis indicators here.

Formatted: Highlight

Formatted: Font: 11 pt, Highlight


Formatted: Highlight

**Table-2: Effect of Chemicals and bio-pesticides on management of whitefly, *B. tabaci* transmitted MYMV in *V. radiata* during, 2018-19 & 2019- 20**


**Comment [pL13]:** Re-phrase!!! It's very vague

Treatments (Brand names of Pesticides)		15 Days after 1 <sup>st</sup> Spray			15 Days after 2 <sup>nd</sup> Spray			% Reduction over control			Yield (q/ha)		
		2018-19	2019-20	Mean	2018-19	2019-20	Mean	2018-19	2019-20	Mean	2018-19	2019-20	Mean
1	Dupont Tm Rekord TM	23.06 (28.61)	22.04 (27.88)	22.55	24.76 (29.75)	23.21 (28.68)	23.98	48.90	50.03	49.46	7.35	7.62	7.48
2	A-One)	17.55 (24.65)	16.63 (24.05)	17.09	18.70 (25.53)	17.60 (24.76)	18.15	61.41	62.10	61.75	9.04	9.16	9.10
3	Rogor Plus	20.75 (27.03)	19.09 (25.86)	19.92	21.51 (27.48)	20.92 (27.21)	21.21	55.61	54.96	55.28	8.13	8.54	8.33
4	Success	19.75 (26.26)	18.90 (25.70)	19.32	20.88 (27.15)	19.48 (26.15)	20.18	56.91	58.06	57.48	8.37	8.80	8.58
5	Maxtara	22.33 (28.15)	21.01 (27.70)	21.67	25.58 (30.36)	22.59 (28.33)	24.08	47.21	51.36	49.28	7.04	7.92	7.48
6	Daman	24.24 (29.49)	21.41 (27.17)	22.82	26.41 (30.91)	23.82 (29.21)	25.11	45.50	48.71	47.10	6.15	7.46	6.80
7	Vertifire-L	28.75 (32.42)	25.78 (30.45)	27.26	28.87 (32.46)	26.25 (30.70)	27.56	40.42	43.48	41.95	4.51	5.44	4.97
8	Neem Baan	25.87 (30.48)	24.36 (29.56)	25.11	27.60 (31.65)	25.94 (30.59)	26.77	43.04	44.15	43.59	6.03	6.72	6.37
9	Control	47.88 (43.78)	47.11 (43.34)	47.49	48.46 (44.11)	46.45 (42.96)	47.45	-	-		3.49	4.23	3.86
	SEm±	1.69	1.86	1.77	1.65	1.47	1.56				0.51	0.52	0.51
	CD 5%	5.06	5.56	5.31	4.95	4.41	4.68				1.54	1.56	1.55
	CV %	9.71	11.07	10.39	9.21	8.54	8.87				6.03	6.08	6.05

**Comment [pL12]:** What is this number represent? Is it percent of management? Average number of populations? Please clarify!!



**Fig. 1: Population of whitefly, *B. tabaci* in Chemicals and bio-pesticides on management of transmitted MYMV in *V. radiata* during 2018-19 & 2019- 20 [add statistical analysis to the graph](#)**



**Fig. 2: Effect of Chemicals and bio-pesticides on management of whitefly, *B. tabaci* transmitted MYMV in *V. radiata* during 2018-19 & 2019- 20**

**Reference:**

**Comment [pL14]:** Please support your introduction with more recent references!

1. Akram M, Naimuddin. Management of Mung bean yellow mosaic disease and effect on grain yield. *Indian J. Plant Protection*.2016; **44** (1): 127-131.
2. Borah B, Dasgupta I. Begomovirus research in India: A critical appraisal and the way ahead. *J. Biosci.* 2012; **37**(4):791- 806.
3. El Shafie HAF and Abdel raheem, BA. Field evaluation of three biopesticides for integrated management of major pests of tomato, *Solanum lycopersicum* L. in Sudan. *Agric. Biol. J. N. Am.* 2012; **3**(9): 340-344.
4. Hossain ASMA, Islam SMAS, Akhter N. and Muqit A.Effect of plant extracts, insecticides and cultural practices on growth characters and disease severity of Mung bean yellow mosaic. *International Journal of Sustainable Crop Production*.2010; **5**(2): 16-20.
5. Islam Md. S, Hossain Md. B, Saleh A, Shahriar, Fatema Begum & Md N H Sani. Management of Yellow Mosaic Disease of Mung bean Against *Mung bean Yellow Mosaic Virus (MYMV)* in Bangladesh. *Int. J. of App. Sci.* 2018;**1** (2) 105-106.
6. Jayappa Ramappa HK and Devamani BD. Management of Mung bean Yellow Mosaic Virus (MYMV) in Mung bean (*Vigna radiata* L.). *J. Entomol. Zool. Stud.*2017;**5**(5): 596-601.
7. Nariani, T.K. (1960). Yellow mosaic of mung (*Phaseolus aureus* L.). *Indian Phytopathol.***13**: 24-29.
8. Nene, Y. L., (1973) Viral diseases of some warm weather pulse crops in India; *Plant Dis. Rep.* **57**: 463-467.
9. Pulses revolution from food to nutritional security. Min. of Agri. & FW (DAC&FW), GOI. 2017-18.
10. Reang D, Devi MR, Dhar A and Nath PS. Comparative Evaluation of Botanicals and Chemical Insecticides and their Suitable Use for the Management of Yellow Mosaic Virus (YMV) of Mung bean. *Int. J. Curr. Microbiol. App. Sci.*2018;**7**(9): 2094-2101.

11. Sharma M, Budha PB and Pradhan SB. Efficacy Test of Bio-pesticides against Tobacco Whitefly *Bemisia tabaci* (Gennadius, 1889) on Tomato Plants in Nepal. *J. of Inst. of Sci. and Tech.*, 2015; **20** (2):11-17.
12. Sharma SR, Varma A. Effect of cultural practices on virus infection in cowpea. *Z fur Acker Pflanzenbau*. 1984; 153:21-31.
13. Sujatha B and Bharpoda TM Bio-efficacy of Biopesticides against Sucking Pests in Green Gram Grown during *Kharif*. *Int. J. Pure App. Biosci.* 2017; **5** (4): 1827-1834.

UNDER PEER REVIEW