

Performance Effect of Super Absorbent Polymer and Humic Acid on Maize Yield and Nutrient Uptake under on Rainfed Alfisols

ABSTRACT

A field study was carried out to study the performance effect of superabsorbent polymer at 2.5 and 4.5 kg ha⁻¹ and humic acid at 15 and 30 kg ha⁻¹ alone and their combinations with recommended 100% fertilizers on yield, uptake, and photosynthetic pigments of maize grown on rainfed alfisols under rainfed conditions at Regional Agricultural Research Station, Palem, Telangana. Results showed that Conjoint application of hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ along with 100% RDF package significantly increased the pooled grain and stover yield (7136 and 8457 kg ha⁻¹) of maize. Irrespective level of hydrogel and humic acid combinations with 100% RDF increased the macronutrients contained in uptake by grain, stover and total uptake and showed a similar pattern was also observed in corresponding to grain and stover yield. The higher-chlorophyll "a", "b" and total chlorophyll content (1.81, 1.69 1.54; 0.69, 0.62, 0.55 and 2.65, 2.46 and 2.24 mg g⁻¹ in fresh plant weight at 30, 60 and 90 DAS) were significantly influenced with by the soil application of hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹. In conclusion, The present investigation indicates the positive interaction of between humic acid with and super absorbent polymer to twich improved nutrient uptake and maize yield and uptake.

Keywords: Super absorbent polymer, Humic acid, Hydrogel, Recommended dose of fertilizer, Maize and Chlorophyll content

1. INTRODUCTION

In many regions of the world, including India, drought stress is one of the most important factors that decrease agricultural crop production (Sah *et al.*, 2020). Maize is the third leading cereal crop of the world after wheat and rice, and it seems to be moderately forbearing the moisture stress in Alfisols. In India, the crop is grown more than 76.03 lakh hectares (use international unit) under rainfed conditions (Agricultural Market Intelligence Centre, PJTSAU, 2021) with the erratic distribution of rainfall, which leads terminal and dry spell This finally leading leads to reduced yields. In view of global climate change, drought stress aggravates soil moisture availability especially during grain development, and which is responsible for 20–30 % yield losses mainly due to undersized kernels (Gao *et al.*, 2017). According to this basis, one One of the ways to improve the soil moisture in Alfisols is by applying super absorbent polymer along with organic substances that supply hold water and nutrients for crop roots (Pawlowski *et al.*, 2009).

Hydrogels are superabsorbent polymers (SAPs) which is capable to on an average, hold 332-465 times water of its weight and release it slowly releases water during in drought stress conditions in light soils (Dehkordi, 2016). It to regulate soil water availability to for crops, and to favor the efficiency of nutrient absorption and to improve the crop yields (Mendonça *et al.*, 2013). Several research workers highlighted the positive benefits (Rehman *et al.*, 2011; Singh, 2012; Langaroodi *et al.*, 2013) in terms of moisture conservation, as well as yield improvement in many crops. On the other hand Meanwhile, humic acid is an organically charged bio-stimulant that significantly affects plant growth, and development to increase and crop yield. It has been extensively investigated (Liu *et al.*, 2019) that humic acid stimulates microbial activities in the rhizosphere due to the biochemical decomposition of animal and plant residues (El-Howeity *et al.*, 2019), improves photosynthetic

efficiency and soil-water properties (Shah *et al.*, 2018b). Many authors have highlighted the production advantages of humic acid application on various crops (Morozesk *et al.*, 2017; Olaetxea *et al.*, 2018), further it also helps in moisture retention and mitigation of salinity (Yoon- Ha Kim *et al.*, 2012). The studies made to understand the combined application of superabsorbent polymers with and humic acid substances were lacking. Apart from this the physiological basis for improvement of yield when applied SAP materials with and humic acid needed to be understood in detail.

2. MATERIAL AND METHODS

A field experiment was conducted at Regional Agricultural Research Station, of Professor Jayashankar Telangana State Agricultural University, Palem, Telangana state, India during *kharif*, (.....) 2015 and 2016. The station is located in a under-rainfed conditions area. Objective of the research was to find out the performance effect of conjoint application of hydrogel and humic acid on maize yield and nutrient uptake in Alfisols. The characteristics of the soils at experimental site soil was sandy loam in texture, slightly alkaline (pH 7.61) in reaction, non-saline (0.27 dS m^{-1}), low in organic carbon (0.44 percent), available N ($159.61 \text{ kg ha}^{-1}$), high in available P_2O_5 (61.38 kg ha^{-1}) and medium in K_2O ($317.21 \text{ kg ha}^{-1}$). The experiment was laid out in a randomized block design with ten treatments replicated three times. thrice, consists combination of two factors i.e., hydrogel and humic acid with control treatment where no input was applied. The maize hybrid DHM 117 was used as test crop by imposing treatments composition as follows: T₁: 100% RDF, T₂:100% RDF + Humic acid 15 kg ha^{-1} , T₃:100% RDF + Humic acid 30 kg ha^{-1} , T₄: 100% RDF + Hydrogel 2.5 kg ha^{-1} , T₅:100% RDF + Hydrogel 4.5 kg ha^{-1} , T₆:100% RDF + Humic acid $15 + \text{Hydrogel } 2.5 \text{ kg ha}^{-1}$, T₇:100% RDF + Humic acid $15 + \text{Hydrogel } 4.5 \text{ kg ha}^{-1}$, T₈:100% RDF + Humic acid $30 + \text{Hydrogel } 2.5 \text{ kg ha}^{-1}$, T₉:100% RDF + Humic acid $30 + \text{Hydrogel } 4.5 \text{ kg ha}^{-1}$. All the treatments received a uniform recommended dose of fertilizers i.e., 200 kg N , $60 \text{ kg P}_2\text{O}_5$ and $50 \text{ kg K}_2\text{O ha}^{-1}$ through urea, diammonium phosphate, and muriate of potash. Respectively. The basal dose of fertilizers (33 % N, 100 % P and 50% K) were applied at the time of sowing, and the remaining 67 % N applied in two split doses at 25 and 55 DAS, and 50% K was applied at 25 DAS by pocketing method at the base of individual plants. As per the treatment inception, required quantities of hydrogel and humic acid were applied at a depth of 8-10 cm in rows, where the test crop was sown with a spacing of $60 \times 20 \text{ cm}$. The plant samples were collected for the analysis of photosynthetic pigments by following Lichtenthaler and Wellbum (1983) method and nutrient uptake studies described by Piper (1966). The grain and stover yield was were recorded at the time of harvesting from the randomly tagged five plants and expressed grain and stover yield as kilograms per hectare (Kg ha^{-1}).

Further, the data was analyzed statistically in a randomized block design using OPSTAT, and significance of the treatment effect was determined using the F-test. The least significant differences were calculated at the 5% probability level to determine the significance of the difference between two treatments (Gomez and Gomez, 1984).

3. RESULTS AND DISCUSSION

3.1 Grain and Stover Yield (kg ha^{-1}) of Maize

The grain and stover yield of maize were influenced significantly by the application of superabsorbent polymers in combination with humic acid and 100% RDF package, and is presented in Table 1. Grain and stover yield of maize varied from 4458 to 6977 and 5158 to 8363 kg ha^{-1} respectively during *kharif*, 2015, and ranged 4731 to 7294 and 5346 to 8551 kg ha^{-1} respectively during *kharif*, 2016. Soil application of hydrogel@ 4.5 kg ha^{-1} + humic acid@ 30 kg ha^{-1} along with 100% RDF recorded-produced the highest grain and stover yields (6977 & 8363 and 7294 & 8551 kg ha^{-1}) during *kharif*, 2015 and 2016, and was significantly higher over rest of the treatments. It may be attributed to improvement in the water holding capacity induce by hydrogel as well as of the soil due to humic acid coupled with supersorbing properties of the hydrogel (reference.....) which absorbs the t- water and releases it slowly to the growing plants. Available water as per thefor crop requirement stimulates

along with nutrients uptake which to enhance photosynthetic activity which further resultant in higher-high yields. Significantly lower-low grain and stover yields were noticed when maize was grown only with recommended dose of fertilizers (6022 & 7208 and 6319 & 7028 kg ha⁻¹) in both the years, and pooled yield found as 6171 and 7122 kg ha⁻¹, compared to the significantly The highest yield was on registered treatment T_e (100% RDF + Humic acid 30 + Hydrogel 4.5 kg ha⁻¹). The ultimate lowest grain and stover yields were registered in-ion the control treatment (4458 & 5158 and 4731 & 5346 kg ha⁻¹ respectively), during *kharif*, 2015 & 2016; and pooled yield was noticed as 4595 & 5252 kg ha⁻¹ where no input was added. Further, pooled grain and stover yield varied from 4595 to 7136 and 5252 to 8457 kg ha⁻¹. Among the treatments, integrated soil application of hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ with 100% RDF package has resulted in significantly higher-high pooled grain and stover yield (7136 and 8457 kg ha⁻¹), even when maize crop experienced moisture stress during crop critical stages like a knee-high stage, tasseling, cob initiation, and soft dough stage during *kharif*, 2015 and 2016. The positive effect of superabsorbent polymers in increasing the t-yields-yield of mize was reported by Gunes *et al.*, (2016) and Kumari *et al.*, (2017) in-maize-. Similarly, Baldotto *et al.*, (2019) reported an increase of 15% in corn grain yield with the humic acid application (Oktem *et al.*, 2018).

3.2 Macro-Nutrient Uptake (kg ha⁻¹) of Maize in Grain and Stover

The pooled macronutrient (NPK) uptake-contained in grain and stover of test crop at harvest, was significantly affected by the judicious use of super absorbent polymers + humic acid along with recommended fertilizer package of test crop-(Fig 1) at-harvest. Among different treatments, integration of hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ with 100% RDF was found to be the best nutrient management practice which resulted in significantly higher N, P, K uptake in grain and stover (88.86, 17.90, 88.84 and 61.47, 21.42, 93.14 kg ha⁻¹), and total uptake (150.33, 39.32 and 131.99 kg ha⁻¹) respectively in comparison to alone application of 100% RDF (67.97, 11.81, 26.45 and 37.55, 13.84, 67.40 kg ha⁻¹), with total uptake of 103.11, 25.83 and 95.29 kg ha⁻¹ respectively. The higher NPK uptake of maize grain and stover registered with the application of 100% RDF along with hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ than alone 100% RDF package may be attributed to better crop growth due to higher-removal-of-high nutrients-availability under-by- availing moisture for nitrification process as reported by Mandal *et al.* (2015) in maize crop. With the-available-favourable soil moisture content, the nutrients may also be released from the saturated zone resulted in higher-high-solubility, availability and uptake of nutrients (Shahid *et al.*, 2016).

3.3 Chlorophyll 'a', 'b' and total chlorophyll content (mg g⁻¹) of Maize

The pooled data on chlorophyll "a", "b" and total chlorophyll content of maize was influenced at different growth stages are presented in Fig 2. Soil-application-Application of 100% RDF along with hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ recorded-resulted in significantly higher chlorophyll "a" (1.81, 1.69 and 1.54 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively), chlorophyll "b" (0.69, 0.62 and 0.55 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively) and total chlorophyll content (2.65, 2.46 and 2.24 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively). Whereas, significantly lower chlorophyll "a" (1.54, 1.37 and 1.22 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively), chlorophyll "b" (0.45, 0.42 and 0.38 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively) and total chlorophyll content (2.13, 1.97 and 1.75 mg g⁻¹ fresh weight at 30, 60 and 90 DAS respectively) was noticed in maize with the alone application of recommended fertilizer package. Application of 100% RDF coupled with hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ has recorded significantly higher chlorophyll "a", "b" and total chlorophyll content compared to the maize grown with the alone application of recommended fertilizer dosage and it confirms the positive relationship with grain yield of maize. According to Akhter *et al.*, (2004), addition of 0.1, 0.2 and 0.3% hydrogel increased the moisture retention at field capacity with a linear correlation linearly of (r= 0.988), thus the amount of plant-available water was significant in both sandy loam and loam

soils. Application of hydrogel can help the crop further to avoid moisture deficit and to improve crop yield (Mohammadkhani and Heidari, 2007), compared to without hydrogel application

4. CONCLUSION

The results of this experiment indicated that soil a Application of hydrogel@4.5 kg ha⁻¹ + humic acid@30 kg ha⁻¹ along with 100% recommended fertilizer package on Alfisols resulted in the highest pooled grain and stover yield of mize, i.e. of 7136 and 8457 kg ha⁻¹ respectively and was found better than other treatments. Conjoint application of super absorbent polymers along with humic acid substances had a positive influence on yield and by increasing survivability of maize plants under moisture deficit conditions on rainfed Alfisols, especially at critical growth stages of maize grown alfisols under rainfed conditions. favored the physiological activities and

Table 1: Influence of different levels of Hydrogel and Humic acid on maize grain and stover yield (kg ha⁻¹)

Treatments	2015		2016		Pooled	
	Grain Yield	Straw Yield	Grain Yield	Straw Yield	Grain Yield	Straw Yield
	(Kg ha ⁻¹)					
T ₁ : 100% RDF	6022	7028	6319	7216	6171	7122
T ₂ : 100% RDF + Humic acid 15 kg ha ⁻¹	6205	7332	6583	7520	6394	7426
T ₃ : 100% RDF + Humic acid 30 kg ha ⁻¹	6313	7521	6602	7709	6458	7615
T ₄ : 100% RDF + Hydrogel 2.5 kg ha ⁻¹	6361	7496	6674	7684	6518	7590
T ₅ : 100% RDF + Hydrogel 4.5 kg ha ⁻¹	6484	7584	6711	7772	6598	7678
T ₆ : 100% RDF + Humic acid 15 kg ha ⁻¹ + Hydrogel 2.5 kg ha ⁻¹	6653	7962	6948	8150	6801	8056
T ₇ : 100% RDF + Humic acid 15 kg ha ⁻¹ + Hydrogel 4.5 kg ha ⁻¹	6875	8101	7152	8289	7014	8195
T ₈ : 100% RDF + Humic acid 30 kg ha ⁻¹ + Hydrogel 2.5 kg ha ⁻¹	6891	8139	7186	8327	7039	8233
T ₉ : 100% RDF + Humic acid 30 kg ha ⁻¹ + Hydrogel 4.5 kg ha ⁻¹	6977	8363	7294	8551	7136	8457
T ₁₀ : Control	4458	5158	4731	5346	4595	5252
SEm (+)	16.30	33.07	10.52	21.38	14.69	28.51
CD (P = 0.05)	51.73	87.21	27.39	60.14	39.05	73.29




Fig 1. Effect of different levels of Hydrogel and Humic acid on pooled grain NPK uptake (kg ha⁻¹) of maize




Fig 2. Influence of different levels of Hydrogel and Humic acid on pooled stover NPK uptake (kg ha⁻¹) of maize




Fig 3. Influence of different levels of Hydrogel and Humic acid on pooled total NPK uptake (kg ha⁻¹) of maize




Fig 4. Effect of different levels of Hydrogel and Humic acid on pooled chlorophyll "a" (mg g⁻¹) of maize




Fig 5. Effect of different levels of Hydrogel and Humic acid on pooled chlorophyll "b" (mg g^{-1}) of maize




Fig 6. Effect of different levels of Hydrogel and Humic acid on pooled total chlorophyll (mg g^{-1}) of maize

REFERENCES

1. Akhter, J., Mahmood, K., Malik, K.A., Mardan, A., Ahmad, M., and Iqbal, M.M. 2004. Effects of hydrogel amendment on water storage of sandy loam and loam soils and seedling growth of barley, wheat and chickpea. *Pl. Soil and Env.*, 50(10): 463-469.
2. Baldotto, M. A.; Melo, R. O.; Baldotto, L. B. Field corn yield in response to humic acids application in the absence or presence of liming and mineral fertilization. *Semina: Ciências Agrárias*, v.40, p.3299-3304, 2019. <https://doi.org/10.5433/1679-0359.2019v40n6Supl2p3299>
3. Barron J, Okwach G. Run-off water harvesting for dry spell mitigation in maize (*Zea mays* L.) results from on-farm research in semi-arid Kenya. *Agric. Water Manage.* 2005; 74:1-21.
4. Dar SB, Mishra D, Zahida R, Afshana BB. Hydrogel: To enhance crop productivity per unit available water under moisture stress *Agriculture. Bull. Env. Pharmacol. Life Sci.* 2017; 6(10):129-135. Dehkordi,

- D.K. 2016. The effects of superabsorbent polymers on soils and plants. *Pertanika. J. Trop. Agric. Sci.*, 39(3): 267-298.
5. El-Howeity, M. A.; Abdel-Gwad, S. A.; Elbaalawy, A. M. Effect of bio-organic amendments on growth, yield, nodulation status, and microbial activity in the rhizosphere soil of peanut plants under sandy soil conditions. *Journal of Soil Sciences and Agricultural Engineering*, v.10, p.299-305, 2019. <https://doi.org/10.21608/jssae.2019.43220>
 6. Gao, Z., Liang, X. G., Zhang, L., Lin, S., Zhao, X., Zhou, L. L., et al. (2017). Spraying exogenous 6-benzyladenine and brassinolide at tasseling increases maize yield by enhancing source and sink capacity. *Field Crops Res.* 211, 1–9. doi: 10.1016/j.fcr.2017.05.027
 7. Gomez, K. A., and Gomez, A. A., 1984. *Statistical procedures for Agricultural research*. John-wiley and sons, Inc, New York, pp-680.
 8. Gunes, A., Nurul Kitir, Metin Turan, Erdal Elkoca, E, Yildirim, E., and Nazmiye, 2016. Evaluation of effects of water-saving superabsorbent polymer on corn (*Zea mays* L.) yield and phosphorus fertilizer efficiency. *Turkish J. Agri. Forestry*, 40: 365-378.
 9. Kumari, S., Solanki, N. S., Dashora, L. N., and Upadhyay, B., 2017. Effect of superabsorbent polymer and plant geometry on growth and productivity of maize (*Zea mays* L.). *J. Pharmacognosy Phytochemistry*, 6(4): 179-181.
 10. Langaroodi, N.B.S., Ashouri, M., Dorodian, H.R., and Azarpour, E. 2013. Study effects of super absorbent application, saline water and irrigation management on yield and yield components of peanut (*Arachis hypogaea* L.). *Ann. Biolog. Res.*, 4(1): 160-169.
 11. Lichtenthaler, H. K., and Wellbum, A. R., 1983. Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochem. Soc. Trans.*, 11: 591-592.
 12. Liu, M.; Wang, C.; Wang, F.; Xie, Y. Maize (*Zea mays* L.) growth and nutrient uptake following integrated improvement of vermicompost and humic acid fertilizer on coastal saline soil. *Applied Soil Ecology*, v.142, p.147-154, 2019. <https://doi.org/10.1016/j.apsoil.2019.04.024>
 13. Mandal UK, Sharma KL, Venkanna K, Korwar GR, Reddy KS, Pushpanjali NN *et al.* Evaluating hydrogel application on soil water availability and crop productivity in semiarid tropical red soil. *Indian J. Dryland Agric. Res. Dev.* 2015; 30(2):01-10.
 14. Mendonça, T. G.; Urbano, V. R.; Peres, J. G.; Souza, C. F. Hidrogel como alternativa no aumento da capacidade de armazenamento de água no solo. *Water Resources and Irrigation Management*, v.2, p.87-92, 2013.
 15. Mohammadkhani, N., and Heidari, R., 2007. Effects of water stress on respiration, photosynthetic pigments and water content in two maize cultivars. *Pakistan J. Biol. Sci.*, 10(22): 4022-4028.
 16. Morozesk, M.; Bonomo, M. M.; Souza, I. da C.; Rocha, L. D.; Duarte, I. D.; Martins, I. O.; Dobbss, L. B.; Carneiro, M.T. W. D.; Fernandes, M. N.; Matsumoto, S. T. Effects of humic acids from landfill leachate on plants: An integrated approach using chemical, biochemical and cytogenetic analysis. *Chemosphere*, v.184, p.309-317, 2017. <https://doi.org/10.1016/j.chemosphere.2017.06.007>
 17. Oktem, A.; Celik, A.; Oktem, A.G. Effect of humic acid seed treatment on yield and some yield characteristic of corn plant (*Zea mays* L. indentata). *Journal of Agricultural, Food and Environmental Sciences*, v.72, p.142-147, 2018.
 18. Olaetxea, M.; Hita, D. de; Garcia, C. A.; Fuentes, M.; Baigorri, R.; Mora, V.; Garnica, M.; Urrutia, O.; Erro, J.; Zamarreno, A. M.; Berbara, R. L. Hypothetical framework integrating the main mechanisms involved in the promoting action of rhizospheric humic substances on plant root-and shoot-growth. *Applied Soil Ecology*, v.123, p.521-537, 2018.
 19. Pawlowski A, Lejcus K, Garlikowski D, Orezesyna H (2009). Geocomposite with superabsorbent as an element improving water availability for plants on slopes. *Geophys. Res. Abstr.*, 11: 1-2
 20. Piper CS. *Soil and plant analysis*. Hans Publisher, Bombay, 1996.
 21. Rehman, A., Ahmad, R., and Safdar, M. 2011. Effect of hydrogel on the performance of aerobic rice sown under different techniques. *Pl. Soil and Env.*, 57(7): 321-325.
 22. Sah, R.P., Chakraborty, M., Prasad, K. *et al.* Impact of water deficit stress in maize: Phenology and yield components. *Sci Rep* 10, 2944 (2020). <https://doi.org/10.1038/s41598-020-59689-7>
 23. Shah, Z. H.; Rehman, H. M.; Akhtar, T.; Alsamadany, H.; Hamooh, B.T.; Mujtaba, T.; Daur, I.; Al Zahrani, Y.; Alzahrani, H. A.; Ali, S.; Yang, S. H. Humic substances: Determining potential molecular regulatory processes in plants. *Frontiers in Plant Science*, v.9, p.263, 2018b. <https://doi.org/10.3389/fpls.2018.00263>

24. Shahid BD, Ram H. Grain yield, nutrient uptake and water-use efficiency of wheat (*Triticum aestivum*) under different moisture regimes, nutrient and hydrogel levels. *Indian J Agron.* 2016; 61(1):101-104.
25. Singh, H. 2012. Effect of hydrogel on growth, yield and water use efficiency in pearl millet (*Pennisetum glaucum*) production. *Forage Res.*, 38(1), 27–28.
26. Tavakkoli AR, Oweis TY. The role of supplemental irrigation and nitrogen in producing bread wheat in the highlands of Iran. *Agric. Water Manag.* 2004; 65:225-236.
27. Yoon-Ha Kim YH, Khan AL, Shinwari ZK, Kim DH, Waqas M, Kamran M, In-J L, 2012. Silicon treatment to rice (*Oryza sativa* L cv 'Gopumbyeo') plants during different growth periods and its effects on growth and grain yield. *Pak J Bot* 44(3): 891-897
28. Zahedi H, Noormohammadi G, Shirani-Rad AH, Habibi D, Mashhadi-Akbar-Boojari M (2009). The effects of Zeolite and foliar applications of selenium on growth, yield and yield components of three canola cultivars under drought stress. *World Appl. Sci. J.*, 7(2): 255-262.A

UNDER PEER REVIEW