

Original Research Article

Response of Date of Sowing and Varieties on Growth and Yield of Niger Under Eastern Ghat High Land Zone of Odisha

ABSTRACT

Niger (*Guizotia abyssinica* (L. f.) Cass.), a minor oil seed crop is generally cultivated in marginal and sub marginal lands under poor agronomic practices without giving much importance to its optimum time of sowing which causes great losses in its yield. However, if the sowing time is optimized the crop has got high yield potential. Also the farmers always demand new varieties having greater production potential suitable to the particular agro-climatic situation. In view of this, a field experiment was carried out under All India Coordinated Research Project on Niger at Regional Research and Technology Transfer Station (OUAT), Semiliguda of Koraput district under Eastern Ghat High Land zone of Odisha in acidic soil during three consecutive *kharif* seasons from 2015 to 2017 to study the response of date of sowing and varieties on growth and yield of niger. The experiment consisted of six dates of sowing and two varieties of niger with split plot design. The pooled data for three years revealed that out of six different dates of sowing, the crop which was sown on 15th July resulted significantly the highest seed yield (563 kg ha⁻¹) with a net monetary return (Rs. 8083 ha⁻¹) and benefit cost ratio (1.6). Niger variety Utkal Niger-150 recorded higher seed yield and economic returns.

Key words: *Niger*; agroclimatic situation; -minor oilseed crop; -seed yield; Economics

1. INTRODUCTION

Niger (*Guizotia abyssinica* (L.f.) Cass.) is one of the most important minor oilseed crop. It is grown in India in an area of about 2.99 lakh ha with a production of 0.98 lakh tonnes and a productivity of 327 kg ha⁻¹. In the state of Odisha, it covers an area of 0.65 lakh ha with a production of 0.23 lakh tonnes and productivity of 360 kg ha⁻¹ [11]. The rainfall of the Eastern Ghat High Land zone of Odisha is very erratic and varies temporally and spatially. 75-80% of the total rainfall is received during the monsoon season only. The production and productivity of upland paddy is subjected to the drought like situation and thus poor yield. The upland area under paddy is 21,000 ha. To get a sustainable yield from this upland area, niger is found to be a good substitute due to its low water requirement and drought resistance capacity. Economic yield of crop may be limited by source (photosynthesizing organs) or sink (storage organs) or the capacity to translocate assimilates from one part to other limiting processes such as water or nutrient uptake or transport. In the crops like niger the source and sink are developed simultaneously, the adjustments of two components may be or readily achieved. The seasonal sequence of conditions play a major role in determining whether source or sink is more limiting. Establishment of a good stand is the essential pre-requisite for attaining high yields. It depends on time, depth and method of sowing. Sowing very early in the season may not be advantageous, say for example sowing rainfed niger in June may result in failure of the crop if there is prolonged dry spell from the second week of June to second week of July. However, sowing early in certain situations increases the yield. In other words, delayed sowing invariably reduces yield. Rainfed crop yields are reduced due to delay in sowing beyond June. The reduction in yields is attributed to early induction of flowering, unfavourable temperature and erratic rainfall. Most of the tropical crops are short day plants. Day length starts falling from July onwards, but the reduction in day length is steep from October onwards. Sowing the crop at optimum time increases yields due to suitable environment at all growth stages of crop. Flowering is induced after sufficient vegetative growth. Moisture stress or dry spells may be avoided during critical stages. Due to varied distribution pattern of rainfall, it is pertinent to know the exact sowing window of the crop for optimum yield. Good quality

Comment [A1]: Optimized how? Describe

Comment [A2]: To describe:
1- the periods
2- the varieties
3- the methodology for evaluating production and economic returns
4- results/conclusion

Remembering: It should briefly describe the purpose of the work, techniques and methods used, major findings with important data and conclusions.

Comment [A3]: This keyword is already in the title. I suggest that authors use keywords that are not repeated in the title, so you increase the visibility of your work in indexed databases

seed is a prime importance in agriculture. They should germinate rapidly and uniformly when sown with high yield potential. Emergence should be prompt and early vegetative growth should be vigorous and suitable for the particular agroclimatic situation. Farmers often demand the new varieties for their farming situation. Considering this, the varieties are also to be screened to test its suitability in this agroclimatic zone. The exact date of sowing and the suitable variety plays an important role in sustainable production. This could be attributed to overall improvement in plant vigour leading to better manifestation of yield attributes in niger. In niger, though research information on these aspects are available and very meager, so still a few more technical information are to be found out. Therefore, the present investigation is being carried out to study the response of date of sowing and the varieties on growth and yield of niger.

2. MATERIALS AND METHODS

2.1 Experimental site

To achieve the objectives of the programme, a field experiment was conducted under All India Coordinated Research Project (AICRP) on Niger at Regional Research and Technology Transfer Station (OUAT), Semiliguda under Koraput district in Eastern Ghat High Land zone of Odisha during three consecutive *kharif* seasons of 2015, 2016 and 2017. The farm is located in the geographical parallels of 18°42'N latitude, 82°30'E longitude and an altitude of 884.0 m. The region is marked by its warm and humid climate with an average annual rainfall of 1500 mm, most of which is received from middle of June to middle of October.

2.2 Soil Characteristics

The soil samples collected from field were air dried, grounded with mortar pestle and sieved in 2 mm sized mesh [3]. The soil of experimental site was red, sandy to clay loam in texture and acidic in reaction (pH 5.8) with available N (170 kg ha⁻¹), available P (16 kg ha⁻¹) and available K (145 kg ha⁻¹).

2.3 Experimental Design

The experiment consists of six dates of sowing with 15 days interval viz. 15th July, 30th July, 14th Aug, 29th Aug, 13th Sept and 28th Sept with two varieties of niger namely Deomali and Utkal Niger-150 (Table 1). The experiment was evaluated in split plot design with three replications for statistical analysis. The niger crop was sown with seed rate of 10 kg ha⁻¹ with a row spacing of 30 cm. The intra-row spacing of 10 cm was maintained by thinning operation. The thinning and weeding operations were carried out on 15 and 21 days after sowing in every year under the experimentation. The recommended dose of fertilizer (40:40:20 NPK kg ha⁻¹) was applied to the crop. Full dose of P, K and half N in form of Diammonium Phosphate (DAP), Muriate of Potash (MOP) and Urea were applied as basal and rest half N was applied after three weeks after sowing. The crop was harvested at physiological maturity. The periodical biometric and post harvest observations were taken at regular interval.

The weather data (Table 2) presented in Fig.1 showed that the rainfall received during crop period was 806.9 mm during 2015. Maximum temperature (Tmax) varied from 27.7 to 29.4 °C and minimum temperature (Tmin) varied from 4.9 to 17.2 °C. The total Sunshine hour received during the crop season was 880.5 hours. The maximum relative humidity varied from 85.3% to 95.3%. The weather data (Table 3) presented in Fig. 2 showed that the rainfall received during crop period was 1083.3 mm during 2016. The Tmax varied from 26.6 C to 28.1 °C and Tmin varied from 5.9 to 14.4 °C. The total sunshine hours received during the crop season was 115.4 hours. The maximum relative humidity varied from 95.6% to 96.9%. The weather data (Table 4) presented in Fig. 3 showed that the rainfall received during crop period was 1166.8 mm during 2017. The Tmax varied from 26.2 C to 28.6 °C and Tmin varied from 11.1 to 20.8 °C. The total Sunshine hour received during the crop season is 834.4 hours. The maximum relative humidity varied from 70.6% to 99.5%.

2.4 Statistical Analysis

The experimental data collected during the crop growth and harvest were analysed statistically following the procedure as described by Gomez and Gomez [4]. Treatment differences were tested at

Comment [A4]: I suggest write more about the varieties. There is a hypothesis about them? Or about the date of sowing?

Comment [A5]: The introduction has an interesting logical sequence, however, it lacks scientific information to support the manuscript. I suggest a strong literature review that will properly cite various written informations in the manuscript. Also to have new and more updated bibliographic references

Comment [A6]: Is the word correct?

Comment [A7]: This information could be in your abstract

Comment [A8]: I suggest reworking figures 1, 2 and 3 to allow readers to compare (even visually) on the same scale. By having 5 variables measured, it might be more interesting to divide them into more figures and that they have the same scale. It's very confused the way it is

5% level of significance by F test and using analysis of variance (ANOVA) for making comparison among treatment means for various yield and yield components of niger. Least significance difference (LSD) was done at $P=0.05$.

3. RESULTS AND DISCUSSION

3.1 Effect on Growth Parameters

The perusal of data shown in Table 5 revealed that the 15th July sown crop resulted significantly taller plants (225.7 cm) with maximum number of branches plant⁻¹ (7.8) at harvest. Shorter plant height (68.0 cm) with minimum number of branches plant⁻¹ (4.5) were recorded by the last sown crop on 28th September among various dates of sowing. Maximum plant height under 15th July sowing might be assigned due to the reason that the crop did not experience moisture stress at critical stages. The crop was exposed to favourable temperatures and high relative humidity. Delay in sowing times up to 28th September significantly reduces the plant height and number of branches plant⁻¹ in niger. Similar results were also reported by Agarwal et al. [1]; Jagtap et al. [5]; Nayak and Paikaray [9]; Paul et al. [11]; Mili et al. [12]. In case of varieties, Deomali recorded significantly taller plants (157.8 cm) which is 5.95% higher than Utkal Niger-150 (148.4 cm). This may be due to the genetic constituent of variety Deomali. Similar variability in growth parameters among the varieties was also reported by Mishra et al. [7] and Mohan Kumar et al. [8] in niger crop. The interaction effect of date of sowing x variety was found to be non-significant.

Comment [A9]: The author used others analysis, as Two-Way Anova. Must describe in the methodology all analysis that was used. Besides, need a improvement of description, according to the results.

3.2 Days to 50% Flowering and Maturity

Pooled data of 3 years (Table 5) showed a decreasing trend in days to 50% flowering and days to maturity with delay in date of sowing from 15th July to 28th September with an exception of 13th September (50% flowering). Maximum days to 50% flowering and days to maturity were recorded with date of sowing on 15th July (73 and 127 respectively). Minimum days to 50% flowering (42) and days to maturity (90) were recorded on 13th and 28th September respectively. This result was in conformity with the findings of Jagtap et al. [5]. The variation in growth parameters viz., earliness in flowering and maturity among the date of sowing might be due to the influence of environmental factors such as stress or less rainfall condition, temperature, sunshine, rainfall and relative humidity in late sown condition of the crop. In regards to varieties, both the varieties were significantly different with each other. Days to 50% flowering and days to maturity were significantly higher with respect to variety Deomali (55 and 108) as compared to Utkal Niger-150 (54 and 106).

Comment [A10]: A hypothesis could be created for this result. This is a interesting result

3.3 Effect on Yield Attributes

Pooled data of 3 years (Table 5) revealed that the number of capitula plant⁻¹ and number of seeds capitula⁻¹ varied significantly with different dates of sowing. Both these yield attributes showed a decreasing trend with delay in date of sowing from 15th July to 28th September. Similar finding was also reported by Ranjit et al., 2012. Date of sowing on 15th July recorded significantly higher number of capitula plant⁻¹ (67.6) and number of seeds capitula⁻¹ (29.5). Minimum number of capitula plant⁻¹ (17.0) and number of seeds capitula⁻¹ (16.5) were observed on last date of sowing on 28th September. Similar results were also reported by Agarwal et al. [1]; Jagtap et al. [5]; Nayak and Paikaray [9]; Paul et al. [11]. In case of varieties both the number of capitula plant⁻¹ and number of seeds capitula⁻¹ were found to be non-significant.

3.4 Seed Yield

The data on influence of date of sowing, variety and their interaction on seed yield is presented in Table 6 and Fig.4. The first date of sowing on 15th July recorded highest seed yield of 519.0 kg ha⁻¹ which was 78% more than that of last date of sowing on 28th September (113.9 kg ha⁻¹) during the year 2015. But during the year 2016, the second date of sowing on 30th July resulted significantly higher seed yield of 716.0 kg ha⁻¹ followed by the first date of sowing on 15th July (671.2 kg ha⁻¹). During the year 2017, the seed yield was noticed to range from 424.9 to 498.4 kg ha⁻¹. However, the pooled data analysis revealed that the first date of sowing on 15th July reported to give significantly

highest seed yield of 562.9 kg ha⁻¹ followed by the second date of sowing on 30th July (496.3 kg ha⁻¹). The seed yield of first date of sowing (15th July) was 12 to 77% higher than other dates of sowing up to the last date of sowing (28th September). Such significant reduction in seed yield of niger due to delay in sowing time has also been reported earlier by Singh et al. [13] and Mandal et al. [6]. The variation for seed yield might be due to variation in expression of characters fully in the favourable environmental conditions. The higher yield might be due to optimum temperature in early sown crop which was beneficial for the early establishment of crop and subsequent proper growth, resulting in producing more height, foliage and higher number of seeds per capitula which ultimately resulted in higher yield and full expression of the varietal characters during favourable conditions, which existed during early sown crops. These results were also in conformity with the reports of Mishra et al. [7] in niger and Babalad et al. [2] in soybean. Among the varieties, except during the year 2015, there was no significant difference in seed yield during 2016, 2017 and pooled data analysis. However, Utkal Niger-150 significantly out yielded Deomali with a percent increase of 4% over Deomali. It was observed that environment played major influence on yield level of variety than varieties themselves. The important environmental parameters were less dry spell, no wet spell, high sunshine hours and moderate temperature.

3.5 Economics

The input and output prices of commodities prevailed during each year of investigation were taken into consideration for calculating cost of production, net monetary return (NMR) and benefit: cost ratio (B:C ratio). The economic indices such as NMR and B:C ratio were worked out, statistically analyzed and presented in Table 7. 15th July date of sowing recorded significantly higher NMR during the year 2015 and 2017. The mean data showed that the sowing of niger on 15th July recorded highest NMR of Rs.8083 ha⁻¹ and B:C ratio 1.6 followed by 30th July and 14th August with NMR Rs.5414 ha⁻¹ and Rs.4623 ha⁻¹, respectively and B:C ratio 1.3 which was superior over rest of the dates of sowing. The last two dates of sowing (13th and 28th September) of niger showed negative NMR due to much delayed in date of sowing. In case of varieties Utkal Niger-150 recorded significantly higher NMR of Rs. 1390 ha⁻¹ with B:C ratio 1.1.

4. CONCLUSION

Based on the luxurious growth in the first sown crop associated with higher amount of rainfall, sunshine hours and optimum growing condition and yield attributes, seed yield and economic indices, it can be concluded that the most suitable date of sowing niger variety Utkal Niger-150 is 15th July for getting higher seed yield and monetary returns under Eastern Ghat High Land zone of Odisha.

REFERENCES

1. Agarwal KK, Jain KK, Sharma RS and Kashyap ML. Response of winter niger (*Guizotia abyssinica*) to sowing time and fertility levels. Journal of Oilseeds Research. 1996; 13(1): 122-123.
2. Babalad HB, Lingaraju BS, Radder GD. Studies on soybean cultivation during non-traditional rabi/summer under irrigated conditions. Journal of Oilseeds Research. 1996; 13(1): 66-69.
3. Brady N. The nature and properties of soils. 10th Edition, Macmillan Publishing Company, Cranbury. 1990.
4. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. 2nd Edition, John Wiley and Sons, New York. 1984; p. 680.
5. Jagtap PK, Sandipan PB, Patel KM, Patel MC. Effect of sowing date on yield potential of niger crop in rainfed condition. Plant Archives. 2014; 14(2): 995-997.
6. Mandal, MRI, Biswas MKP, Raman MM, Khealque MA. Effect of dates of sowing on the yield and yield component of niger. Bangladesh J Agric Sci. 1991; 18:153-155.
7. Mishra RC, Pradhan K, Paikray RK, Sahu PK, Panda BS. Seasonal adaptability of niger varieties. Oil Crops News Ietr. 1991; 8: 26-27.
8. Mohan Kumar BN, Basavegowda, Vyakaranahal BS, Deshpande VK, Kenchanagoudar PV. Influence of sowing dates on production of seed yield in niger (*Guizotia abyssinica* Cass.). Karnataka Journal of Agricultural Sciences. 2011; 24 (3): 289 -293.

Comment [A11]: Interesting information here again. All important information could be in the abstract. Remove all introduction of abstract and put interesting information to catch readers attention

Comment [A12]: This is a methodology, not a result

Comment [A13]: Great. According this results the author could create a scientific question. This question can be used as a way to improve interest of readers. For example: What is the best sowing period? which conditions or attributes influence the sowing of niger?

Comment [A14]: I missed a strong literature review and also a more current literature

9. Nayak BC, Paikray RC. Effect of time of sowing and plant density on yield of niger. Journal of Oilseeds Research. 1991; 8: 254- 258.
10. Indian Institute of Oil Seeds Research, IIOSR. Oilseeds Statistics - A Compendium, ICAR-Indian Institute of Oil Seeds Research, Rajendranagar, Hyderabad. 2015; 350-372 <https://krishi.icar.gov.in/jspui/bitstream/123456789/2820/1/COMPENDIUM%20-CONTENTS.pdf>
11. Paul, SR, Suhrawardy J, Guha B. Effect of different dates of sowing on growth and yield of niger under rainfed condition. Journal of Oilseeds Research. 1995; 12(1): 123-125.
12. Mili R, Gogoi B, Bora PC. Influence of time of sowing on yield and yield attributes of niger (*Guizotia abyssinica* L.) under rainfed situation of Assam. Agricultural Science Digest. 2012; 32(3): 263 – 265.
13. Singh PP, Singh RV, Singh MP, Jain D, Khanddait SL. Effect of sowing dates fertility level and plant density on the growth and seed yield of winter niger (*Guizotia abyssinica* Cass). Indian Journal of Applied and Pure Biology. 1990; 5(2):89-92.




Fig.1 Weather data during 2015

Comment [A15]: Need describe better your subtitle of figures. Enrich it with more informations (take from methodology)




Fig. 2 Weather data during 2016




Fig.3 Weather data during 2017




Fig.4 Influence of date of sowing, variety and their interaction on seed yield of niger

Table 1. Rainfall (mm) and dry spell (days) of the experimental station during the crop growth period

| Month | Rainfall (mm) | | | Normal rainfall (mm) | Dry spell (days) | | |
|-----------|---------------|-------|-------|----------------------|------------------|------|------|
| | 2015 | 2016 | 2017 | | 2015 | 2016 | 2017 |
| July | 64.3 | 410.7 | 378.7 | 375.6 | 21 | 10 | 10 |
| August | 202.6 | 305.0 | 284.5 | 393.6 | 18 | 14 | 15 |
| September | 477.8 | 310.1 | 259.6 | 256.3 | 19 | 11 | 16 |
| October | 33.2 | 55.2 | 266.0 | 126.1 | 30 | 25 | 19 |
| November | 18.8 | 3.0 | 18.0 | 32.6 | 30 | 29 | 28 |
| December | 10.2 | 0.0 | 0.0 | 6.5 | 30 | 31 | 31 |

Table 2. Meteorological information of Semiliguda centre during crop period growth in 2015

| Month | Tmin | Tmax | Rainfall (mm) | Sunshine hours (hr:mm) | Relative Humidity (%) | |
|-----------|------|------|---------------|------------------------|-----------------------|------|
| | | | | | Min | Max |
| July | 17.2 | 28.0 | 64.3 | 73.0 | 73.4 | 91.2 |
| August | 17.1 | 27.7 | 202.6 | 59.5 | 79.6 | 94.0 |
| September | 15.9 | 27.7 | 477.8 | 105.3 | 72.5 | 95.3 |
| October | 12.8 | 29.4 | 33.2 | 214.4 | 55.3 | 92.5 |
| November | 4.9 | 27.9 | 18.8 | 187.9 | 66.4 | 90.2 |
| December | 7.8 | 28.5 | 10.2 | 240.4 | 66.3 | 85.3 |

Note: Tmin= Minimum temperature; Tmax= Maximum temperature; hr= Hour

Table 3. Meteorological information of Semiliguda centre during crop period growth in 2016

| Month | Tmin | Tmax | Rainfall (mm) | Sunshine duration (hr:mm) | Relative Humidity (%) |
|-----------|------|------|---------------|---------------------------|-----------------------|
| July | 14.4 | 26.6 | 410.7 | 40.1 | 96.1 |
| August | 13.9 | 26.6 | 305.0 | 83.5 | 95.6 |
| September | 14.0 | 27.2 | 310.1 | 54.1 | 96.2 |
| October | 9.7 | 27.3 | 55.2 | 173.4 | 96.4 |
| November | 5.9 | 28.1 | 3.0 | 246.0 | 96.9 |
| December | 7.4 | 27.4 | 0.0 | 248.4 | 96.7 |

Note: Tmin= Minimum temperature; Tmax= Maximum temperature; hr= Hour

Table 4. Meteorological information of Semiliguda centre during crop growth period in 2017

| Month | Tmin | Tmax | Rainfall (mm) | Sunshine duration (hr:mm) | Relative Humidity (%) |
|-----------|------|------|---------------|---------------------------|-----------------------|
| July | 20.7 | 26.2 | 378.7 | 69.0 | 99.5 |
| August | 20.8 | 27.0 | 284.5 | 48.5 | 86.9 |
| September | 20.4 | 28.4 | 259.6 | 86.7 | 81.9 |
| October | 17.5 | 28.6 | 226.0 | 198.4 | 70.6 |
| November | 15.1 | 27.1 | 18.0 | 212.4 | 98.8 |
| December | 11.1 | 27.1 | 0.0 | 219.4 | 97.4 |

Note: Tmin= Minimum temperature; Tmax= Maximum temperature; hr= Hour

Table 5. Growth parameters and yield attributes of niger as influenced by date of sowing and variety (pooled data of 3 years)

| Treatment | Plant height (cm) | No. of branches per plant | Days to 50% flowering | Days to maturity | No. of capitula per plant | No. of seeds per capitula |
|----------------------------|-------------------|---------------------------|-----------------------|------------------|---------------------------|---------------------------|
| Date of sowing (D) | | | | | | |
| 15 th July | 225.7 | 7.8 | 73 | 127 | 67.6 | 29.5 |
| 30 th July | 202.6 | 7.0 | 65 | 122 | 58.0 | 26.0 |
| 14 th August | 170.3 | 6.3 | 55 | 112 | 47.3 | 25.2 |
| 29 th August | 145.3 | 5.6 | 49 | 99 | 35.6 | 21.3 |
| 13 th September | 106.5 | 5.3 | 42 | 93 | 30.6 | 18.8 |
| 28 th September | 68.0 | 4.5 | 43 | 90 | 17.0 | 16.5 |
| S.E (m) ± | 5.8 | 0.2 | 0.5 | 0.5 | 2.4 | 1.1 |
| CD (0.05) | 18.4 | 0.8 | 1.6 | 1.5 | 7.6 | 3.4 |
| Variety (V) | | | | | | |
| Deomali | 157.8 | 6.0 | 55 | 108 | 42.7 | 23.1 |
| Utkal Niger-150 | 148.4 | 6.1 | 54 | 106 | 42.7 | 22.6 |
| S.E (m) ± | 3.4 | 0.1 | 0.3 | 0.3 | 0.9 | 0.5 |
| CD (0.05) | 10.3 | 0.4 | 0.9 | 0.8 | 2.7 | 1.4 |
| Interaction (D X V) | | | | | | |
| S.E (m) ± | 8.2 | 0.3 | 0.7 | 0.7 | 2.1 | 1.1 |
| CD (0.05) | 25.4 | 0.9 | 2.1 | 2.0 | 6.5 | 3.4 |
| Interaction (V X D) | | | | | | |
| S.E (m) ± | 4.8 | 0.2 | 0.4 | 0.4 | 1.6 | 0.8 |
| CD (0.05) | 15.0 | 0.6 | 1.3 | 1.2 | 5.1 | 2.4 |

Comment [A16]: Even though it is understandable for scientists the meaning, it is recommended that these acronyms be described to the non-scientific community. Remembering that your manuscript is about production and can be read by producers who may not understand academic language.

Same to CD and interaction (including de meaning in the methodology)

Table 6. Seed yield (kg ha⁻¹) of niger as influenced by date of sowing and variety

| Treatment | Seed yield (kg ha ⁻¹) | | | |
|----------------------------|-----------------------------------|-------|-------|--------|
| | 2015 | 2016 | 2017 | Pooled |
| Date of sowing (D) | | | | |
| 15 th July | 519.0 | 671.2 | 498.4 | 562.9 |
| 30 th July | 334.1 | 716.0 | 438.8 | 496.3 |
| 14 th August | 296.8 | 673.6 | 459.2 | 476.5 |
| 29 th August | 166.7 | 576.8 | 467.7 | 403.7 |
| 13 th September | 117.2 | 240.0 | 424.9 | 260.7 |
| 28 th September | 113.9 | 42.4 | 227.9 | 128.1 |
| S.E (m) ± | 7.4 | 27.7 | 21.5 | 18.9 |
| CD (0.05) | 23.3 | 87.4 | 67.7 | 59.5 |
| Variety (V) | | | | |
| Deomali | 241.0 | 480.0 | 420.2 | 380.4 |
| Utkal Niger-150 | 274.9 | 493.3 | 419.0 | 395.7 |
| S.E (m) ± | 8.3 | 17.9 | 11.7 | 12.6 |
| CD (0.05) | 25.7 | 55.0 | 35.9 | 38.9 |
| Interaction (D X V) | | | | |
| S.E (m) ± | 20.4 | 43.7 | 28.6 | 30.9 |
| CD (0.05) | 62.9 | 134.8 | 88.0 | 95.2 |
| Interaction (V X D) | | | | |
| S.E (m) ± | 9.4 | 24.0 | 17.0 | 16.8 |
| CD (0.05) | 29.0 | 74.6 | 53.1 | 52.2 |

Table 7. Net Monetary Return and Benefit: Cost ratio as influenced by date of sowing and variety

| Treatment | 2015 | | 2016 | | 2017 | | Mean | |
|----------------------------|-------------------------------|--------------|-------------------------------|--------------|-------------------------------|--------------|-------------------------------|--------------|
| | NMR (Rs ha ⁻¹) | B:C Ratio | NMR (Rs ha ⁻¹) | B:C Ratio | NMR (Rs ha ⁻¹) | B:C Ratio | NMR (Rs ha ⁻¹) | B:C Ratio |
| Date of sowing (D) | | | | | | | | |
| 15 th July | 8697 | 1.7 | 9897 | 1.6 | 5655 | 1.4 | 8083 | 1.6 |
| 30 th July | 1300 | 1.1 | 11689 | 1.7 | 3252 | 1.2 | 5414 | 1.3 |
| 14 th August | -192 | 1.0 | 9993 | 1.6 | 4068 | 1.3 | 4623 | 1.3 |
| 29 th August | -5398 | 0.6 | 6121 | 1.4 | 4410 | 1.3 | 1711 | 1.1 |
| 13 th September | -7378 | 0.4 | -7351 | 0.6 | 2694 | 1.2 | -4012 | 0.7 |
| 28 th September | -7509 | 0.4 | -15255 | 0.1 | -5184 | 0.6 | -9316 | 0.4 |
| S.E (m) ± | 156.5 | 0.02 | 290.2 | 0.03 | 266.4 | 0.06 | 237.7 | 0.04 |
| CD (0.05) | 493.3 | 0.08 | 914.3 | 0.10 | 839.4 | 0.19 | 749.0 | 0.12 |
| Variety (V) | | | | | | | | |
| Deomali | -2423 | 0.8 | 2249 | 1.1 | 2507 | 1.2 | 778 | 1.0 |
| Utkal Niger-150 | -1070 | 0.9 | 2782 | 1.2 | 2459 | 1.2 | 1390 | 1.1 |
| S.E (m) ± | 104.1 | 0.03 | 157.6 | 0.02 | 96.4 | 0.03 | 119.4 | 0.03 |
| CD (0.05) | 320.7 | 0.08 | 485.7 | 0.05 | 296.9 | 0.10 | 367.8 | 0.08 |
| Interaction (D X V) | | | | | | | | |
| S.E (m) ± | 254.9 | 0.07 | 386.1 | 0.04 | 236.1 | 0.08 | 292.4 | 0.06 |
| CD (0.05) | 785.6 | 0.21 | 1189.6 | 0.13 | 727.5 | 0.25 | 900.9 | 0.20 |
| Interaction (V X D) | | | | | | | | |

| | | | | | | | | |
|-----------|-------|------|-------|------|-------|------|-------|------|
| S.E (m) ± | 137.9 | 0.03 | 230.0 | 0.03 | 181.5 | 0.05 | 183.1 | 0.04 |
| CD (0.05) | 428.9 | 0.10 | 717.3 | 0.08 | 568.4 | 0.15 | 571.5 | 0.11 |

Note: NMR= Net monetary return; B:C ratio= Benefit - cost ratio

UNDER PEER REVIEW