Original Research Article

Mitigation option for GHG emission from wetland rice cultivation.

ABSTRACT

The study on the effect of soil amendments and fertilizers on the mitigation of CH_4 and N_2O from rice field was conducted in the Wetland farm of Tamil Nadu Agricultural University farm, Coimbatore, Tamil Nadu, India during *rabi* season did reveal that use of Gypsum and Fly Ash along with recommended dose of fertilizer reduced the emission of methane whereas urea with neem treatment had reduced the N_2O emission from the conventional water logged rice field. Methane emission was found to peak during the panicle initiation stage while nitrous oxide emission was noted only during the maturity stage. The application of Fly Ash along with gypsum had cut down the methane emission to a greater extent at all the stages. Nitrous oxide emission was reduced with the application of slow release N fertilizer (neem treated urea).

Key words: Fly ash, Neem, Gypsum, GHG, Methane and Nitrous oxide.

INTRODUCTION:

The most predominant form of carbon reserve is the soils and the soil carbon content plays a major role in the fertility of the soil. In case of paddy cultivation, the soil has been kept under submergence leading to anaerobic condition which contributes towards higher methane emission and to a lesser extent nitrous oxide emission (Tiwari *et al.*, 2020). But the untimely application and faulty methods of nitrogenous fertilizer application leads to N_2O emission.

Comment [p1]: Mention should be made of other works

UNDER PEER REVIEW

Rice being the stable food of major Asian population, the rising demand for food commodities has caused increasing pressure on paddy cultivation. Paddy field emits approximately 11 per cent and 30 per cent of global agricultural Nitrous oxide and Methane emission, respectively (Gupta *et al.*, 2021). It is therefore, pertinent to develop technologies to reduce emission of GHGs from rice field.

Fly ash, an amorphous mixture contains high amount of silica, iron, manganese oxides, zinc and copper which could act as a good soil amendment to suppress methane emission (Yadav and Pandita, 2019). These elements enact to enhance the electron acceptors leading to depression in methanogenesis. Gypsum has very high concentration of electron acceptor like SO_4^{2-}; its application might be effective on reducing CH_4 emission during rice cultivation (Thakur and Solanki, 2021). Nitrification inhibitors such as neem coated urea could slow down the process of nitrification in soil which could reduce emission of N_2O as well as methane from soil by 10-15 per cent (Singh et al., 2019).

The research was done with the objective of developing a mitigation strategy to reduce the emission of CH_4 and N_2O from the rice fields with selected soil amendments like fly ash, gypsum and Neem Treated Urea (NTU).

MATERIALS AND METHODS:

The field experiment was conducted at wetland farm, Tamil Nadu Agricultural University, Coimbatore during *rabi* with CO(R) 51, a short duration rice variety and the experiment was laid out in randomized complete block design and replicated thrice.

Comment [p2]: A map of the study area must be presented

Crop husbandry:

Nursery was raised with a seed rate of 40 kg ha^{-1}. The seeds were soaked for 12 hrs and treated with *Azospirillum* and *Pseudomonas* at the rate of 200 g and 100 g per 10 kg of

Comment [p3]: How you calculated the seeding rate?

UNDER PEER REVIEW

seeds and kept in dark for another 24 hours. Twenty days old seedlings were transplanted with a spacing of 20 x 10 cm in the plots that were treated with the selected soil amendments and fertilizers according to the treatments. Submergence in the main field was maintained till 15 days prior to harvest.

Treatments:

The eight treatments studied were as follows:

T_1 : Urea + Murate Of Potash (MOP) + Single Super Phosphate (SSP) (150:50:50)

T_2 : Neem Treated Urea (NTU) + MOP + SSP

T_3 : Urea + MOP + SSP + Gypsum (500 kg ha^{-1})

T_4 : NTU+ MOP + SSP + Gypsum (500 kg ha^{-1})

T_5 : Urea + MOP + SSP + Fly Ash (20 tonnes ha^{-1})

T_6 : NTU + MOP + SSP + Fly Ash (20 tonnes ha^{-1})

T_7 : Urea + MOP + SSP + Gypsum (500 kg ha^{-1}) + Fly Ash (20 tonnes ha^{-1})

T_8 : NTU + MOP + SSP + Gypsum (500 kg ha^{-1}) + Fly Ash (20 tonnes ha^{-1})

Neem treated urea was prepared by blending urea with crushed neem seed or neem cake 20% by weight. Powdered neem cake to pass through 2mm sieve before mixing with urea and kept overnight before use. Phosphorus, Gypsum and Fly Ash were was applied as basal, Potassium and Nitrogen were applied in four equal splits as per the treatment at basal, active tillering, panicle initiation and 50 per cent flowering stages.

Gas collection:

Comment [p4]:) you have to explain how you calculated the different doses? SSP? GYPSUM and Fly Ash

Comment [p5]: THE CHARACTERISTICS OF THE SOIL MUST BE PROVIDED (pH, EC.....OF SOIL)

UNDER PEER REVIEW

Gas samples were collected from the field using static closed chamber technique. The gas chambers were fabricated as per the recommendations made from several studies (Denmead, 2008). Open-bottom perplex chambers using 4 mm acrylic sheets with a dimension of 50 cm x 50 cm x 100 cm were fabricated. A battery (12V) operated fan was fixed for air circulation (to avoid plant suffocation) to mix the air inside the chamber.

As described by Khosa *et al.*, (2010) each chamber was placed on the soil surface with 4-5 cm inserted into the soil. Care was taken not to disturb the vegetation during the whole measurement programme. After covering the plants with the chamber, four air samples were collected in Tedlar bags starting with zero time and subsequent sampling was done at an interval of 15 minutes using one way valve pump. As described by Jayadeva *et al.* (2009), the air samples were collected in the morning (09:00-10:00 hours) and in the evening (14:00-15:00 hours) and the average of morning and evening fluxes were used as the flux value for the day.

Gas estimation:

The CH_4 and N_2O were estimated in a Shimadzu GC-2014 gas chromatograph equipped with FID and ECD. The gas samples were introduced into the analyzer by filling the fixed loop (1.0 ml) on the sampling valve. Samples were injected into the column system by starting the analyzer which automatically activated the valve and back flush the samples according to the time programmed. The GC was calibrated before and after each set of measurements using 1ppm, 2.3ppm and 5ppm of standards (Chemtron® science laboratories Pvt. Ltd., Mumbai) as primary standard curve linear over the concentration ranges used. CH_4 and N_2O concentration were expressed as mg m^{-2} hr^{-1} using the equation given by Lantin *et al*., (1995). The obtained CH_4 and N_2O concentrations were determined by peak area and flux

UNDER PEER REVIEW

was calculated based the equation proposed by Rolston (1986) to estimate methane and nitrous oxide.

$$f = (V/A)\ (\Delta C/\ \Delta t)$$

Where f is equal to greenhouse gas emission rate (mg m^{-2} h^{-1}), V is equal to volume of chamber above soil (m^3), A is equal to cross-section of chamber (m^2), ΔC is equal to concentration difference between zero and t times (mg cm^{-3}), and Δt is equal to time duration between two sampling periods (h).

RESULTS AND DISCUSSION:

Methane emission

The emission strategies were quantified for the four growth stages (active tillering, panicle initiation, 50 per cent flowering and harvest stages) and data are presented in figure 1. Significantly higher methane emission was recorded in treatment with NPK alone (T_1 & T_2) at all the four stages. In the present investigation the treatment in which fly ash and gypsum were used along with RDF (T_8) did record lower CH4 emissions as compared to other treatments. Though application of gypsum and fly ash along with RDF (T_8) did reduce the methane emission significantly, from the results it was truly observed that recommended fertilizer and gypsum (T_3) alone stood next to the treatments with NPK alone (T_1) on the higher emission of methane. The reduction of methane with the application of fly ash and gypsum along with NTU, MOP and SSP (T_8) over control (T_1) was about 31.53, 24.46, 442 and 3.93 per cent at active tillering, panicle initiation, 50 per cent flowering and maturity stages respectively.

The application of fly ash reduced the methanogens population with the lowest number in the treatments T_5, T_6, T_7 and T_8 at all three stages. The methanogen population at active tillering stage in the treatments T_1 to T_4 i.e. the treatments without fly ash in the range between 11 to

UNDER PEER REVIEW

12 x 10^3 CFU per g of soil whereas it was 7 to 8 x 10^3 CFU per g in the treatments with fly ash as a component. In the panicle initiation stage the population was 18 or 19 x 10^3 CFU per g of soil in non fly ash applied treatments and 12 or 13 x 10^3 CFU per g of soil in fly ash treated fields. During the 50 per cent flowering stage, the population was found to be highest in T_2 (NTU+SSP+MOP+Gypsum), but was not statistically significant over other treatments. The range was between 12 to 15 x 10^3 CFU per g of soil in treatments without fly ash and 8 to 10 x 10^3 CFU per g of soil in treatments with fly ash. The study on methanogen population revealed that the population was lowest in the treatments T_5, T_6, T_7 and T_8 in which Fly Ash was a component at the three stages (active tillering, panicle initiation and 50 per cent flowering) (Pradipa *et al.,* 2016) and the lower emission was also noticed in the same treatments at all the four stages. The methanogen population was also in line with emission results.

Comment [p6]: WHAT MUST BE EXPLAINED a statistical test should be performed to confirm this.

The reduction in methanogen population inturn the methane emission with the increased sulfate concentration, iron and manganese might have resulted into proliferation of other reducing bacteria like Sulphate reducing, that out-compete methanogens for substrates leading to reduction in population of methanogens (Singh et al., 2018).Also increase ferric iron concentration increases the microbial ferric iron reduction to Ferrous iron leading to suppression of Methanogenesis (Gabriel et al., 2020). This might also be result due to shifting of electron flow from methanogenesis to sulfate reduction under anaerobic soil conditions (Conard *et al.,* 2020).

Either, Gypsum or fly ash when applied alone had lesser impact when compared with the treatments where both are applied in combination. Eventhough gypsum is a good source of sulphate, the rate of application was only 500 kg per ha which might not be sufficient enough

UNDER PEER REVIEW

to suppress the activity of methanogens (Pradipa *et al.*, 2016) inturn the methane emission. The suppression of CH_4 emission was due to the increased concentrations of active sulfate in the soil amended by gypsum and fly ash might have controlled methanogens activity by limiting substrates availability (Malyan *et al.*, 2021) and hence resulted in reduced methane emission. The increased supply of iron and manganese compounds from the fly ash might have acted as electron acceptors and thereby, suppressed CH_4 production as well as CH_4 emission during rice cultivation (Sapkota *et al.*, 2020). Application of NTU urea has no influence on methane emission from rice fields. The reduction in methane emission during the later stages was prominent that too in the treatments with fly ash and gypsum in combination with RDF, since the antagonistic microbial mass might need time for its proliferation and active involvement in electron transport chain.



Figure. 1. Influence of different treatments on methane emission (mg m^{-2} hr^{-1})

Nitrous oxide emission rate

In respect of nitrous oxide, upto 50 per cent of flowering the nitrous oxide emission was below detectable level (50 ppb) and was recorded only at the harvest stage since water was stopped 15 days before harvest (Figure. 2). Once the soil was brought to aerobic condition the nitrous oxide emission was found in the present investigation. At this stage, the data were found to be significant, wherein significantly higher nitrous oxide emission was noted in treatments T_7 and T_1.

In the treatments in which NTU was applied instead of urea showed a decline in N_2O emission as reported by Davamani *et al.*, (2021). The lowest nitrous oxide emission was noted with the treatment T_2 in which NTU was used as a slow nitrogen release fertilizer.

Comment [p7]: WHAT MUST BE EXPLAINED

The mean nitrous oxide emission during the maturity stage recorded was 0.5008 mg m^{-2} hr^{-1}. The high pitched nitrous oxide emission rate of 0.5940 mg m^{-2} hr^{-1} was with T_1 (control) followed by T_5 and the most modest value of 0.3996 mg m^{-2} hr^{-1} was recorded with the treatment T_2. The reduction was about 32.72 per cent.



Figure. 2. Influence of different treatments on nitrous oxide emission at maturity stage (mg $m^{-2}hr^{-1}$)

UNDER PEER REVIEW

CONCLUSION:

Based on the results it is concluded that fly ash or/and gypsum along with RDF reduces the emission of CH_4 by providing alternate electron acceptors whereas NTU, did reduce the N_2O emission by inhibiting nitrification. The amendments in combination ca have a good impact on emission of the two major GHG (i.e. CH_4 & N_2O) from the rice field.

Reference:

Conrad, R. (2020). Importance of hydrogenotrophic, aceticlastic and methylotrophic methanogenesis for methane production in terrestrial, aquatic and other anoxic environments: a mini review. Pedosphere, 30(1), 25-39.

Davamani, V., Poornima, R., Arulmani, S., Parameswari, E., John, J. E., & Deepasri, M. (2021). Mitigation of nitrous oxide emission through fertigation and 'N'inhibitors–A sustainable climatic crop cultivation in tomato. Science of The Total Environment, 152419.

Denmead O T. (2008). Approaches to measuring fluxes of methane and nitrous oxide between landscapes and the atmosphere. Plant Soil. 309: 5 - 24.

Gabriel, G. V., Oliveira, L. C., Barros, D. J., Bento, M. S., Neu, V., Toppa, R. H., ... & Navarrete, A. A. (2020). Methane emission suppression in flooded soil from Amazonia. Chemosphere, 250, 126263.

Gupta, K., Kumar, R., Baruah, K. K., Hazarika, S., Karmakar, S., & Bordoloi, N. (2021). Greenhouse gas emission from rice fields: a review from Indian context. Environmental Science and Pollution Research, 1-22.

UNDER PEER REVIEW

Jayadeva, H.M., Prabhakara Setty, T.K., Gowda, R.C., Devendra, R., Mallikarjun, G.B., and Bandi, A.G. (2009) "Methane emission as influenced by different crop establishment techniques and organic manures," Agricultural Science Digest, 29, 241-245.

Jyotsnarani Jena, Sanak Ray, Haragobinda Srichandan, Anuradha Das and Trupti Das. (2013). Role of Microorganisms in Emission of Nitrous Oxide and Methane in Pulse Cultivated Soil Under Laboratory Incubation Condition. Indian Journal of Microbiology. 53(1): 92–99.

Khosa M K, B S Sidhu and D K Benbi. (2010). Effect of organic materials and rice cultivars on methane emission from rice field. Journal of Environmental Biology. 31: 281 – 285.

Lantin R S, J B Aduna and A M J Javellana. (1995). Methane measurements in rice fields. Instruction manual and methodologies, maintenanace and troubleshooting guide. In: A joint undertaking by International Rice Research Institute (IRRI), United states Environmental Protection Agency (US-EPA) and United Nation Development Program (UNDP). 12-15.

Malyan, S. K., Kumar, S. S., Singh, A., Kumar, O., Gupta, D. K., Yadav, A. N., ... & Kumar, A. (2021). Understanding Methanogens, Methanotrophs, and Methane Emission in Rice Ecosystem. In Microbiomes and the Global Climate Change (pp. 205-224). Springer, Singapore.

Pradipa C, N. Maragatham, M. Maheswari, V. Geethalakshmi. (2016). Effect of fertilizers and soil amendments on Methanogen population in Rice field. In: Proceedings of National conference on Climate change impacts and Natural Resource Management. Anna University, Chennai

UNDER PEER REVIEW

Rolston D E. (1986). Gas Flux, in: Methods of soil analysis. In: Part 1, 2nd ed., Klute A. (Ed.)., Agronomy Monograph No. 9. ASA and SSSA, Madison, WI. 1103–1119.

Sapkota, A., Haghverdi, A., Avila, C. C., & Ying, S. C. (2020). Irrigation and greenhouse gas emissions: a review of field-based studies. Soil Systems, 4(2), 20.

Singh, A., Jaswal, A., & Singh, M. (2019). Impact of neem coated urea on rice yield and nutrient use efficiency (NUE). Agricultural Reviews, 40(1), 70-74.

Singh, N. K., Patel, D. B., & Khalekar, G. D. (2018). Methanogenesis and Methane Emission in Rice/Paddy Fields. In Sustainable Agriculture Reviews 33 (pp. 135-170). Springer, Cham.

Thakur, S., & Solanki, H. (2021). Role of Methane in Climate Change and Options for Mitigation-A Brief Review.

Tiwari, S., Singh, C., & Singh, J. S. (2020). Wetlands: a major natural source responsible for methane emission. In Restoration of Wetland Ecosystem: A Trajectory Towards a Sustainable Environment (pp. 59-74). Springer, Singapore.

Yadav, V. K., & Pandita, P. R. (2019). Fly ash properties and their applications as a soil ameliorant. In Amelioration Technology for Soil Sustainability (pp. 59-89). IGI Global.

Yusminah Hala, Oslan Jumadi, Abd. Muis, Hartati and Kazuyuki Inubushi. (2014). Development of Urea Coated with Neem (Azadirachta indica) to Increase Fertilizer Efficiency and Reduce Greenhouse Gases Emission. Jurnal Teknologi (Sciences & Engineering). 69:5, 11–15

UNDER PEER REVIEW