

Comparative Assessment of Soil Carbon Sequestration and Carbon dioxide emissions from Agroforestry Systems in Kogi East Nigeria

ABSTRACT

This study was conducted to assess the below ground carbon sequestration (soil carbon stock per unit land area) and carbon dioxide (CO₂) emissions from agroforestry systems (AFSs) in Kogi East (Ankpa, Dekina, Ofu, Olamaboro, and Omala local government areas) Nigeria. Stratified random sampling was used to select study locations of the agroforestry systems in Kogi East, Nigeria. Four AFSs were selected in each local government area (LGA) - this consisted majorly of smallholder farmer's farm with silvoarable systems in the region (4 communities per LGA, total of 20 communities). The selection criteria for AFS was based on farm size not less than 1 hectare. The results from the analysis revealed that highest soil carbon stock [C stock (Mg Cha-1)] was recorded from AFSs in Dekina (334.43 Mg Cha-1) while no significant difference in carbon stock was observed from the soils of AFSs in Ankpa, Ofu, Olamaboro, and Omala LGAs (69.01, 159.21, 142.58, 117.33 Mg Cha-1 respectively). Nonetheless, the soils from AFSs in Dekina and Ofu LGAs had highest CO₂ emissions (186.23 and 159.40 gCO₂ emitted/50g wet soil slice respectively) while the lowest CO₂ emissions (104.15 and 88.88 gCO₂ emitted/50g wet soil slice) were recorded from Ankpa and Omala LGAs respectively.

Keywords: Agroforestry, Climate change, Soil Carbon Sequestration, Soil Management.

1. INTRODUCTION

Carbon (C) sequestration can be described as the process of capturing atmospheric C and safely storing it in long-lived pools (United Nations Framework Convention on Climate Change- UNFCCC, 2007; Zheng *et al.*, 2020). Globally, carbon sequestration in terrestrial vegetation systems is recognised to have potential to mitigate the increasing levels of CO₂ in the atmosphere (Intergovernmental Panel on Climate Change - IPCC, 2007; Keenan and Williams, 2018; Lal *et al.*, 2018). On the other hand, agroforestry can be referred to as combination of agriculture (crops and/or livestock) and forestry (trees and shrubs) on the same land management unit (FAO, 2010; World Agroforestry Centre, 2012). Carbon sequestration in agroforestry involves the process of taking up atmospheric CO₂ during photosynthesis and the transfer of fixed C into vegetation, detritus, and soil pools for long-term storage (Nair *et al.*, 2010). Agroforestry can prevent the deliberate harmful circle of deforestation, soil erosion and other environmental problems in Nigeria (Sobola *et al.*, 2015; Meena *et al.*, 2020).

Carbon sequestration in agroforestry systems can be categorised into: 1) aboveground segment of trees and herbaceous parts like leaves, stems, etc. and 2) belowground segment comprising of the roots, C stored in different soil horizons, and soil organisms (Nair, 2011a). Under the same ecological conditions, the above and below ground C sequestration of agroforestry systems (woody perennial-based land use systems) are higher than monocultures of crops or pasture due to the ability of trees to absorb atmospheric carbon to store in their tissues and soils for a longer period of time (Kirby and Potvin, 2007; Jose, 2009; Rigueiro-Rodríguez *et al.*, 2009; Niether *et al.*, 2020). Soil organic matter and nutrient stocks in agroforestry systems are improved by the abundant and frequent addition of leaf litter and/or prunings including root biomass over a period of time which is vital for soil carbon dynamics (Albrecht and Kandji, 2003; Sileshi *et al.*, 2020)..

Agroforestry as a practice can create an integrated and sustainable land use systems (Mosquera-Losada *et al.*, 2009; Jose *et al.*, 2021). It can increase the productivity of land while applying management practices that are environmentally and socially acceptable

(Rigueiro-Rodríguez *et al.*, 2009; Sharma *et al.*, 2021). The environmental benefits of AFS include: 1) **soil quality improvement**: Agroforestry systems have improved nutrient cycling through leaf litter production and decomposition as well as deep nutrient capture by their root systems (the roots of the trees in agroforestry system are deep and strong accessing nutrients deeper in soil profile that are most times not available for monocultures) (Sileshi *et al.*, 2020), they enhance soil organic carbon and greater soil microbial dynamics compared to monocultures (Dollinger and Jose, 2018; Marsden *et al.*, 2020); 2) **climate change adaptation and mitigation**: for example soil carbon sequestration. Intergovernmental Panel on Climate Change (IPCC, 2000) posited that assuming a global implementation of agroforestry systems, about 1.1 to 2.2 Pg of carbon can be captured from the atmosphere globally over 50 years. This is reported to have a compensating effect on greenhouse gas (GHG) emissions (10 – 15 % reduction in CO₂ emissions annually) in terms of climate change adaptation and mitigation strategies (Albrecht and Kandji, 2003). Furthermore, it is projected that a C sequestration of 0.586 Tg C per year can be achieved by 2040 by converting 630 million ha of unproductive croplands and grasslands to agroforestry (IPCC, 2000).; 3) **water quality management**: agroforestry can reduce water contamination and eutrophication by reduction in the use of inputs such as fertilizers (nitrate and phosphate fertilizers), herbicides and pesticides (Rigueiro-Rodríguez *et al.*, 2009). Trees in agroforestry systems act as dispersion barriers to pest reducing the use of pesticides and herbicides. Also, the deep and strong rooting zones of trees in agroforestry systems uptake surplus nutrients that would, otherwise contaminate rivers. Furthermore, water quality can be protected by riparian buffer strips (strips of perennial vegetation - tree/shrub/grass) either natural or planted between croplands/pastures and water sources like streams, lakes, wetlands, and ponds to reduce non-point source pollution from agricultural lands (Schultz *et al.*, 2004; Mosquera-Losada *et al.*, 2009; Russell *et al.*, 2010). The riparian buffer strips will help decrease sediment and nutrient load from soil erosion, and also filter surface water and groundwater runoff (Dosskey, 2001; Lee *et al.*, 2003); and 4) **conservation of biodiversity**: AFS provides habitat for biodiversity to live and breed (Harvey and Gonzalez-

Comment [DM2]:

Comment [DM3]:

Comment [DM4]: ,

Villalobos, 2007; Jose, 2009; Cornell and Miller, 2011, Nair, 2011b). The combination of mulching and shading effects created by trees in an agroforestry system helps to improve the microclimatic conditions (temperature, water vapour content of air and wind speed) which lowers soil surface temperature as well as reduced rates of evaporation of soil moisture. This modified microclimatic conditions have beneficial roles on the system such as enhancing biodiversity and animal well-being, improved soil quality, pest and disease control (Briggs, 2012; Jose, 2019).

In terms of socioeconomic benefits, agroforestry systems are source of nutrition as well as additional income for farmers engaged in it. The farmers are gainfully employed with reduced level of poverty and improved standard of living (IFAD, 2011). The diversification of farm outputs in an agroforestry system is helpful in the reduction of risks from total crop failure compared to monoculture system in periods of extreme weather events including floods and droughts (Cornell and Miller, 2011). In addition to production of food crops, agroforestry systems provide different products such as fuel wood, timber, fruits, nuts, fibre, fodder and forage, gums and resins, hatching and hedging materials, gardening materials, craft products, medicinal products, and shade for animals and farm workers including recreation. Socioeconomic development (diversification of rural economies, skills, and products) can be sustained by the sales these timber and non-timber products by the farmers (Cornell and Miller, 2011, World Agroforestry Centre, 2012).

The aim of this study was to assess soil carbon sink of agroforestry systems in Kogi East Nigeria. This can provide insights on the possible contribution of AFS as an adaptation measure to climate change impacts in Kogi East, Nigeria.

2. Methodology

2.1 Study Location

Four agroforestry systems were selected from five local government areas (Ankpa, Dekina, Ofu, Olamaboro, and Omala) within Kogi East, Nigeria. This consisted majorly of smallholder farmer's farm with silvoarable systems in the region (4 communities per LGA, total of 20 communities) (Table 1). The communities within Ankpa LGA were Odagbo, Oje Elanyi, Ojogobi Olaji, and Okaba. Dekina LGA communities were Anyigba, Dekina, Egume, and Odu Ogbaloto. Ofu LGA communities were Ogbulu, Ugwolawo, Ejule, and Ochadamu. Olamaboro LGA communities were Ejoka, Igoti Ade, Unobe, and Ubalu while Omala LGA were Ajedibo, Ajomakoji, Odumukpo, and Okugba.

Table 1: Description of Study Locations

Local Government Area	Location of Agroforestry System	Vegetation/Cultivated Crops (Agroforestry System)	Coordinate		Topography
			Latitude	Longitude	
Ankpa	Odagbo	Trees: Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>), Teak tree (<i>Tectona grandis</i>), and African Locust Bean (<i>Parkia biglobosa</i>). Crops: Cassava (<i>Manihot esculenta</i>) and Maize (<i>Zea mays</i>).	7°47'05"N	7°73'55"E	Undulating
	Oje Elanyi	Trees: Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>), Mango (<i>Mangifera indica</i>), Mahogany (<i>Swietenia</i>), Iron Tree/Prosopis africana (Guill., Perrott, and Rich.) (Taub.), and African Locust Bean (<i>Parkia biglobosa</i>). Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>), and Groundnut (<i>Arachis hypogaea</i>).	7°36'25"N	7°62'37"E	Nearly flat
	Ojogobi Olaji	Trees: Oil Palm (<i>Elaeis guineensis</i>), and African Locust Bean (<i>Parkia biglobosa</i>). Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>), and Groundnut (<i>Arachis hypogaea</i>).	7°18'63"N	7°57'54"E	Nearly flat
	Okaba	Trees: Oil Palm (<i>Elaeis guineensis</i>), Kolanut tree (<i>cola nitida</i>), Iron Tree/Prosopis africana (Guill., Perrott, and Rich.) (Taub.), Teak tree (<i>Tectona grandis</i>), and African Locust Bean (<i>Parkia biglobosa</i>). Crops: Cassava (<i>Manihot esculenta</i>), and Maize (<i>Zea mays</i>).	7°46'94"N	7°73'92"E	Gentle undulating
Dekina	Ayingba	Trees: African Locust Bean (<i>Parkia biglobosa</i>), Oil Palm (<i>Elaeis guineensis</i>), Teak tree (<i>Tectona grandis</i>). Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>), Yam (<i>Dioscorea spp</i>).	7°29'10"N	7°11'32"E	Nearly flat
	Dekina	Trees: Cashew (<i>Anacardium occidentale</i>), Mango (<i>Mangifera indica</i>), African Locust Bean (<i>Parkia biglobosa</i>), and Iron Tree/Prosopis africana (Guill., Perrott, and Rich.) (Taub.) Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Soybean (<i>Glycine max</i>)	7°41'13"N	7°12'10"E	Lower slope
	Egume	Trees: Cashew (<i>Anacardium occidentale</i>), Plantain (<i>Musa x paradisiaca</i>), African Locust Bean (<i>Parkia biglobosa</i>), and Iron Tree/Prosopis africana (Guill., Perrott, and Rich.) (Taub.) Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>)	7°28'45"N	7°12'10"E	Undulating
	Odu Ogbaloto	Trees: Oil Palm (<i>Elaeis guineensis</i>), Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Yam (<i>Dioscorea spp</i>).	7°29'28"N	7°10'15"E	Undulating
Ofu	Ogbulu	Trees: Cashew (<i>Anacardium occidentale</i>), Oil Palm (<i>Elaeis guineensis</i>), African Locust Bean (<i>Parkia biglobosa</i>), Mango (<i>Mangifera indica</i>). Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Cowpea (<i>Vigna unguiculata</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>).	7°23'22"N	7°3'20"E	Nearly flat
	Ugwolawo	Trees: Teak tree (<i>Tectona grandis</i>), Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>), Mango (<i>Mangifera indica</i>), Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Groundnut (<i>Arachis hypogaea</i>), Okra (<i>Abelmoschus esculentus</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>).	7°23'22"N	7°3'20"E	Nearly flat
	Ejule	Trees: Teak tree (<i>Tectona grandis</i>), Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>).	7°23'22"N	7°3'20"E	Flat

		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Okra (<i>Abelmoschus esculentus</i>), Cowpea (<i>Vigna unguiculata</i>).			
	Ochadamu	Trees: Neem tree (<i>Azadirachta indica</i>), and Oil Palm (<i>Elaeis guineensis</i>).	7°23'37"N	7°2'7"E	Undulating
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>).			
Olamaboro	Ejoka	Trees: Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>), Mahogany (<i>Swietenia</i>), Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.), and African Locust Bean (<i>Parkia biglobosa</i>).	7°31'68"N	7°62'67"E	Nearly flat
		Crops: Cassava (<i>Manihot esculenta</i>) and Maize (<i>Zea mays</i>).			
	Igoti Ade	Trees: Oil Palm (<i>Elaeis guineensis</i>), Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.), Plantain (<i>Musa x paradisiaca</i>), Wild mango/Ogbono (<i>Irvingia gabonensis</i>), and African Locust Bean (<i>Parkia biglobosa</i>).	7°24'05"N	7°59'10"E	Nearly flat
		Crops: Cassava (<i>Manihot esculenta</i>), Yam (<i>Dioscorea spp</i>), and Maize (<i>Zea mays</i>).			
	Unobe	Trees: Cashew (<i>Anacardium occidentale</i>), Teak tree (<i>Tectona grandis</i>), Oil Palm (<i>Elaeis guineensis</i>), Plantain (<i>Musa x paradisiaca</i>).	7°23'22"N	7°3'20"E	Nearly flat
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Okra (<i>Abelmoschus esculentus</i>).			
	Ubalu	Trees: Cashew (<i>Anacardium occidentale</i>), Oil Palm (<i>Elaeis guineensis</i>), Plantain (<i>Musa x paradisiaca</i>), and Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.).	7°23'22"N	7°3'20"E	Undulating
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Yam (<i>Dioscorea spp</i>), Pigeon pea (<i>Cajanus cajan</i>).			
Omala	Ajedibo	Trees: Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.) and African Locust Bean (<i>Parkia biglobosa</i>).	7°74'58"N	7°61'04"E	Undulating
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), guinea corn (<i>Sorghum bicolor</i>), and Pigeon pea (<i>Cajanus cajan</i>).			
	Ajomakoji	Trees: Oil Palm (<i>Elaeis guineensis</i>), Cashew (<i>Anacardium occidentale</i>), Mango (<i>Mangifera indica</i>), Mahogany (<i>Swietenia</i>), Teak tree (<i>Tectona grandis</i>), Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.), and African Locust Bean (<i>Parkia biglobosa</i>).	7°91'12"N	7°51'62"E	Nearly flat
		Crops: Cassava (<i>Manihot esculenta</i>) and Maize (<i>Zea mays</i>).			
	Odumukpo	Trees: Teak tree (<i>Tectona grandis</i>) and African Locust Bean (<i>Parkia biglobosa</i>).	7°54'35"N	7°30'89"E	Nearly flat
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), and Yam (<i>Dioscorea spp</i>).			
	Okugba	Trees: Oil Palm (<i>Elaeis guineensis</i>), Mahogany (<i>Swietenia</i>), Iron Tree/ <i>Prosopis africana</i> (Guill., Perrott, and Rich.) (Taub.), Plantain (<i>Musa x paradisiaca</i>), and African Locust Bean (<i>Parkia biglobosa</i>).	7°43'82"N	7°36'99"E	Undulating
		Crops: Cassava (<i>Manihot esculenta</i>), Maize (<i>Zea mays</i>), Egusi/Melon (<i>Cucumeropsis mannii</i>), and Pigeon pea (<i>Cajanus cajan</i>).			

2.2 Sample Size and Sampling Techniques

Stratified sampling was used to select study locations that gave a good representation of the AFSs in Kogi East Nigeria. The selection of agroforests was based on farm size not less than 1 hectare. In each Local Government Area (LGA), four (4) AFSs were selected from four communities (1 AFS per community, total of 20 agroforestry systems from 5 LGAs) were selected for the study.

2.3 Soil Sampling and Analysis

Random soil sampling technique was used to collect surface soil samples at 0-15 cm depth from each of the AFS in the selected farms in study locations. A total of 400 samples (20 samples per community, 80 per LGA) were collected and bulked to 200 composite samples (10 samples per community, 40 per LGA) for soil carbon stock and carbon dioxide emissions. The soils were prepared (air-dried, crushed, **grinded**, and passed through a 2mm sieve and **material** larger than 2mm were discarded). Soil samples for carbon dioxide analysis were taken at 0 - 15 cm depth using a tube soil auger and transferred into zip lock bags on the field so as to preserve samples from contamination and drying.

Comment [DM5]: ground

Comment [DM6]:

Comment [DM7]:

2.4 Determination of Soil Carbon Stock per Unit Land Area

Nair *et al.* (2011a) reported that analysis of C content in the soil (mass per unit mass of soil, for example g C per 100 g soil) is the most common method for calculating the amount of C sequestered in soils. Soil Organic Carbon (SOC) Stocks at fixed depth (0-15 cm) was determined using the formula from Carter and Gregorich (2007):

$$SOC_{FD} = \sum_1^n D_{cs} C_{cs} L_{cs} \times 0.1$$

where SOC_{FD} is the SOC stock to a fixed depth (Mg Cha⁻¹ to the specified depth), D_{cs} is the density of core segment (g cm⁻³), C_{cs} is the organic C concentration of core segment (mg C g⁻¹ dry soil), and L_{cs} is the length of core segment (cm). Soil organic carbon concentration

was determined using the Walkley-Black wet oxidation method. The method involved the oxidation of organic carbon (OC) with dichromate and tetraoxosulphate VI (H₂SO₄); the residual dichromate was titrated against ferrous sulphate (Walkley and Black, 1934).

Comment [DM8]: sulphuric acid

Comment [DM9R8]:

Comment [DM10]:

2.5 Determination of Carbon dioxide Emissions

Comment [DM11]: please state reference

Comment [DM12]:

The reagents and equipment used for the study include: 0.5 M sodium hydroxide (NaOH), 0.2 M hydrochloric (HCl) acid, 0.4 M Barium chloride (BaCl₂), Phenolphthalein indicator, 125 ml conical flasks, Burettes and Respiration flasks (1 litre air tight sealable Agee jars). In the laboratory, 50 g each of soil sample were placed in pre-weighed Agee jars. The weight of each soil sample and Agee jar was weighed so as to obtain the wet weight of the soil slice. 10 ml of 0.5M of sodium hydroxide (NaOH) solution was dispensed into 125 ml conical flask and placed inside each of the Agee jar containing the soil samples. A control made up of three blank Agee jars containing 125 ml conical flask of NaOH with no soil was set up. The lids of all the jars were screwed tightly and kept to incubate for fourteen days. The Agee jars were ventilated every three days for two minutes. On the fourteenth day, the conical flasks were removed and the amount of CO₂ produced were analysed by volumetric titration. 4 ml of 0.5M NaOH trapping solution from the control jar was pipetted into a 50 ml conical flask and 10ml of 0.4M barium chloride was added to the content of the flask followed by 4 drops of phenolphthalein indicator which now gives the content of the flask a yellow coloration. This was titrated with 0.2M hydrochloric acid solution until a colourless solution was obtained (end point). The volume of HCL acid used in the titration process was read from the burette and noted. This procedure was repeated for the other two blanks and for the trapping solution used in the other jars.

The carbon dioxide emitted per gram of wet soil slice (gCO₂ emitted/g wet soil slice) was computed as=

$$\frac{\text{moles of NaOH reacted with CO}_2 \times 44\text{g}}{2}$$

2

2.6 Statistical Analysis

All measured variables was subjected to descriptive statistics (mean and standard deviation). Analysis of variance (ANOVA) was carried out on measured variables using GENSTAT Discovery Software while treatment means were separated using Duncan Multiple Range Test (DMRT) at $\leq 5\%$ probability level.

Comment [DM13]: were

3. Results and Discussions

3.1 Carbon Stock of Soils from Agroforestry Systems in Kogi East, Nigeria

The highest carbon stock was recorded from the soils from agroforestry systems in Dekina ($334.43 \text{ Mg Cha}^{-1}$) while no significant difference in carbon stock was observed from the soils of AFS in Ankpa, Ofu, Olamaboro, and Omala LGAs ($69.01, 159.21, 142.58, 117.33 \text{ Mg Cha}^{-1}$ respectively) (Table 2). Although previous studies such as Nair *et al.* (2009); and Nair (2011a) posited that factors that can influence the total amount of carbon sequestered include previous land use, tree species and density (broadleaves are higher sequesters compared to coniferous and deciduous trees), the type of agroforestry system (nature of components), age of perennials like trees (mature stands of trees have the capacity to storage more carbon compared to young stands), ecological region. Furthermore, soil organic carbon sequestration depends primarily on the soil C input and soil stabilization processes. Plant root and rhizosphere inputs, in particular, make a large contribution to SOC (Terrer *et al.*, 2021).

Table 2: Carbon Stock of Soils from Agroforestry Systems in Kogi East, Nigeria

Local Government Area	C stock per hectare (Mg Cha-1)	Statistics		
		Max	Min	SEM
		531.00	56.92	24.98
Ankpa	69.01b			
Dekina	334.43a			
Ofu	159.21b			
Olamaboro	142.58b			
Omala	117.33b			

Note: Means in a column with different letters are statistically significant at probability level of 5 % ($p = 0.05$), Max= Maximum, Min = Minimum, Mg Cha-1 = Mega gram carbon per hectare, and CO_2 = Carbon dioxide, and gCO_2 = grams of carbon dioxide.

3.2 Carbon dioxide emissions of Soils from Agroforestry Systems in Kogi East, Nigeria

The soils from agroforestry systems in Dekina and Ofu LGAs had highest carbon dioxide emissions (186.23 and 159.40 gCO₂ emitted/50g wet soil slice respectively) (Table 3). The lowest CO₂ (104.15 and 88.88 gCO₂ emitted/50g wet soil slice) were recorded from the soils of Ankpa and Omala LGAs respectively. The absence of variation in CO₂ emission levels in some of the locations studied can be attributed to similar land management practices like tillage, bush burning and soil fertility management. This could contribute to increase or decrease in carbon emissions as well as soil organic carbon (Reang *et al.*, 2021).

Comment [DM14]:

Comment [DM15]: (elucidate a reason)

Table 3: Carbon dioxide emissions of Soils from Agroforestry Systems in Kogi East, Nigeria

Local Government Area	Carbon dioxide Emissions (gCO ₂ emitted/50g wet soil slice)	Statistics		
		Max	Min	SEM
		195.80	84.04	9.62
Ankpa	104.15cd			
Dekina	186.23a			
Ofu	159.40ab			
Olamaboro	138.51bc			
Omala	88.88d			

Note: Means in a column with different letters are statistically significant at probability level of 5 % (p = 0.05), Max= Maximum, Min = Minimum, CO₂ = Carbon dioxide, and gCO₂ = grams of carbon dioxide.

4. Summary and Conclusions

This study was conducted to assess the below ground carbon sequestration- C stock per unit land area (Mg C ha^{-1}) and carbon dioxide emissions (gCO_2 emitted/50g wet soil slice) of agroforestry systems in Kogi East (Ankpa, Dekina, Ofu, Olamaboro, and Omala) Nigeria. Stratified random sampling was used to select study locations that gave a good representation of the AFS in the Kogi East Nigeria. The results from the analysis revealed that highest carbon stock was recorded from the soils of AFSs in Dekina while no significant difference in carbon stock was observed from the soils of AFSs in Ankpa, Ofu, Olamaboro, and Omala LGAs. Furthermore, the results further indicated that the soils from AFSs in Dekina and Ofu LGAs had highest CO_2 emissions while the lowest CO_2 emissions were recorded from Ankpa and Omala LGAs respectively.

References

- Albrecht, A., and Kandji, S.T. (2003). Carbon sequestration in tropical agroforestry systems. *Agriculture, Ecosystems and Environment*, 99, 15–27.
- Briggs, S. (2012). Agroforestry: a new approach to increasing farm production. A Nuffield Farming Scholarships Trust report sponsored by the NFU Mutual Charitable Trust. Pp.82.
- Carter, M. R., and Gregorich, E. G. (2007). Soil sampling and methods of analysis. CRC press.
- Cornell, J.D., Miller, M. (2011). "Agroforestry." In: Encyclopedia of Earth. Eds.Cutler J. Cleveland (Washington, D.C.: Environmental Information Coalition, National Council for Science and the Environment). Retrieved from <http://www.eoearth.org/article/Agroforestry>
- Dollinger, J., and Jose, S. (2018). Agroforestry for soil health. *Agroforestry systems*, 92(2), 213-219.
- Dosskey, M.G. (2001). Toward quantifying water pollution abatement in response to installing buffers on crop land. *Environmental Management*, 28(5), 577-598.
- FAO (2010). Climate smart agriculture: policies, practices and financing for food security, adaptation and mitigation, Rome.
- Harvey, C.A. and Gonzalez-Villalobos, J.A. (2007). Agroforestry systems conserve species-rich but modified assemblages of tropical birds and bats. *Biodiversity and Conservation*, 16, 2257-2292.
- IFAD (2011). Addressing climate change in west and central Africa: forest management in Cameroon, the Democratic Republic of the Congo, Gabon and Nigeria. International Fund for Agricultural Development, Rome.
- Intergovernmental Panel on Climate Change (IPCC, 2000). Land use, land-use change, and forestry. A special report of the IPCC. In: Watson RT, Noble IR, Bolin B, Ravindranath NH, Verardo DJ, Dokken DJ (eds.). Cambridge University Press, Cambridge, 375 Pp.
- ~~Intergovernmental Panel on Climate Change (IPCC, 2000). Land use, land-use change, and forestry. A special report of the IPCC. In: Watson RT, Noble IR, Bolin B, Ravindranath NH, Verardo DJ, Dokken DJ (eds.). Cambridge University Press, Cambridge, 375 Pp.~~
- Intergovernmental Panel on Climate Change (IPCC, 2007). Intergovernmental panel on climate change 2007. Synthesis report. Retrieved from http://www.ipcc.ch/pdf/assessment-report/ar4/syr/ar4_syr.pdf.

- Jose, S. (2009). Agroforestry for ecosystem services and environmental benefits: an overview. *Agroforestry Systems* 76(1), 1-10.
- Jose, S. (2019). Environmental impacts and benefits of agroforestry. In Oxford Research Encyclopedia of Environmental Science.
- Jose, S., Garrett, H. E. G., Gold, M. A., Lassoie, J. P., Buck, L. E., and Current, D. (2021). Agroforestry as an integrated, multifunctional land use management strategy. *North American Agroforestry*, 1-25.
- Keenan, T. F., and Williams, C. A. (2018). The terrestrial carbon sink. *Annual Review of Environment and Resources*, 43(1), 219-243.
- Kirby, K.R., Potvin, C. (2007). Variation in carbon storage among tree species: implications for the management of a small-scale carbon sink project. *For Ecol Manag* 246, 208–221.
- Lal, R., Smith, P., Jungkunst, H. F., Mitsch, W. J., Lehmann, J., Nair, P. R., .. and Ravindranath, N. H. (2018). The carbon sequestration potential of terrestrial ecosystems. *Journal of Soil and Water Conservation*, 73(6), 145A-152A.
- Lee, K.H., Isenhardt, T.M., and Schultz, R.C. (2003). Sediment and nutrient removal in an established multi-species riparian buffer. *Journal of Soil and Water Conservation*, 58, 1-8.
- Marsden, C., Martin-Chave, A., Cortet, J., Hedde, M., and Capowiez, Y. (2020). How agroforestry systems influence soil fauna and their functions-a review. *Plant and Soil*, 453(1), 29-44.
- Meena, R. K., Meena, M. L., Meena, B. L., Meena, H., and Sharma, Y. K. (2020). The role of agroforestry in ecological sustainability. *AGRICULTURE and FOOD e-NEWSLETTER*, 30.
- Mosquera-Losada, M.R., McAdam, J.H., Romero-Franco, R., Santiago-Freijanes, J.J., and Rigueiro-Rodríguez, A. (2009). Chapter 1: Definitions and Components of Agroforestry Practices in Europe. In: Rigueiro-Rodríguez, A., McAdam, J., and Mosquera-Losada, M. R. (eds.) *Agroforestry in Europe: Current Status and Future Prospects*. *Advances in Agroforestry Book-Series*. Springer 6, 3-19.
- Nair, P.K.R., Kumar, B.M. and Nair, V.D. (2009). Agroforestry as a strategy for carbon sequestration. *Journal of Plant Nutrition and Soil Science*, 172(1), 10-23.
- Nair P.K.R., Nair, V.D., Kumar, B.M., Showalter, J.M. (2010). Carbon sequestration in agroforestry systems. *Adv Agron* 108, 237–307.
- Nair, P.K.R. (2011a). Methodological Challenges in Estimating Carbon Sequestration Potential of Agroforestry Systems. In: Kumar, B.M., and Nair, P.K.R. (eds.) *Carbon Sequestration Potential of Agroforestry Systems: Opportunities and*

- Challenges. *Advances in Agroforestry*, 8,3-16. Springer Dordrecht Heidelberg London New York.
- Nair, P. K. R. (2011b). Agroforestry systems and environmental quality: Introduction. *Journal of Environmental Quality*, 40, 784–790.
- Niether, W., Jacobi, J., Blaser, W. J., Andres, C., and Armengot, L. (2020). Cocoa agroforestry systems versus monocultures: a multi-dimensional meta-analysis. *Environmental Research Letters*, 15(10), 104085.
- Reang, D., Hazarika, A., Sileshi, G. W., Pandey, R., Das, A. K., and Nath, A.J. (2021). Assessing tree diversity and carbon storage during land use transitioning from shifting cultivation to indigenous agroforestry systems: Implications for REDD+ initiatives. *Journal of Environmental Management*, 298, 113470.
- Rigueiro-Rodríguez, A., Fernández-Núñez, E., González-Hernández, P., McAdam, J.H. and Mosquera-Losada, M.R. (2009). Chapter 3 Agroforestry Systems in Europe: Productive, Ecological and Social Perspectives. In: Rigueiro-Rodríguez, A., McAdam, J., and Mosquera-Losada, M. R. (eds.) *Agroforestry in Europe: Current Status and Future Prospects*. Advances in Agroforestry Book-Series. Springer 6, 43-66.
- Russell, D., R. Asare, J.P. Brosius, R. Witter, M. Welch-Devine, K. Spainhower, and Barr., R. (2010). People, Trees and Parks: Is Agroforestry In or Out? *Journal of Sustainable Forestry*, 29, 1-26.
- Schultz RC, Isenhardt TM, Simpkins WW, Colletti JP (2004). Riparian forest buffers in agroecosystems- lessons learned from the Bear Creek Watershed, central Iowa, USA. *Agroforestry Systems*, 61–62(1–3), 35–50.
- Sharma, M., Kaushal, R., Kaushik, P., and Ramakrishna, S. (2021). Carbon Farming: Prospects and Challenges. *Sustainability*, 13(19), 11122.
- Sileshi, G. W., Mafongoya, P. L., and Nath, A. J. (2020). Agroforestry systems for improving nutrient recycling and soil fertility on degraded lands. In *Agroforestry for Degraded Landscapes* (pp. 225- 253). Springer, Singapore.
- Sobola, O. O., Amadi, D. C., and Jamala, G. Y. (2015). The Role of Agroforestry in Environmental Sustainability. *Journal of Agriculture and Veterinary Science (IOSR-JAVS)* 8 (5), 1: 20-25.
- Terrer, C., Phillips, R. P., Hungate, B. A., Rosende, J., Pett-Ridge, J., Craig, M. E., and Jackson, R. B. (2021). A trade-off between plant and soil carbon storage under elevated CO₂. *Nature*, 591(7851), 599-603.

- United Nations Framework Convention on Climate Change (UNFCCC, 2007). Report of the conference of parties on its thirteenth session, Bali, Indonesia. United Nations Framework Convention on Climate Change, Geneva.
- Walkley, A. and Black, C. A. (1934). An examination of the methods for determining soil organic matter and proposed modification of the chromic acid titration method. *Soil Science*. 27: 29-38.
- World Agroforestry Centre (2012). "People and Agroforestry." *Trees On Farms: 2011 International Year of Forests Agroforestry*. Retrieved from <http://treesonfarms.com/agroforestry/people-and-agroforestry>
- Zheng, J., Chong, Z. R., Qureshi, M. F., and Linga, P. (2020). Carbon dioxide sequestration via gas hydrates: a potential pathway toward decarbonization. *Energy and Fuels*, 34(9), 10529-10546.