

**Effect of foliar application of micronutrients on growth, yield and quality of tomato
(*Solanum lycopersicum* L.) cv. Arka Samrat.**

ABSTRACT

The present investigation was carried out with title “**Effect of foliar application of micronutrients on growth, yield and quality of tomato (*Solanum lycopersicum* L.) cv. Arka Samrat**” at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh during the *Rabi*-2021-22 with a view to determine the effect of foliar application of micronutrients on tomato variety ‘Arka Samrat’ for its growth, quality and yield and to work out the economics of various treatments. Under this experiment, overall 8 treatment was taken T₁ Control (water spray), T₂ FeSO₄ @ 0.2% spray, T₃ Ca(NO₃)₂ @ 0.2% spray, T₄ H₃BO₃@0.1% spray, T₅ ZnSO₄ @ 0.2% spray, T₆ FeSO₄ @ 0.2% + Ca(NO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2% spray, T₇ Ca(NO₃)₂ @ 0.2% + ZnSO₄ @ 0.2% spray, T₈ Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1% spray. From the above experimental finding it may be concluded that the treatment T₆ (FeSO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) was found to be best in the terms of growth Yield and quality of tomato. While, the maximum Plant height (167.85 cm), highest No. of leaves per plant(204.33), Minimum Days to 50% flowering(37.33 DAS),Maximum No. of flower per cluster(5.12), Maximum No. of fruit per plant(60.33) and maximum Fruit set per cluster(5.06), Maximum fruit weight(94.22g), Maximum average yield per plant(5.68kg), Maximum average yield per plot(34.10), Maximum average yield per hectare(189.43t/ha),_Maximum Total soluble solid_(5.23),_Maximum Ascorbic Acid_(26.11).

Keywords: Foliar application, micronutrients, H₃BO₃.

Introduction

Tomato, botanically known as *Solanum lycopersicum* L. or *Lycopersicon esculentum* Mill. It is one of the most popular and widely grown vegetable crops throughout the world and is treated as “protective food” universally. It is rich source of vitamins, vegetable protein and minerals and holds a glorious position among vegetable after the potato and sweet potato. Tomato is known as the poor man’s apple (orange) in India & love apple in England. Tomato is used as soup, salad, pickles, ketchup, puree, sauces, tomato paste, tomato juice and other products. The pulp and juice of tomato fruit are digestible and a mild aperients, a promoter of gastric secretion and a blood purifier.

Tomatoes are a horticulture crop belongs to the family *Solanaceae* bearing chromosome number $2n=2X=24$ (Karpechenko, 1925). It originated from in South America (Vavilov, 1935). The tomato plants typically grow to 1–3 meters (3–10 ft) in height and have a weak stem that often sprawls over the ground and vines over other plants. Flowers are generally borne in clusters of 4 to 8 but small-fruited types may have 30 to 50 flowers per cluster. Tomato plants are dicots, and grow as a series of branching stems, with a terminal bud at the tip that does the actual growing. Tomato plays a major role in human nutrition, fruit contain 93.1% water, 1.9% protein, 0.3 g fat, 0.7% fibre, 3.6% carbohydrates, 23 calorie, 320 IU vitamin A., 0.07 mg vitamin B1, 0.01 mg vitamin B2, 31 mg vitamin C, 20 mg calcium, 36 mg phosphorus and 0.8 mg iron. Tomato has valuable vitamins and cholesterol. Approximately 20–50 mg of lycopene per 100g of fruit weight can be obtained from tomato. Tomato is a warm-season crop. The best fruit colour and quality is are

obtained at a temperature range of 21–24°C. Tomato is one of the most versatile crops in the world because of its fast and wide climate adaption and it is universally treated as protective food. Tomato contributes to a healthy, well-balanced diet. They are rich in minerals, vitamins, essential amino acids, sugars, dietary fibres and it has many other uses tomato seed contains 24% oil and is used as a salad oil and in the manufacture of margarine.

India ranks second in Tomato production producing 30.26% of the world’s Tomato production first being China and is followed by Turkey ranking third in the world. (FAOSTAT, 2020). The area under Tomato production in India accounts for to 46.72 thousand ha with a production of 34.29 million tonnes in year 2019–20. Andhra Pradesh ranks first in area and production of tomatoes in the years. Tomato in year 2019–20 followed by Madhya Pradesh and Karnataka. In Uttar Pradesh area under production is 0.20 lakhs hectares while production is estimated to be 5.29 million tonnes for the years 2019–20. (Source: NHB, Ministry of Agriculture & Farmers Welfare, Government of India, 2020–21).

Role of Fertilizer and Micronutrients on crop plants

The high productive ability of tomato puts tremendous pressure on soil for the removal of nutrients. As such liberal application of nutrients is needed to meet the nutritional requirements of the crops, however, in the wake of the energy crisis, harmful effect on soil health and ever-increasing prices of chemical fertilizer becomes a problem before for the producers. Therefore, a dire need have has been felt to

apply fertilizers in more and more amounts to fulfil the requirements of the crop as well as to nourish the health and fertility status of the soil, but should be applied in appropriate doses to reduce the imparity of soil.

Tomato is one of the most important crops and it has a rich economic importance. Micronutrients promote to produce of higher yield and increase harvest quality, maximizing a plant's plant genetic potential and the presence of micronutrients impacts impact on root development, fruit setting, plant vigour and health. Micronutrients are fundamental for balanced nutrition and a tremendous tool to help farmers in increasing crop yield and quality. This experiment was conducted to help in understanding the “**Effect of foliar application of micronutrients on growth, yield and quality of tomato (*Solanum lycopersicum* L.) cv. Arka Samrat.**

Materials and Methods

The area of the Prayagraj district comes under the subtropical belt in the south east of Uttar Pradesh, which experiences extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46° C-48° C and seldom falls as low as 4°C- 5°C. The relative humidity ranges between 20 to 94 %. The average rainfall in this area is around 1013.4 mm annually. However, occasional precipitation is also not uncommon during the winter months.

The experiment was conducted in Randomized Block Design with 8 treatment and three replications replicated thrice. The treatments were T₀ Control (water spray), T₁ FeSO₄ @ 0.2% spray, T₂ Ca(NO₃)₂ @ 0.2% spray, T₃ H₃BO₃@0.1% spray, T₄ ZnSO₄ @

0.2% spray, T₅ FeSO₄ @ 0.2% + Ca(NO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2% spray, T₆ Ca(NO₃)₂ @ 0.2% + ZnSO₄ @ 0.2% spray, T₇ Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1% spray.

Result

Plant height at 30 DAT

The height of plant significantly varied among different treatment combinations. The maximum plant height (41.98 cm) at 30 DAT was observed with treatment T₆ (Feso₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Znso₄ @ 0.2%) followed by T₈ (Calcium nitrate @ 0.2% + H₃BO₃@0.1%) with 37.73 cm. Minimum plant height (28.15 cm) was observed in T₁ (control), while the remaining treatments are moderate in their growth habit.

Plant height at 60 DAT

The height of plant significantly varied among different treatment combinations. The maximum plant height (91.32 cm) at 60 DAT was observed with treatment T₆ (Feso₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Znso₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) with 86.96 cm. Minimum plant height (77.68 cm) was observed in T₁ (control), while the remaining treatments are moderate in their growth habit.

Plant height at 90 DAT

The height of plant significantly varied among different treatment combinations. The maximum plant height (167.85 cm) at 90 DAT was observed with treatment T₆ (Feso₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Znso₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) with 155.41 cm. Minimum plant height (141.11 cm) was observed in T₁ (control), while the remaining treatments are moderate in their growth habit.

Comment [K4]: Please mention the working method of each factor

Comment [K1]: Need to be complete

Comment [K2]: Please mention the statistical analysis

Comment [K3]: Please mention the time of application

Number of leaves per plant at 30 DAT

It is evident that the number Number of leaves per plant was influenced by different treatments at all successive stages stage of growth. There was significant difference between the treatments at 30 days after planting among the treatments applied, T₆ (FeSO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) with 96.33 increase significantly better Number of leaves per plant followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) with 91.67 whereas the minimum score was observed in treatment T₁ (Control) with 76.67.

Number of leaves per plant at 60 DAT

It is evident that the nNumber of leaves per plant was influenced by different treatments at all successive stages stage of growth. There was significant difference between the treatments at 60 days after planting among the treatments applied, T₆ (FeSO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) with 144.67 increase significantly better Number of leaves per plant followed by T₈ (Calcium nitrate @ 0.2% + H₃BO₃@0.1%) with 136.67 whereas the minimum score was observed in treatment T₁ (Control) with 129.33.

Number of leaves per plant at 90 DAT

It is evident that the nNumber of leaves per plant was influenced by different treatments at all successive stages of growth. There was significant difference between the treatments at 90 days after planting among the treatments applied, T₆ (FeSO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) with 204.33 increase significantly better Number of leaves per plant followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) with 200.67 whereas the minimum score was

observed in treatment T₁ (Control) with (192.67).

The minimum Days to 50% flowering T₆ (FeSO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) with 37.33 days, followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃ @0.1%) with 38.33 days whereas maximum Days to 50% flowering 44.67 days was recorded in control.

At harvest time maximum number of flower per cluster 5.12 was recorded in T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by 4.84 T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) whereas minimum fruits per cluster 3.92 were found in T₁ (control).

The maximum number of fruits per plants (60.33) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) i.e., 57.67 and the lowest number of fruit per plant (50.33) were observed in T₁ (Control).

The maximum number of fruits per plants (5.06) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) i.e., 4.61 and the lowest fruit set per cluster (3.90) were observed in T₁ (Control).

The maximum fruits weight (94.22g) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2% + H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) i.e., 87.94g and the lowest fruits weight (64.74g) were observed in T₁ (Control).

The maximum average yield per plant (5.68 kg) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) i.e., 5.07 kg and the lowest average yield per plant (3.26 kg) were observed in T₁ (Control).

The maximum average yield per plot (34.10 kg/plot) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) i.e., 30.42 kg/plot and the lowest average yield per plot (19.55 kg/plot) were observed in T₁ (Control).

The maximum average yield per hectare (189.43 t/ha) were recorded in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Calcium nitrate @ 0.2% + H₃BO₃@0.1%) i.e., 169.01 t/ha and the lowest average yield per hectare (108.63 t/ha) were observed in T₁ (Control)

The maximum TSS (5.23 °Brix) was observed in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Ca(NO₃)₂ @ 0.2% + H₃BO₃@0.1%) with 4.80 °Brix respectively. The minimum TSS (3.83 °Brix) was noticed in treatment T₁ (Control).

The maximum Ascorbic acid (mg/100g) (26.11mg) was observed in treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + Zn SO₄ @ 0.2%) followed by T₈ (Calcium nitrate @ 0.2% + H₃BO₃@0.1%) with 24.30mg respectively. The minimum Ascorbic acid (17.38mg) was noticed in treatment T₁ (Control).

Conclusion

From the above experimental finding it may be concluded that the treatment T₆ (Fe SO₄ @ 0.2% + (CaNO₃)₂ @ 0.2%+ H₃BO₃ @ 0.1% + ZnSO₄ @ 0.2%) was found to be best in the terms of growth Yield and quality of tomato. While, maximum Plant height, highest No. of leaves per plant, Minimum Days to 50% flowering, Maximum No. of flower per cluster, Maximum No. of fruit per plant and maximum Fruit set per cluster, Maximum fruit weight, Maximum average yield per plant, Maximum average yield per plot, Maximum average yield per hectare, Maximum Total soluble solid, Maximum Ascorbic Acid.

Discussion

The foliar application of micronutrients might have improved the ~~soil's soil~~-physical and chemical properties and ~~led to an adequate leading to the adequate~~ supply of nutrients to the plants which might have promoted ~~the~~ maximum vegetative growth while the minimum plant growth was due to non-availability of nutrients. Similar findings were reported by **Sivaiah *et al.* (2013); Meena *et al.* (2015); Kumar *et al.* (2016); Singh *et al.* (2018) and Swetha *et al.* (2018)** in tomato.

Integration of organic fertilizers and biofertilizers favoured vigorous growth and synthesized more cytokinins in plants, which might have helped to the translocation of cytokinins as well as more quantity of available phosphorus through the xylem vessels and their accumulation in the axillary buds that would have favoured the plant to enter into reproductive phase

(Dange *et al.*, 2002). Similar results have also been reported by **Singh and Tiwari (2013)**, **Dixit *et al.* (2018)** and **Singh *et al.*, (2018)**.

UNDER PEER REVIEW




Table 1:- Effect of foliar application of micronutrients Plant height, height, No. of leaves per plant, Days to 50 % flowering, flowering, Number of flowers per cluster, Number of fruit per plant.

Comment [K5]: Need to be add the mean compression

Comment [K6]: Need to be revised and short

| Notation | Treatment | Plant height | | | No. of leaves per plant | | | Days to 50 % flowering | Number of flowers per cluster | Number of fruit per plant |
|----------|--|--------------|--------|--------|-------------------------|--------|--------|------------------------|-------------------------------|---------------------------|
| | | 30 DAT | 60 DAT | 90 DAT | 30 DAT | 60 DAT | 90 DAT | | | |
| T1 | Control | 28.15 | 77.68 | 141.11 | 76.67 | 129.33 | 192.67 | 44.67 | 3.92 | 50.33 |
| T2 | FeSO ₄ @ 0.2% | 29.80 | 78.99 | 145.17 | 80.67 | 130.33 | 194.33 | 43.67 | 4.39 | 51.67 |
| T3 | Ca(NO ₃) ₂ @ 0.2% | 31.54 | 81.75 | 145.51 | 83.00 | 130.67 | 195.00 | 42.33 | 4.33 | 53.33 |
| T4 | H ₃ BO ₃ @ 0.1% | 33.25 | 83.43 | 150.03 | 85.33 | 132.33 | 197.00 | 40.00 | 4.56 | 54.67 |
| T5 | ZnSO ₄ @ 0.2% | 34.54 | 85.66 | 151.72 | 87.00 | 132.33 | 197.33 | 39.33 | 4.39 | 55.67 |
| T6 | Feso ₄ @ 0.2% + (CaNO ₃) ₂ @ 0.2%+ H ₃ BO ₃ @ 0.1% + ZnsO ₄ @ 0.2%) | 41.98 | 91.32 | 167.85 | 96.33 | 144.67 | 204.33 | 37.33 | 5.12 | 60.33 |
| T7 | CaNO ₃ @ 0.2% + ZnSO ₄ @0.2% | 35.70 | 86.89 | 153.97 | 89.67 | 133.33 | 199.00 | 39.00 | 4.84 | 54.00 |
| T8 | CaNO ₃ @ 0.2% + H ₃ BO ₃ @0.1% | 37.73 | 86.96 | 155.41 | 91.67 | 136.67 | 200.67 | 38.33 | 4.55 | 57.67 |
| | 'F' test | S | S | S | S | S | S | S | S | S |
| | C.V. | 1.50 | 0.45 | 0.95 | 1.82 | 2.09 | 0.70 | 3.15 | 5.40 | 2.09 |
| | C.D. at 5% | 0.90 | 0.67 | 2.54 | 2.77 | 4.95 | 2.44 | 2.26 | 0.43 | 2.02 |
| | SE.d(□) | 0.42 | 0.31 | 1.18 | 1.28 | 2.29 | 1.13 | 1.05 | 0.20 | 0.93 |

Table 2:- Effect of foliar application of micronutrients Fruit set per cluster, Average fruit weight, Average Yield per plant, Average Yield per hectare, Total soluble solid, Ascorbic Acid.

| Notation | Treatment | Fruit set per cluster | Average fruit (g)weight | Average Yield kg/per plant | Average Yield kg/per plot | Average Yield per t/hectare | Total soluble solid | Ascorbic Acid |
|----------|--|-----------------------|-------------------------|----------------------------|---------------------------|-----------------------------|---------------------|---------------|
| T1 | Control | 3.90 | 64.74 | 3.26 | 19.55 | 108.63 | 3.83 | 17.38 |
| T2 | FeSO ₄ @ 0.2% | 4.17 | 71.26 | 3.68 | 22.09 | 122.71 | 4.20 | 18.29 |
| T3 | Ca(NO ₃) ₂ @ 0.2% | 4.30 | 75.57 | 4.03 | 24.18 | 134.34 | 4.25 | 19.27 |
| T4 | H ₃ BO ₃ @ 0.1% | 4.44 | 78.17 | 4.27 | 25.64 | 142.43 | 4.28 | 21.28 |
| T5 | ZnSO ₄ @ 0.2% | 4.33 | 83.25 | 4.64 | 27.80 | 154.47 | 4.44 | 22.23 |
| T6 | Feso ₄ @ 0.2% + (CaNO ₃) ₂ @ 0.2%+ H ₃ BO ₃ @ 0.1% + Znso ₄ @ 0.2%) | 5.06 | 94.22 | 5.68 | 34.10 | 189.43 | 5.23 | 26.11 |
| T7 | CaNO ₃ @ 0.2% + ZnSO ₄ @0.2% | 4.44 | 85.80 | 4.63 | 27.80 | 154.44 | 4.78 | 24.21 |
| T8 | CaNO ₃ @ 0.2% + H ₃ BO ₃ @0.1% | 4.61 | 87.94 | 5.07 | 30.42 | 169.01 | 4.80 | 24.30 |
| | 'F' test | S | S | S | S | S | S | S |
| | C.V. | 4.09 | 0.81 | 1.76 | 1.80 | 1.79 | 3.30 | 1.95 |
| | C.D. at 5% | 0.32 | 1.14 | 0.14 | 0.84 | 4.66 | 0.26 | 0.74 |
| | SE.d(□) | 0.15 | 0.53 | 0.06 | 0.39 | 2.15 | 0.12 | 0.34 |

Formatted: Highlight

Nutrients play an important role in improving the productivity and quality of Tomato. The Added-added dose of nitrogen, phosphorus and other essential nutrients increased the vigour vigor of plants, assimilating area, and size of fruit, thereby resulting into a higher weight of fruit. These results are in close conformity with the findings of Ali et al. (2015); Haleema et al. (2017); Satyamurthy et al. (2017); Pandiyan et al. (2018); Singh et al. (2018) and Shnain et al. (2021) as reported in tomato.

Added dose of nitrogen, phosphorus and other essential nutrients increased the vigor of plants, assimilating area, size of fruit, thereby resulting in a into higher weight of fruit. These results are in close conformity with the findings of Kazemi (2013); Saravaiya et al. (2014); Ali et al. (2015); Haleema et al. (2017); Satyamurthy et al. (2017); Pandiyan et al. (2018); Singh et al. (2018) and Shnain et al. (2021)

References

Ali, M. R., Mehraj, H. and Jamal Uddin, A. F. M. 2015. Effects of foliar application of zinc and boron on growth and yield of summer tomato. *Journal of*

Biological and Agricultural Research. **06(01):** 512-517.

Anonymous, (2015). *Horticultural Statistics at a Glance*, Horticulture statistics division department of agriculture, cooperation & farmers welfare ministry of agriculture & farmers welfare government of India.

Basavarajeshwari, C. P., Hosamani, R. M., Ajjappalavara, P. S., Naik, B. H., Smitha, R. P. and Ukkund, K. C. (2008). Effect of Foliar Application of Micronutrients on Growth and Yield components of Tomato (*Lycopersicon esculentum* Mill.). *Karnataka Journal of Agricultural Sciences.* **21(3):** 428-430.

De Candolle A. Origin of cultivated plants. *Hafner Publication Co.*, New York, (Reprint of 2nd edition. 1959).

Directorate of Economics and Statistics, (2020-21) Ministry of Agriculture & Farmers Welfare (DAC & FW), Government of India, 2020-21).

Dixit, A. Dhananjay, S., Tinku, K. S. and Pappu, L. B. (2018). Effect of Foliar Application of Some Macro and Micronutrients on Growth and Yield of Tomato (*Solanum lycopersicum* L.) cv. Arka Rakshak. *International Journal of Current*

Microbiology and Applied Science.**6**: 197-203.

FAOSTAT (2020).
www.fao.org/faostat/en/data/CC. Food Supply- Crops Primary Equivalent. *Visited on 15/12/2021*.

Fisher, R. A. and Yates, F. (1963). Statistical Tables for Biological, Agricultural and Medical Research. *Oliver and Boyd, London*: 143 p.

Francis, F.J. (2000). Encyclopedia of Food Science and Technology. John Wiley and Sons Inc.

Frankel, O. H., Brown, A. H. D. and Burdon, J. J. (1998). The Conservation of Plant Biodiversity. 2nd ed. Cambridge: *Cambridge University Press*. pp. 56-78.

Gopalan B.V., Sastri Rama S.C. and Balasubramanian K.P. (2004). Nutritive value of Indian foods. *National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India*.

Habibullah, S. N., Saravaiya, Y. N., Tandel, S. K., Patel, B. N. and Golakiya, P. D. (2017). Effect of Foliar Application of Micronutrients on Growth and Yield of Tomato under Protected

Culture. *Trends in Biosciences*. **10**(14): 2491-2495.

Haleema, B., Abdur, R. and Hussain, A. S. (2017). Effect of Calcium, Boron and Zinc Foliar Application on Growth and Fruit Production of Tomato. *Sarhad Journal of Agriculture*. **30**(1): 19-30.

Kumar, N. M., Ajay, K. P. and Amin, M. B. (2017). Growth and Yield of Solanaceous Vegetables in Response to Application of Micronutrients – A Review. *International Journal of Innovative Science, Engineering & Technology*. **3**(2): 611-626.

Karpechenko G. D. (1925) Chromosomes of Beetroot. *Bulletin Application of Botany*, **14**: 143-148.

Kumar, N. S., Ahirwar, K., Khare, A. K., Mishra, U. S., Singh, D. and Jatav, R. (2016). Effect of micronutrients and bio-fertilizers on yield and yield attributes of tomato. *International Journal of Agriculture Sciences*. **8**(38): 1780-1782.

Meena, E. C., Maji, S.J., Meena, K., Govind, R., Kumawat, K. R., Meena, S. K. and Sodh, K. (2015). Improvement of Growth, Yield and Quality of Tomato (*Solanum lycopersicum* L.) cv. Azad T-6 with Foliar Application of Zinc and Boron.

International Journal of Bio-resource and Stress Management. **6**(5):598-601.

NHB(2013).

nhb.gov.in/statistics/2013-14. Area and Production of Horticulture Crops- All India. Visited on 08/12/2021.

Pandiyar, R., Sathyamurthy, V. A. and Pugalenti, L. (2018). Effect of foliar application of Micronutrients on Growth, and Quality of Tomato. *International Journal of Trend in Scientific Research and Development.* **2**(4): 1988-1992.

Reddy, G. P. D., Reddy P. S. R., Chandramohan, G. R. and Thanuja, S. G. (2018). Effect of Foliar Application of Micronutrients on Growth and Yield Parameters in Tomato (*Solanum lycopersicon* L.). *International Journal Pure Applied Bio-sciences.* **6**(2): 929-934.

UNDER PEER REVIEW