

# Growth dynamics and yield of zero till maize as influenced by microbial consortia treated paddy residue incorporation

## ABSTRACT

A field experiment was conducted at College farm, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad during 2020-21 and 2021-22 to assess the influence of paddy residue management practices at different fertility levels on certain growth parameters and yield of *rabi* maize succeeding *kharif* rice in a strip plot design. The growth analysis indicated that the leaf area index, crop growth rate, net assimilation rate, and leaf area duration, remained same with residue management practices at 0-30 DAS and 30-60 DAS but between 60 DAS-harvest, maize leaf area index, crop growth rate, net assimilation rate and leaf area duration was significantly increased by incorporation of crop residues treated with microbial consortia and SSP. Different fertility levels affected leaf area index, crop growth rate and leaf area duration at all the growth intervals whereas net assimilation rate remained unaffected. Interaction effect of residue management practices and fertility levels on leaf area index, crop growth rate, net assimilation rate and leaf area duration was found non significant at 0-30 DAS and 30-60 DAS, while at 60 DAS-harvest integration of residues and microbial consortia along with chemical fertilizers resulted in significant increase of LAI, CGR and LAD with no significant improvement in NAR. In 2020-21 and 2021-22, the grain yield and straw yield were higher in incorporation of crop residues treated with microbial consortia and SSP in combination with 125 % RDF which was on par with 100% RDF and 75 % RDF and was significantly superior over other treatments.

**Keywords:** Growth parameters, fertility levels, microbial consortia, *rabi* maize, residue management, yield

## 1. INTRODUCTION

A large amount of rice residue is annually produced in the rice growing countries. Moreover, the adoption of mechanized farming has resulted in leaving a sizeable amount of rice straw in the field after harvesting the grain. According to National Policy for Management of Crop Residues (NPCMR), India generates about 500 Mt of crop residues annually. Burning of crop residues leads to enormous loss of plant nutrients particularly organic carbon. Nearly 80-90% of nitrogen (N), 25% phosphorus (P), 20% potassium (K), 50% sulphur (S) and entire amount of organic carbon was found to be lost in the form of gaseous and particulate matter (Krishnaveni and Dayana, 2021). The crop residues can be used in a efficient and utilizable manner other than burning. Adoption of appropriate tillage practices by retention of residue helps to reduce physical disturbance of the soil and leads to accumulation of more soil organic matter by enhancing microbial population (Pathak *et al.*, 2011).

In the post green revolution era, intensive agriculture with the use of high analysis inorganic fertilizers alone beyond the threshold requirement of crop and soil resulted in diminishing soil fertility and posed a stern hazard to sustainable crop production. Availability of fertilizers at subsidized rates and faster output in terms of yield increment encouraged the farmers towards imbalanced use of fertilizers. This not only led to downward spiraling of natural resources but also made us to depend on non-renewable resources both of which posed hazard to sustainable agriculture. Thus, there is an immediate necessity for alternate sustainable nutrient management strategy to overcome this dawning challenge.

Microbial consortia are a type of biofertilizers that involve symbiotic interaction of two or more microbial groups for enhancing the efficiency. They intensify the turnover of soil organic

Comment [DS1]: Dynamics

Comment [DS2]: Y

Comment [DS3]: Z

Comment [DS4]: T

Comment [DS5]: M

Comment [DS6]: I

Comment [DS7]: M

Comment [DS8]: C

Comment [DS9]: T

Comment [DS10]: P

Comment [DS11]: R

Comment [DS12]: I

Comment [DS13]: (LAI)

Comment [DS14]: (CGR)

Comment [DS15]: (NAR)

Comment [DS16]: (LAD)

Comment [DS17]: Please write the scientific term instead of its abbreviation

Comment [DS18]: Please write the scientific term instead of its abbreviation

Comment [DS19]: F

Comment [DS20]: M

Comment [DS21]: M

Comment [DS22]: R

Comment [DS23]: Y

Comment [DS24]: Must be supported by reference

Comment [DS25]: Must be supported by reference

matter and mobilize various nutrients for plant use (Odoh *et al.*, 2020) and also aid in metabolic actions like fixation of nitrogen from atmosphere, solubilization of soil P, release of fixed K and Zn, thereby increase their availability to crop (Veerendra *et al.*, 2021) in a sustainable way apart from improving soil health (Laxman *et al.*, 2017). Nutrients supply through the integration of microbial consortium containing *Azotobacter*, *Trichoderma harzianum*, and *Pseudomonas fluorescens* with chemical fertilizers can reduce the requirements on the one hand and can also increase the response of chemical fertilizers on the other. Thus the present experiment was planned with the objective to determine the effect of integrated application of residues treated with microbial consortium along with NPK fertilizers on crop growth dynamics and yield of maize.

**Comment [DS26]:** Must be supported by reference

**Comment [DS27]:** Add a little information about the maize and its scientific name

## 2. MATERIALS AND METHODS

### 2.1 Study Region

A two year study (2020–2022) on rice-maize system was conducted at the Agricultural College Farm of Professor Jayashankar Telangana State Agricultural University with 17°32'22"N latitude and 78°41'11"E longitude and an altitude of 550 m above mean sea level. Prior to initiation of the actual experiment, rice crop was grown uniformly with traditional farmers practice for homogenization of soil fertility and for collection of residues for the experiment. The soil of the experimental field was sandy clay loam, deep and free from gravels. Base line soil samples from 0 to 15 cm depth were collected and analyzed before initiation of the experiment. The soil had 0.18 % of soil organic carbon as determined by the wet digestion method (Walkey and Black, 1934), 145 kg ha<sup>-1</sup> of available soil nitrogen examined by the alkaline permanganate method (Subbaiah and Asija, 1956), 38 kg ha<sup>-1</sup> of available P<sub>2</sub>O<sub>5</sub> analyzed by spectrophotometer using olsen's method and 277 kg ha<sup>-1</sup> of available potassium (K) determined by neutral normal ammonium acetate method and analyzed by flame photometer (Piper *et al.*, 1966). The soil pH was 7.84 (1:2.5, soil and water ratio) and bulk density 1.42 g cm<sup>-3</sup> was estimated by the core sampler method (Blake and Hartge, 1986). Rice was grown during rainy season and harvested in winter season during both the years of study. Maize was sown under zero tillage after the harvest of rice. The amount of rainfall received and temperature variation from sowing to harvest was measured for the entire growing season.

**Comment [DS28]:** It was better to measure the content of NPK within the tissue of the plant

### 2.1 Methodology

The experiment was laid out in a strip plot design with three replications. The experiment consisted of combinations of residue management practices and fertility levels as follows:

M<sub>1</sub>: Burning residue before sowing, M<sub>2</sub>: Surface retention of residues, M<sub>3</sub>: Removal of residues before sowing, M<sub>4</sub>: Incorporation at 15 DAS, M<sub>5</sub>: Incorporation + SSP at equivalent to 'P' dose at 15 DAS, M<sub>6</sub>: Spraying consortia of decomposers @ 10% of residue weight + surface retention, M<sub>7</sub>: Spraying consortia of decomposers @ 10% of residue weight + incorporation at 15 DAS, M<sub>8</sub>: Spraying consortia of decomposers @ 10% of residue weight + incorporation at 15 DAS + SSP at equivalent to 'P' dose and three fertility levels (S<sub>1</sub>: 75 % RDF, S<sub>2</sub>: 100% RDF and S<sub>3</sub>: 125 % RDF).

Leaf area index, crop growth rate, net assimilation rate and leaf area duration at intervals for the entire growing season were computed.

### Leaf Area Index (LAI)

Leaf area of five contiguous plants from the border rows leaving the extreme row was measured at 30, 60 DAS and at harvest in maize by using leaf area meter image analysis system. After computing the leaf area as explained above, leaf area index was calculated by using the following formula as suggested by Watson (1952).

$$\text{LAI} = \text{Leaf area (cm}^2\text{)} / \text{Ground area (cm}^2\text{)}$$

### Crop Growth Rate (CGR)

Crop growth rate can be defined as the rate of dry matter production per unit ground area per unit time. It was calculated during 0– 30 DAS, 30– 60 DAS and 60 DAS – harvest in maize as per the formula given by Watson, 1956.

$$\text{Crop growth rate} = \frac{W_2 - W_1}{t_2 - t_1} \times \frac{1}{P} \text{ (g m}^{-2} \text{ day}^{-1}\text{)}$$

where,

$$t_2 - t_1 \quad P$$

$W_1$  and  $W_2$  are the total dry weight (g) at time  $t_1$  and  $t_2$  respectively and

$P$  is the ground area occupied by the plants ( $\text{m}^2$ ).

### Net Assimilation Rate (NAR)

Net assimilation rate indirectly indicates the rate of net photosynthesis. It is expressed as dry matter produced (g) per unit of leaf area ( $\text{m}^2$ ) per unit time (day). It was calculated by the formula outlined by Gregory, 1926. NAR was calculated at 0 – 30 DAS, 30– 60 DAS and 60 DAS – harvest in maize.

$$\text{Net assimilation rate} = \frac{(W_2 - W_1)}{(t_2 - t_1)} \times \frac{(\text{Log}_e A_2 - \text{Log}_e A_1)}{(A_2 - A_1)} \text{ (g m}^{-2} \text{ day}^{-1}\text{)}$$

where,

$W_1$  and  $W_2$  are the total dry weight (g) at time  $t_1$  and  $t_2$  respectively.

$A_1$  and  $A_2$  are leaf area ( $\text{m}^2$ ) at time  $t_1$  and  $t_2$  respectively.

### Leaf area duration (LAD)

Leaf area duration is the integration part of leaf area over a time period (Watson, 1956). It is a measure of its ability to produce leaf area on unit area of land throughout its life. LAD was calculated 0– 30 DAS, 30– 60 DAS and 60 DAS – harvest in maize.

$$\text{Leaf area duration (days)} = \frac{A_1 + A_2}{2} \times (t_2 - t_1)$$

where,

$A_1$  and  $A_2$  are leaf area index at time  $t_1$  and  $t_2$  respectively

For the determination of grain yield, the kernels from the air-dried cobs from each net plot were separated, dried and threshed after proper cleaning to obtain 14 per cent moisture. Weight of grains of each plot was recorded separately and expressed as grain yield in  $\text{kg ha}^{-1}$ . Stover from the net plot area was weighed after thorough drying under the sun and expressed as stover yield in  $\text{kg ha}^{-1}$ .

## 3. RESULTS AND DISCUSSION

### 3.1 Leaf Area Index

The yield is determined by the extent of assimilate synthesis, and leaf area is a fundamental physiological parameter that controls this process. The canopy design, which in turn is determined by agronomic techniques like plant variety, plant density, crop and fertiliser management, etc., determines optimal leaf area. Up to 60 DAS, all treatments generally improved leaf area; after that point, a downward tendency toward maturity due to leaf senescence was seen. An overview at Table 1 showed that throughout both of the experimentation years, residue management and fertility level treatments did not significantly affect leaf area at 30 and 60 DAS. On the other hand, during both trial years, there were notable changes in leaf area between treatments at harvest. The interaction between residue management and fertility levels, remained significant over the course of both years at harvest stage only.

**Table 1: Leaf area index of maize as influenced by residue management options and fertilizer levels during 2020 and 2021.**

Treatments	2020-21			2021-22		
	30 DAS	60 DAS	Harvest	30 DAS	60 DAS	Harvest
<b>Paddy residue management options</b>						
Burning (M <sub>1</sub> )	0.54	3.81	1.48	0.72	4.62	2.10
Retention (M <sub>2</sub> )	0.67	4.05	1.52	0.86	4.83	2.13
Removal (M <sub>3</sub> )	0.51	3.72	1.45	0.70	4.55	2.02
Incorporation (M <sub>4</sub> )	0.56	3.30	1.71	0.73	4.01	2.42
Incorporation + SSP (M <sub>5</sub> )	0.58	3.46	2.07	0.75	4.10	2.67
Retention + consortium (M <sub>6</sub> )	0.69	4.02	1.64	0.89	4.76	2.17
Incorporation + consortia (M <sub>7</sub> )	0.60	3.52	2.45	0.80	4.30	2.92
Incorporation + consortium + SSP (M <sub>8</sub> )	0.63	3.63	2.84	0.83	4.48	3.15
Mean	0.59	3.68	1.89	0.78	0.55	2.44
SE(m) <sub>±</sub>	0.067	0.142	0.114	0.106	0.163	0.067
CD (p=0.05)	NS	NS	0.346	NS	NS	0.202
<b>Fertilizer levels</b>						
75 % RDF (S <sub>1</sub> )	0.48	3.47	1.59	0.72	4.29	2.03
100 % RDF (S <sub>2</sub> )	0.61	3.68	1.91	0.77	4.49	2.47
125 % RDF (S <sub>3</sub> )	0.71	3.92	2.18	0.86	4.59	2.85
Mean	0.59	3.68	1.89	0.78	0.55	2.44
SE(m) <sub>±</sub>	0.02	0.047	0.039	0.013	0.021	0.069
CD (p=0.05)	0.08	0.18	0.15	0.05	0.08	0.27
<b>Interaction</b>						
<b>R×F</b>						
SE(m) <sub>±</sub>	0.06	0.12	0.09	0.07	0.11	0.08
CD (p=0.05)	NS	NS	0.26	NS	NS	0.26
<b>F×R</b>						
SE(m) <sub>±</sub>	0.16	0.29	0.19	0.12	0.17	0.26
CD (p=0.05)	NS	NS	0.55	NS	NS	0.78

During 2020-21, leaf area index was higher with incorporation of residues treated with microbial consortia and SSP followed by incorporation treated with consortia and incorporation treated with SSP. The boost in leaf area index was due to a significant increase in leaf primordials, a high rate of cell division and cell expansion (Gul *et al.*, 2015). On the other hand, significantly lower LAI was noticed with removal of residues which was in equivalence with

incorporation, retention + consortia, surface retention and *in-situ* burning. During 2021-22, similar trend was observed with incorporation +consortia +SSP, incorporation+consortia and incorporation+SSP while bare incorporation was found superior over retention + consortia, surface retention, *in-situ* burning and removal.

Significantly higher leaf area index was registered with fertility level of 125 % RDF over 100 % RDF and 75 % RDF. The greater amounts of cellular protoplasm as well as increased amounts of proteins all of which contributed to increase in leaf area with higher fertilizer levels.

The results find support from the findings of Amanullah *et al.* (2007) who reported increase in leaf area index with increase in nitrogen rate.

**Table 2: Interaction effect of residue management and fertilizer levels on leaf area index of maize at harvest during 2020-21**

Treatment	Leaf area index (harvest)			
	Fertilizer levels			
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
Burning (M <sub>1</sub> )	1.12	1.44	1.89	1.48
Retention (M <sub>2</sub> )	1.09	1.50	1.96	1.51
Removal (M <sub>3</sub> )	1.01	1.44	1.89	1.44
Incorporation (M <sub>4</sub> )	1.33	1.68	2.13	1.71
Incorporation + SSP (M <sub>5</sub> )	1.73	2.22	2.27	2.07
Retention + consortium (M <sub>6</sub> )	1.20	1.72	2.00	1.64
Incorporation + consortia (M <sub>7</sub> )	2.42	2.45	2.47	2.44
Incorporation + consortium + SSP (M <sub>8</sub> )	2.83	2.84	2.86	2.84
Mean	1.6	1.9	2.2	
For comparison the mean of	SEM±		CD	
Residue management	0.11		0.34	
Fertilizer levels	0.03		0.15	
Main plot at same level of sub plot	0.09		0.26	
Sub plot at same level of main plot	0.19		0.55	

**Table 3: Interaction effect of residue management and fertilizer levels on leaf area index of maize at harvest during 2021-22**

Treatment	Leaf area index (harvest)	
	Fertilizer levels	


Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
Burning (M <sub>1</sub> )	1.58	2.05	2.68	2.10
Retention (M <sub>2</sub> )	1.58	2.19	2.62	2.13
Removal (M <sub>3</sub> )	1.51	2.01	2.54	2.02
Incorporation (M <sub>4</sub> )	1.70	2.42	3.15	2.42
Incorporation + SSP (M <sub>5</sub> )	2.33	2.81	2.87	2.67
Retention consortium (M <sub>6</sub> ) +	1.61	2.19	2.72	2.26
Incorporation consortia (M <sub>7</sub> ) +	2.84	2.91	3.00	2.91
Incorporation consortium + SSP (M <sub>8</sub> ) +	3.08	3.18	3.19	3.15
Mean	2.02	2.47	2.84	
For comparison the mean of	SEM <sub>±</sub>		CD	
Residue management	0.06		0.20	
Fertilizer levels	0.06		0.27	
Main plot at same level of sub plot	0.08		0.26	
Sub plot at same level of main plot	0.26		0.78	

No interaction effect was found at 30 and 60 DAS. At harvest, perusal of the data revealed that incorporation of residues treated with consortia and SSP in combination with 75 % RDF resulted in higher leaf area index which was similar with 100 % RDF and 75 % RDF. Similar trend was observed with incorporation of residues treated with consortia alone. Lower leaf area index was observed under removal with 75 % RDF, *in-situ* burning with 75 % RDF, retention with 75 % RDF, retention + consortia with 75 % RDF and incorporation with 75 % RDF. Similar findings of increased leaf area in the chickpea crop with combined application of chemical fertilisers and microbial inoculants (rhizobium + PSB + PGPR) were also reported by Rani *et al.* (2017).


### 3.2 Crop Growth Rate

Crop growth rate increased from sowing to 60 DAS thereafter showed a declining trend which might be due to assimilate translocation to the seed, the senescence and leaf fall at later stage. Residue management practices had significant impact on crop growth rate at 60 DAS – harvest

only (Table 1 & 2) during 2020-21 and 2021-22. Crop growth rate was higher under incorporation + microbial consortia + SSP, while it was lower in removal, *in-situ* burning of residues, incorporation and with the residue retention under zero tillage which were on par with each other during 2020-21. During 2021-22, similar results were attained with incorporation + microbial consortia + SSP but mere incorporation was found effective over removal, *in-situ* burning, retention and retention + microbial consortia indicating incorporation effect dominating over the burning, removal and retention of residues. This is similar to results from Meena *et al.* (2015) where residue addition resulted in higher crop growth rate under zero till conditions.



**Fig 1: Crop growth rate as influenced by residue management and fertility levels during 2020-21.**



**Fig 2: Crop growth rate as influenced by residue management and fertility levels during 2021-22.**

Fertility levels also influenced the crop growth rate at all the intervals of observation which was enhanced significantly by 125 % RDF over 100 % RDF and 75 % RDF. Decrease in the crop growth rate of maize when supplied with 75% RDF might be due to decrease in leaf area index at low rate of fertilizer application. These results corroborate with those obtained by Pandey *et*

al. (2015), who reported a significant increase in CGR and NAR in maize hybrids grown under sufficient phosphorous levels.

**Table 4: Interaction effect of residue management and fertilizer levels on crop growth rate ( $\text{g m}^{-2} \text{day}^{-1}$ ) of maize at 60 DAS-harvest during 2020-21**

Treatment	CGR ( $\text{g m}^{-2} \text{day}^{-1}$ )			
	Fertilizer levels			
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
Burning (M <sub>1</sub> )	17.3	19.5	22.2	19.7
Retention (M <sub>2</sub> )	17.7	19.4	22.7	19.9
Removal (M <sub>3</sub> )	17.7	19.3	21.7	19.6
Incorporation (M <sub>4</sub> )	18.0	20.8	23.3	20.7
Incorporation + SSP (M <sub>5</sub> )	21.6	23.4	23.8	22.9
Retention + consortium (M <sub>6</sub> )	13.6	20.3	26.2	20.0
Incorporation + consortia (M <sub>7</sub> )	24.3	25.2	25.9	25.1
Incorporation + consortium + SSP (M <sub>8</sub> )	26.2	27.9	28.1	27.4
Mean	19.6	22.0	24.2	
For comparison the mean of	SEM <sub>±</sub>		CD	
Residue management	0.71		2.17	
Fertilizer levels	0.50		1.98	
Main plot at same level of sub plot	0.66		1.92	
Sub plot at same level of main plot	1.73		5.01	

Crop growth rate was not influenced by the interaction of both residue management practices and fertility levels at 0-30 DAS and 30-60 DAS but at 60 DAS-harvest, incorporation + microbial consortia + SSP in combination with 125 % RDF recorded higher crop growth rate which was on par with incorporation + microbial consortia+ SSP with 100 % RDF and incorporation + microbial consortia + SSP with 75 % RDF. Similar trend was observed with

combination of incorporation + microbial consortia with 125 % RDF and incorporation + SSP with 125 % RDF. Significantly lower crop growth rate was obtained with the treatment combination of removal of residues with 75 % RDF which was at par with *in-situ* burning of residues with 75 % RDF, surface retention with 75 % RDF and retention + microbial consortia with 75 % RDF.

**Table 5: Interaction effect of residue management and fertilizer levels on crop growth rate ( $\text{g m}^{-2} \text{day}^{-1}$ ) of maize at 60 DAS-harvest during 2021-22**

Treatment	CGR ( $\text{g m}^{-2} \text{day}^{-1}$ )			
	Fertilizer levels			
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
Burning (M <sub>1</sub> )	19.8	25.0	27.5	24.1
Retention (M <sub>2</sub> )	19.9	24.8	28.3	24.3
Removal (M <sub>3</sub> )	20.2	24.1	27.3	23.9
Incorporation (M <sub>4</sub> )	22	27.4	32.0	27.1
Incorporation + SSP (M <sub>5</sub> )	26.4	30.3	30.4	29.0
Retention consortium (M <sub>6</sub> ) +	21.1	25.3	28.7	25.0
Incorporation consortia (M <sub>7</sub> ) +	30.2	31.3	31.6	31.0
Incorporation consortium + SSP (M <sub>8</sub> ) +	32.6	33.0	33.5	33.0
Mean	24.0	27.7	29.9	
For comparison the mean of	SEM $\pm$		CD	
Residue management	0.59		1.80	
Fertilizer levels	0.56		2.23	
Main plot at same level of sub plot	0.64		1.87	
Sub plot at same level of main plot	1.88		5.45	

### 3.3 Net Assimilation Rate ( $\text{g m}^{-2} \text{day}^{-1}$ )

The residue management effect was significant at 60 DAS - harvest only, with highest increase in net assimilation rate under incorporation + microbial consortia + SSP which was at par with incorporation + microbial consortia, incorporation + SSP and incorporation (Table 3). During

first and second year of study, lower values of net assimilation rate were observed under surface retention which was equivalent with *in-situ* burning, removal and retention + microbial consortia.

Fertility levels resulted in a non significant increase of net assimilation at 0 – 30 DAS, 30 – 60 DAS and 60 DAS – harvest intervals during both the years of experimentation. Similarly no interaction effect was found between residue management practices and fertility levels during the *rabi* seasons of 2020-21 and 2021-22.




Fig 3 : Net assimilation rate ( $\text{g m}^{-2} \text{ day}^{-1}$ ) of maize as influenced by residue management and fertilizer levels during 2020-21.





Fig 4 : Net assimilation rate ( $\text{g m}^{-2} \text{ day}^{-1}$ ) of maize as influenced by residue management and fertilizer levels during 2021-22.


### 3.4 Leaf Area Duration(days)

Leaf area duration (LAD) indicates duration for which a certain canopy size is maintained in the field. It is a combined measurement between leaf area index and time. From Table 4 & 5, it

can be inferred that fertility levels exerted significant influence on LAD at all the intervals while, residue management practices exerted a significant influence at 60 DAS –harvest only. Among the residue management treatments, during 2021-22 and 2021-22, at 60 DAS–harvest, incorporation + microbial consortia + SSP resulted in significantly higher leaf area duration over other treatments viz., incorporation + consortia, incorporation + SSP, incorporation, retention + consortia, retention, *in-situ* burning and removal.



**Fig 5: Leaf area duration (days) of maize as influenced by residue management and fertilizer levels during 2020-21**



**Fig 6: Leaf area duration (days) of maize as influenced by residue management and fertilizer levels during 2021-22**

Leaf area duration was higher with 125 % RDF over 100 % RDF and 75 % RDF. Colomb *et al.* (2000) reported a 13.5 % reduction in leaf area duration in field grown maize under phosphorous deficiency. Higher leaf area duration at higher fertility level might be due to efficient interception and utilization of solar radiation per unit area.

The interaction effect on leaf area duration was significantly influenced by both residue management practices and fertility levels during 2021-22. At 60 DAS – harvest, significantly higher leaf area duration was recorded with the combination of incorporation + microbial consortia + SSP with 125 % RDF which was on par with incorporation + microbial consortia + SSP with 100 % RDF and incorporation + microbial consortia+ SSP with 75 % RDF. Significantly lower leaf area duration was recorded with the treatment combination of removal of residues combined with 75 % RDF which was at par with *in-situ* burning of residues with 75 % RDF, surface retention with 75 % RDF and retention + microbial consortia with 75 % RDF.

**Table 6: Interaction effect of residue management and fertilizer levels on leaf area duration (days) of maize at 60 DAS-harvest during 2021-22**

Treatment	LAD (days)			
	Fertilizer levels			
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
Burning (M <sub>1</sub> )	136.6	151.0	166.4	151.3
Retention (M <sub>2</sub> )	140.0	160.8	169.1	156.6
Removal (M <sub>3</sub> )	129.2	147.2	167.0	147.8
Incorporation (M <sub>4</sub> )	125.0	144.8	164.1	144.6
Incorporation + SSP (M <sub>5</sub> )	142.3	156.2	158.5	152.3
Retention consortium (M <sub>6</sub> ) +	138.5	157.7	172.2	156.1
Incorporation consortia (M <sub>7</sub> ) +	158.0	163.0	165.8	162.3
Incorporation consortium + SSP (M <sub>8</sub> ) +	167.5	171.8	175.7	171.7
Mean	142.1	156.6	167.4	
For comparison the mean of	SEM <sub>±</sub>		CD	
Residue management	3.47		10.52	
Fertilizer levels	1.83		7.18	
Main plot at same level of sub plot	2.85		8.26	
Sub plot at same level of main plot	6.54		18.95	

### 3.5 Grain yield (kg ha<sup>-1</sup>)

Data furnished in Table 6 &7 indicated that grain yield was significantly influenced by residue management, fertility levels and their interaction.

Among all the residue management practices tested, incorporation + microbial consortia + SSP recorded significantly higher grain yield in comparison to rest of the practices. During 2020-21 and 2021-22, on average, there was an increase of 23 and 26 %, respectively in incorporation + consortia + SSP treatment over *in-situ* burning treatment. In this study, it is observed that surface retention, removal, *in-situ* burning and retention + consortia recorded lower maize yield in comparison to incorporation. This was obviously associated with an increase in root biomass carbon in residue incorporated plots, which significantly improved the yield of crops (Mandal *et al.*, 2008). The effect of incorporation on enhancing grain yields was reported by several researchers (Ammaji, 2014; Bahrani *et al.*, 2007; Egbe and Ali, 2010).

With respect to different fertility levels tested, higher maize grain yield was obtained with 125 % RDF followed by 100 % RDF and 75 % RDF. The results are supported by the findings of Aulakh, 2001 that higher fertility results in higher yields.

The interaction effect when tested with residue management practices and fertility levels was found significant with the treatment combinations tested. Higher grain yield was evident from incorporation + microbial consortia + SSP with 125 % RDF but it was found equivalent with that of incorporation + microbial consortia + SSP with 100 % RDF and incorporation + consortia + SSP with 75 % RDF treatment combinations. Significantly lower grain yield was recorded with the treatment combination of removal with 75 % RDF which was on par with that of *in-situ* burning with 75 % RDF, surface retention with 75 % RDF and retention + consortia with 75 % RDF combinations. Similar results were obtained by Singh *et al.* (2014) that application of 75 % RDF + *Rhizobium* inoculation increased the root volume through better root development, nodulation, more nutrient availability resulting in vigorous plant growth and dry matter production which in turn resulted in better flowering, pod formation and ultimately seed yield.

**Table 7: Yield (Kg ha<sup>-1</sup>) of maize as influenced by residue management and fertility levels during 2020-21.**

--	--	--

Treatment	Grain yield (Kg ha <sup>-1</sup> )				Straw yield (Kg ha <sup>-1</sup> )			
	Fertilizer levels							
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
<b>Burning (M<sub>1</sub>)</b>	4412	5161	5991	5188	5030	6232	7052	6105
<b>Retention (M<sub>2</sub>)</b>	4488	5438	5869	5265	4295	6245	7941	6160
<b>Removal (M<sub>3</sub>)</b>	4039	5255	6220	5171	5284	6180	6666	6043
<b>Incorporation (M<sub>4</sub>)</b>	4564	5418	6239	5407	4649	6553	8061	6421
<b>Incorporation + SSP (M<sub>5</sub>)</b>	5381	5923	6088	5797	5611	7492	7956	7020
<b>Retention + consortium (M<sub>6</sub>)</b>	4724	5355	5941	5340	5537	6254	7268	6353
<b>Incorporation + consortia (M<sub>7</sub>)</b>	6132	6211	6273	6205	7508	7581	7702	7597
<b>Incorporation + consortium + SSP (M<sub>8</sub>)</b>	6487	6529	6787	6601	7932	8164	8357	8151
<b>Mean</b>	5028	5661	6176		5731	6838	7625	
<b>For comparison the mean of</b>	SEM±		CD		SEM±		CD	
<b>Residue management</b>	125.2		379.8		159.1		482.7	
<b>Fertilizer levels</b>	127.7		501.6		184.0		722.5	
<b>Main plot at same level of sub plot</b>	139.9		405.3		217.9		631.2	
<b>Sub plot at same level of main plot</b>	413.9		1199.1		666.9		1932.1	

### 3.6 Straw yield (kg ha<sup>-1</sup>)

The data on maize straw yield (kg ha<sup>-1</sup>) as influenced by residue management and fertility levels is furnished in Table 6 & 7 .

Significantly higher maize straw yield ( $\text{kg ha}^{-1}$ ) after harvest was attained by the adoption of different residue management methods. Incorporation + microbial consortia+ SSP registered significantly higher straw yield followed by incorporation + microbial consortia, incorporation + SSP and incorporation. On the other hand, significantly lower straw yield was observed under removal which was on par with *in-situ* burning, surface retention, retention + microbial consortia among the different residue management methods.

Performance of maize was better under residue addition due to the improvement of soil fertility by N mineralization from the residues resulting in higher grain and stover yields. Also, it is apparent that benefits of zero tillage are accrued only when residues are retained than burnt or removed from the soil. Better crop growth and yield under zero tillage with residue addition as compared to conventional tillage with residue removal have been reported earlier by many researchers (Jat *et al.*, 2019; Thapa *et al.*, 2019; Meena *et al.*, 2015).

Irrespective of residue management practices, higher straw yield was obtained with 125 % RDF as compared to 100 % RDF. Significantly lower straw yield was reported with 75 % RDF among fertility levels. This corroborates with earlier reports by Patel *et al.* (2021), who reported that higher straw yield was obtained with 100 % RDF compared to 50 % RDF and 75 % RDF due to optimum supply of nitrogen and phosphorous which plays a crucial role in physiological processes in a plant resulting in increased growth and yield.

Significant interaction effect was noticed with residue management practices and fertility levels. Among all the tested combinations, incorporation + microbial consortia + SSP with 125 % RDF recorded significantly higher straw yield but it was statistically at par with that of incorporation + microbial consortia + SSP with 100% RDF and incorporation + microbial consortia + SSP with 75 % RDF combinations. On the other hand, significantly lower straw yield was recorded with the treatment combinations of removal with 75 % RDF which was statistically comparable with that of *in-situ* burning with 75 % RDF, retention + microbial consortia with 75 % RDF and surface retention with 75 % RDF during *rabi*, 2020-21 and 2021-22.

**Table 8: Yield (Kg ha<sup>-1</sup>) of maize as influenced by residue management and fertilizer levels during 2021-22.**

Treatment	Grain yield (Kg ha <sup>-1</sup> )				Straw yield (Kg ha <sup>-1</sup> )			
	Fertilizer levels							
Residue management	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean	75 % RDF (S <sub>1</sub> )	100 % RDF (S <sub>2</sub> )	125 % RDF (S <sub>3</sub> )	Mean
<b>Burning (M<sub>1</sub>)</b>	4828	5210	5807	5282	5170	6396	7219	6262
<b>Retention (M<sub>2</sub>)</b>	4909	5359	5723	5330	4816	6417	7797	6343
<b>Removal (M<sub>3</sub>)</b>	4520	5403	5801	5241	4352	6221	8041	6205
<b>Incorporation (M<sub>4</sub>)</b>	5383	5771	6292	5815	6206	6873	7887	6989
<b>Incorporation + SSP (M<sub>5</sub>)</b>	5867	6291	6489	6216	6970	7750	7814	7511
<b>Retention + consortium (M<sub>6</sub>)</b>	4976	5377	5854	5402	5060	6490	7839	6463
<b>Incorporation + consortia (M<sub>7</sub>)</b>	6518	6631	6693	6614	7915	8055	8143	8038
<b>Incorporation + consortium + SSP (M<sub>8</sub>)</b>	6929	6971	7096	6999	8269	8568	8828	8555
<b>Mean</b>	5491	5877	6219		6095	7096	7946	
<b>For comparison the mean of</b>	SEM <sub>±</sub>		CD		SEM <sub>±</sub>		CD	
<b>Residue management</b>	125.1		379.7		169.2		513.3	
<b>Fertilizer levels</b>	82.8		325.2		167.4		657.3	
<b>Main plot at same level of sub plot</b>	100.1		290.0		218.2		632.3	
<b>Sub plot at same level of main plot</b>	238.7		691.7		647.4		1875.4	

#### **4.CONCLUSION**

From the present study, it can be concluded that incorporation of residues treated with microbial consortia and SSP with 75 % RDF led to enhanced growth parameters and improved yield over *in-situ* burning and removal with 125 % RDF.

UNDER PEER REVIEW

## REFERENCES

1. Amanullah K, Rahman H, Shah Z and Shah P. Plant density and nitrogen effects on growth dynamics, light interception and yield of maize. *Archives of Agronomy and Soil Science*. 2008; 54(4): 401-411.
2. Ammaji P. Productivity of maize as affected by crop residue incorporation and nitrogen levels in legume-cereal sequence. *Ph.D Thesis*, submitted to Acharya N.G Ranga Agricultural University, Andhra Pradesh. 2014.
3. Aulakh MS, Khera TS, Doran JW and Bronson JF. Managing crop residues with green manure, urea and tillage in a rice-wheat rotation. *Soil Science Society of American Journal*. 2001;65: 820-827.
4. Bahrani MJ, Raufat MH and Ghadiri H. Influence of wheat residue management on irrigated corn grain production in a reduced tillage system. *Soil and Tillage Research*. 2007; 94: 305-309.
5. Blake GR and Hartge KH. Bulk density. In: Klute, A. (Ed.), *Methods of Soil Analysis, Part I*, ASA Monograph No. 9. Lewis Publishers, Madison, WI. 1986;363-376.
6. Colomb B, Kiniry JR, and Debaeke P. Effect of soil phosphorous on leaf development and senescence dynamics of field-grown maize. *Agronomy Journal*. 2001;92: 428-435.
7. Egbe OM and Ali A 2010. Influence of soil incorporation of common food legume stover on the yield of maize in sandy soils of Moist Savanna Woodland of Nigeria. *Agriculture and Biology Journal of North America*. 2001;1(2): 156-162.
8. Gul S, Khan MH, Khanday BA and Sabeena N. Effect of sowing methods and npk levels on growth and yield of rainfed maize (*Zea mays* L.). *Scientifica*. 2015; 10(2): 1-6.
9. Gregory FG. The effect of climatic condition on the growth of barley. *Annals of Botany*.1926;40: 1-26.
10. Jat ML, Gathala MK, Saharawat YS, Ladha JK and Singh Y. Conservation agriculture in intensive rice-wheat rotation of western Indo-gangetic plains: Effect on crop physiology, yield, water productivity and economic profitability. *International Journal of Environmental Sciences and Natural Resources*.2019;18(3): 88-95.
11. Krishnaveni AS and Dayana. A review on crop residue management. *International Journal of Current Microbiology and Applied Sciences*, 2021;10(3): 28-33.
12. Laxman T, Ramprakash T, Kumar A and Srinivas A. Microbial activity in *Bt* cotton soils as influenced by the biofertilizer consortia and foliar nutrition under rainfed. *International Journal of Current Microbiology and Applied Sciences*.2017;6(8): 3551-3554.
13. Mandal B, Majumdar B, Adhya TK, Bandyopadhyay PK, Bandyopadhyay A, Sarkar D and Mishra AK. Potential of double cropped rice ecology to conserve organic carbon under sub-tropical climate. *Global Change Biology*.2008;14: 1-13.

14. Meena JR, Behera UK, Chakraborty D and Sharma AR. Tillage and residue management effects on soil properties, crop performance and energy relations in greengram under maize based cropping systems. *International Soil and Water Conservation*. 2015;3: 261-272.
15. Odoh CK, Sam K, Zabbey N, Eze CN, Nwankwegu AS, Laku C and Dumpe BB. Microbial Consortium as biofertilizers for crops growing under the extreme habitats. In book: *Plant Microbiomes for Sustainable Agriculture*.2020;381-424.
16. Pandey R, Khetarpal S, Jain V and Kushwaha SR. Phosphorous fertilization improves growth analysis traits and reduces anthesis-to-silking interval leading to increased grain yield in maize. *Indian Journal of Plant Physiology*. 2015;20(4): 385-390.
17. Patel, R., Shroff, P.M., Parmar, P.M. and Shah, S.N. (2021). Effect of varying fertility levels and biofertilizer on growth, yield and economics of fenugreek. *Biological Forum-An international Journal*, 13(3a): 210-213.
18. Pathak H, Saharat YS, Gathala M, Ladha JK. Impact of resource conserving technologies in the rice-wheat system. *Greenhouse Gas Science and Technologies*. 2011;261-277.
19. Piper CS. *Soil and Plant analysis*. Inter Science Publisher. New York. USA. 1966; 59.
20. Rani S, Kumar P and Yadav AK. Effect of biofertilizers in conjunction with chemical fertilizers on growth behaviour and profitability of field pea (*Pisum sativum* L.) grown in western plains of Haryana. *Chemical Science Review and Letters*. 2017. 6(22): 801-805.
21. Subbiah BV and Asija GL. A rapid method for the estimation of nitrogen in soil. *Current Science*. 1956;26: 259-260.
22. Thapa B, Khanal BR and Marahatta S. Effect of conservation agriculture on yield and yield attributing properties of wheat. *Advances in Plants and Agricultural Research*.2019; 9(2): 329-335.
23. Veerendra MB, Padmaja B, Malla Reddy M and Triveni S. Yield, Nutrient uptake, quality and economicsof foxtail millet cultivation as influenced by integrated nutrient management with bacterial consortia and liquid manures. *International Journal of Current Microbiology and Applied Sciences*.2021;10(3): 1703-1711.
24. Walkey A and Black CA. An estimation of digestion method for determining soil organic matter and proposed modification of chromic acid titration method. *Soil Science*. 1934;37(1): 29-34.
25. Watson DJ. The physiological basis of variation in yield. *Advances in Agronomy*.1952;6:103-109.

UNDER PEER REVIEW