

Original Research Article

Impact assessment of different level of mycorrhiza on the growth parameters and nutrient content of *Capsicum annum*

Abstract:- Soil microbes play an important role in the biogeochemical cycling of nutrients and their availability to plants, which is important for sustaining soil health and agricultural sustainability. In light of this, vesicular arbuscular mycorrhiza (VAM) plays an important function. Hyphae of VAM can reach far beyond the plant root zone, allowing it to obtain nutrients from a considerably larger region of soil. Mycorrhiza improves plant nutrition by facilitating the uptake of minerals such as phosphorus and immobile trace elements such as zinc, cobalt, magnesium, iron, copper, molybdenum, and others. Mycorrhiza improves plant growth, productivity and yield by increasing the rate of photosynthesis. In the present study *Glomus fasciculatum* was tested on *Capsicum annum* plant with different inoculum levels (100, 150, 200 and 400 chlamydo spores/kg soil) and found that Plant height, Root length, Dry weight of root and shoot, NPK content, per cent mycorrhizal colonization and sporocarp number were maximum when 400 spores were used for inoculation and minimum were found in untreated plants. SPAD chlorophyll content was highest at 90 DAT among all the observation periods.

Keyword: Soil microbes, arbuscular mycorrhiza, chlorophyll, plant nutrition.

Introduction:- Arbuscular mycorrhizal (AM) fungi can be found in all soils and colonise the roots of a wide range of plant species. These fungi can boost plant growth and reproduction by improving nutrient uptake, particularly minerals that are immobile in soil like phosphorus. Plants can also benefit from AM fungus because they stimulate growth-regulating chemicals, increase photosynthesis, improve osmotic adjustment under drought and salinity conditions, and increase pest resistance (Al-Karaki, 2006). Plant growth and nutrient uptake are the key effects of AM on their host plant (Ortas *et al.*, 2001). In addition, mycorrhizal inoculation minimises the amount of fertiliser required for inoculated plants (Charron *et al.*, 2001). The arbuscular mycorrhizal fungi (AMF) found associated with the majority of land plants including those of the arid areas (Stutz *et al.*, 2000), once it is established in soil, ~~than then~~ it increases mineral nutrition uptake, mainly phosphorus and enhance plant growth. VAM not only increase the uptake of phosphorous, but also helps in uptake of zinc, copper, sulphur, potassium and calcium (Cooper and Tinker, 1978). Additionally In addition, it protect plants against environmental stress such as soil salinity (Giri *et al.*, 2003), drought (Al-Karaki *et al.*, 2004) and pathogens such as Fusarium wilt (Habte *et al.*, 1999). Therefore studies were conducted on association of AM fungi with *Capsicum annum* to observe growth and nutrient content in pot experiment.

Material and methods:- The present study entitled “Impact assessment of different level of mycorrhiza on the growth parameters and nutrient content of *Capsicum annum*” was executed in screen house and laboratories of Plant Pathology department CCS HAU, Hisar during 2018. The AM fungi *i.e.* *Glomus fasciculatum* was taken from the department of Plant Pathology, Chaudhary Charan Singh Haryana Agricultural University, Hisar. One hundred gram of mycorrhizal inoculum containing about 500 extramatrical chlamydo spores and infected root bits were put in upper 5 cm soil layer per pot. Thirty days old seedling of *Capsicum annum* cv. Pusa Jwala was sown in the pots (Four numbers of sets were maintained) and statistical design was CRD. These plants were watered regularly. Observations, number of mycorrhiza spores/100 gram of soil (Gerdemann and Nicolson, 1963) and Percentage mycorrhizal colonization in roots (Phillips and Hayman, 1970) were taken. Plant height, root length, dry weight of root and shoot at 30, 45, 60 and 90 day after transplanting (DAT) and Nutrient content (NPK) of root and shoot at 90 DAT were calculated.

Statistical analysis: Statistical analysis of experiment was carried out using opstat software at <http://hau.ac.in>.

Maintenance of *Glomus fasciculatum*:-

In 20 cm diameter earthen pots the mycorrhizal fungi (*Glomus fasciculatum*) were maintained on wheat (*Triticum aestivum*) and pearl millet (*Pennisetum typhoides*). These pots were filled with 5 kg sterilized river sand. In upper 5 cm soil layer put one hundred g of mycorrhizal inoculum which contain about 450-500 chlamydo spore and root bit and then ten seeds of wheat or pearl millet were sown and watered regularly. Hoagland’s nutrient solution was applied @ 10 ml/pot after every 30 days of transplanting. After 90 days shoot portion of plant were cut at soil level and left the soil in pots to air dry. The soil was crumbled and cut the rootlets into 1 cm segments. This soil was used as a mycorrhizal inoculum.

Chart 1: Hoagland Solution (Plant Nutrition Solution)

Components	Stock Solution	mL stock solution/1L
2M KNO ₃	202 g/L	2.5ml/L
2M Ca(NO ₃) ₂ •4H ₂ O	236 g/0.5 L	2.5 ml /L
Iron (Sprint 138 iron chelate)	15 g/L	1.5 ml /L
2M MgSO ₄ •7H ₂ O	493 g/L	1 ml /L
NH ₄ NO ₃	80g/L	1 ml /L
Minors:		

H ₃ BO ₃	2.86 g/L	1 ml /L
MnCl ₂ •4H ₂ O	1.81 g/L	1 ml /L
ZnSO ₄ •7H ₂ O	0.22 g/L	1 ml /L
CuSO ₄ •5H ₂ O	0.051g/L	1 ml /L
H ₂ MoO ₄ •H ₂ O or	0.09 g/L	1 ml /L
Na ₂ MoO ₄ •2H ₂ O	0.12 g/L	1 ml /L
1M KH ₂ PO ₄ (pH)	136 g/L	1 ml /L

Mycorrhizal colonization

Mycorrhizal colonization was calculated by Staining of roots by following procedure given by Phillips and Hayman (1970).

Staining of root – Paragraph needs paraphrasing to make sense

Roots were cut into 1 cm segments, heat the roots in 10 per cent KOH at 90°C for one hour (Tense? Paraphrase), washed these roots with fresh (10 per cent) KOH solution, immersed roots in alkaline hydrogen peroxide (H₂O₂) for 30 minutes. Then rinsed? with distilled water to remove the excess of H₂O₂ and acidified with 5 N HCL for 30 minutes. Roots were simmering?? in trypane blue in lactophenol (0.05%) for 5 min. Finally, roots were put in lactophenol to remove the extra dye and examine the roots under microscope.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

$$\text{Mycorrhizal colonization (\% in roots)} = \frac{\text{Sum of all numerical ratings} \times 100}{\text{Total number of sample assessed} \times \text{Maximum scale}}$$

Estimation of sporocarp in soil- should be in past tense since this is what you did

Estimation of sporocarp in soil was done by Wet Sieving and Decantation Technique given by Gerdemann and Nicolson (1963). Firstly, The soil sample was mixed well and then 100 g soil was suspended in a pan A add one liter of water and mix it well. Wait for 30 seconds. Suspension was passed through 20 mesh sieve and filtrate was collected into a pan B. Material of pan A was discard. Suspension of B pan was stirred with hand and allows it for few second to settled down then passed through 60 mesh sieve. Filtrate was collected in pan C. Suspension of pan C was passed through the 100 mesh sieve. Maximum mature sporocarps were collected on 100 mesh sieve. One hundred mesh sieve residue was collected into a beaker after washing in order to remove the excess soil and other particles. (you cannot start a sentence with a number). One ml of this solution was taken in counting dish and examined under stereomicroscope microscope and count the sporocarp population in soil.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Estimation of Nitrogen content, phosphorous content and potassium content

For the estimation of nitrogen content, The Lindner method (1944) was adopted. For the estimation of phosphorous content, Vanadomolybdophosphoric yellow color method given by Koenig and Johnson, (1942) was adopted. Potassium was determined in the acid digest of plant samples by using flame photometer (Elico CL 361, India) by direct reading.

Chlorophyll content

Chlorophyll content of the plant was calculated by using SPAD chlorophyll meter.



Fig: 1 Estimation of chlorophyll content of *Capsicum annum* using SPAD meter

Results:-

The present study was done with the objective of determining the best dose of mycorrhization in *Capsicum annum* plants. A pot experiment was conducted in screen house of Plant Pathology, CCS HAU, Hisar, to see evaluate the effect of *Glomus fasciculatum* with different doses may not be relevant here. Mycorrhizal colonization in roots and number of sporocarps in soil was calculated at 30, 45, 60 and 90 days after transplanting. Plant height, root length, dry weight of root and shoot and SPAD chlorophyll content at 30, 45, 60 and 90 days after transplanting were recorded. Nutrient content (NPK) at 90 days after transplanting was calculated this is part of methodology.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight



Fig:2 Effect of different doses of mycorrhiza on plant height

Table: 1 Effect of mycorrhizal application on plant height and root length of *Capsicum annum* plant at 30, 45, 60 and 90 days after transplanting

Plant height (cm)						Root length (cm)				
Inoculum levels (Sporocarp/Kg soil)	Days After Transplanting (DAT)					Days After Transplanting (DAT)				
	30	45	60	90	Mean B	30	45	60	90	Mean B
100	10.10	11.67	17.13	20.03	14.73	7.43	8.80	9.30	10.04	8.89
150	12.13	12.50	18.53	20.87	16.01	7.80	9.24	11.23	11.87	10.03
200	12.87	15.80	19.67	25.27	18.40	8.53	11.83	12.13	12.80	11.33
400	15.60	19.33	21.40	29.23	21.39	8.77	13.40	13.60	14.60	12.59
Control	9.57	11.43	14.93	19.77	13.93	7.23	8.40	9.20	9.93	8.69
Mean A	12.05	14.15	18.33	23.03		7.95	10.33	11.09	11.85	
CD at 5 % level	DAT = 0.46 Inoculum level = 0.52 DAT× Inoculum level= 1.03					Inoculum level = 0.31 DAT = 0.28 DAT× Inoculum level = 0.62				

Effect of soil application with different doses of AM fungi on plant height and root length of *Capsicum annum* (Table 1) was observed and found that application of the mycorrhizal species *Glomus fasciculatum* with different doses significantly increases in plant height of *Capsicum annum* at 30, 45, 60 and 90 days after transplanting. The maximum plant height (29.23 cm) was observed, when 400 chlamydo spores/kg soil were used for inoculation followed by 200 chlamydo spores/kg soil (25.27 cm) among the all the inoculum levels and minimum plant height was recorded in control (19.77 cm). Irrespective of inoculum level maximum plant height was observed at 90 DAT. Highest root length (14.60 cm) was observed at 90 DAT, when 400 chlamydo spores/kg soil were used followed by 200 chlamydo spores/kg soil (12.80 cm) among the all the inoculum levels and the lowest root length was recorded in control (9.93 cm). Among all the inoculum levels maximum plant height and root length was observed, when 400 chlamydo spores/kg soil were used for inoculation as compare to control.

Table: 2 Effect of different doses of mycorrhiza on dry shoot weight and dry root weight of *Capsicum annum* plant at different inoculum levels

Dry shoot weight (g)						Dry root weight (g)				
Inoculum levels (Sporocarp/Kg soil)	Days After Transplanting (DAT)					Days After Transplanting (DAT)				
	30	45	60	90	Mean B	30	45	60	90	Mean B
100	0.44	0.92	0.93	1.84	1.03	0.83	0.93	1.05	1.17	1.00
150	0.62	1.00	1.06	1.91	1.15	0.94	1.09	1.24	1.39	1.16
200	0.84	1.12	1.33	2.08	1.34	1.16	1.23	1.38	1.56	1.33
400	0.97	1.27	1.48	2.17	1.47	1.23	1.47	1.63	1.79	1.53
Control	0.47	0.96	1.22	1.70	1.09	0.48	0.80	1.03	1.11	0.86
Mean A	0.67	1.05	1.20	1.94		0.93	1.10	1.27	1.40	
CD at 5 % level	DAT = 1.18 Inoculum level = 1.18 DAT× Inoculum level= 2.36					DAT = 1.65 Inoculum level = 1.65 DAT× Inoculum level= 3.30				

The effect of mycorrhizal inoculation on dry shoot weight and dry root weight of *Capsicum annum* was observed (Table 2) and found that maximum dry shoot weight was observed, when 400 spores were inoculated (2.17 g), followed by 200 spores (2.08 g), 150 spores (1.91 g), 100 spores (1.84 g) and minimum was found in control (1.09 g). Irrespective of the inoculum level the maximum dry root length was observed, when *Glomus fasciculatum* was inoculated (90 DAT). Among all the four observations period (30 DAT, 45 DAT, 60 DAT and 90 DAT) the maximum dry shoot weight was observed at 90 DAT. All the treatments were significantly different with each other. The maximum dry root weight was recorded, when 400 spores of *Glomus fasciculatum* (1.79 g) were inoculated and minimum was recorded in control (1.11 g) at 90 DAT.

Table: 3 Impact of soil application with different doses of AM fungi on Mycorrhizal colonization (%) and Sporocarp/ 100g soils

Mycorrhizal colonization (%)			Sporocarp number (per 100g soil)
Inoculum levels (Sporocarp/Kg soil)	Days After Transplanting (DAT)		Days After Transplanting (DAT)

	30	45	60	90	Mean B	30	45	60	90	Mean B
100	23.33	35.00	57.33	60.00	43.92	23.0	35.0	44.7	51.0	38.4
150	28.00	40.00	59.00	63.67	47.67	24.3	42.7	55.3	64.7	46.8
200	31.33	46.67	67.33	72.00	54.33	34.7	55.3	66.7	75.0	57.9
400	35.00	51.33	72.33	77.33	59.00	40.3	69.0	81.0	89.3	69.9
Control	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Mean A	29.42	43.25	64.00	68.25		30.6	50.5	61.9	70.0	
CD at 5 % level	DAT = 1.18 Inoculum level = 1.18 DAT× Inoculum level= 2.36					DAT = 1.65 Inoculum level = 1.65 DAT× Inoculum level= 3.30				

Mycorrhizal root colonization (%) and sporocarp number were calculated at 30, 45, 60 and 90 days after transplanting (Table 3). The maximum root colonization (77.33 %) and sporocarp number (89.3) were found when 400 chlamydo spores/kg soil were used and minimum root colonization (60 %) and sporocarp number (51) were calculated when 100 chlamydo spores/kg soil were used.

Table 4: Effect of mycorrhization on chlorophyll content and NPK content of *Capsicum annuum*

Treatments	Chlorophyll content					NPK content			
	Days after transplanting (DAT)					Mean B	N	P	K
	30	45	60	90					
100	17.33	19.73	20.70	21.43	19.80	1.14	0.38	1.58	
150	20.00	21.50	22.87	25.07	22.36	1.16	0.4	1.67	
200	24.03	24.87	25.57	28.87	25.83	1.18	0.48	1.77	
400	26.57	27.23	29.50	32.63	28.98	1.19	0.56	1.84	
C	16.77	17.80	19.50	20.67	18.68	1.12	0.38	1.43	
Mean A	20.94	22.23	23.63	25.73					
CD at 5 % level	DAT= 0.83 Inoculum level = 0.93 DAT×inoculum level = N/A						N/A	0.03	0.09




Plate: 1 Effect of different inoculum levels of mycorrhiza on chlorophyll content of *Capsicum annum*

Application of *Glomus fasciculatum* with different doses (100, 150, 200 and 400 sporocarp/kg soil) significantly increase the chlorophyll content of *Capsicum annum* (Table 4). Data was statistically analysed and found that the chlorophyll content varied significantly at 30, 45, 60 and 90 days after transplanting. The maximum chlorophyll was recorded, when 400 chlamydo spores/kg soil were inoculated (32.63) and minimum was recorded in control (20.67) at 90 DAT among all the inoculum levels. The maximum NPK content was recorded, when 400 spores of *Glomus fasciculatum* were inoculated (1.19 per cent N, 0.56 per cent P and 1.84 per cent K at 90 DAT at 90 DAT and minimum NPK content was recorded in control (1.12 per cent N, 0.38 per cent P and 1.43 per cent K). Irrespective of the days and species the maximum NPK content was observed when 400 sporocarp/kg soil was inoculated. Among all the four observation period *i.e.* 30 DAT, 45 DAT, 60 DAT and 90 DAT, the maximum NPK content was observed at 90 DAT.

Discussion:

In the present investigation effect of *Glomus fasciculatum* with different doses (100, 150, 200 and 400 sporocarps/kg soil) were observed on growth parameters, mycorrhizal per cent colonization and sporocarp number, NPK content and chlorophyll content of *Capsicum annum* plant and found that with increase in inoculum level of mycorrhiza corresponding increase in plant height, root length, dry weight of shoot and root, chlorophyll content and NPK content of *Capsicum annum* plants was obtained. Maximum plant height and growth parameters were found in inoculated plants and minimum was observed in uninoculated (control). Similarly, Thilagar and Bagyaraj (2013) studied on different species of arbuscular mycorrhizal fungi (*Acaulospora laevis*, *Gigaspora margarita*, *Glomus bagyarajii*, *G. etunicatum*, *G.*

fasciculatum, *G. intraradices*, *G. leptotichum*, *G. macrocarpum*, *G. monosporum*, *G. mosseae* and *Scutellospora calospora*) and found that all different species of mycorrhiza significantly increase the plant growth parameters. Ibrahim *et al.* (2011) conducted an experiment to evaluate the effect of five arbuscular mycorrhizal (AM) fungi *viz.*, *Glomus mosseae*, *G. clarum*, *G. caledonium*, *G. intraradices* and *G. etunicatum* and their mixture were taken on maize and found that inoculated plants with different mycorrhizal fungi, increased in shoot, root dry weight, P, Zn and seedling quality content, seedlings flowered earlier as compared to control plants. Similar results were observed by Ortas *et al.* 2011; Bona *et al.* 2018 and Anli *et al.* 2020. The mutualistic association between roots of higher plants and soil borne fungi was reported by Menendez *et al.* (1999). Mycorrhiza fungi benefit the host plant by translocate phosphorus through a wide network of external hyphae and maximize the capability of the root system to absorb phosphorus. Colonization of AMF greatly increased chlorophyll content and root activity. The macro- and micro nutrient contents *i.e.* N, P, K, S, Ca, Cu, Fe, Mn, Mg, and Zn was also improved in roots (Chen *et al.* 2017) similarly application of compost and AMF significantly improved plant growth, stomatal conductance and chlorophyll fluorescence compared to infected and non-infected controls (Rahou *et al.* 2021).

Conclusion:

The present experiment showed that all the inoculum levels (100, 150, 200 and 400 sporocarps/kg soil) of *Glomus fasciculatum* have growth stimulating effects on *Capsicum annum*. Our results clearly suggest that in *Capsicum annum* var. Pusa Jwala, inoculation of AM fungi (100, 150 and 200 sporocarps/kg soil) showed better vegetative growth of all the plants while 400 sporocarps/kg soil inoculation showed the most promising and synergistic effects on vegetative growth of *Capsicum annum*. Colonization of AMF greatly increased chlorophyll content and NPK content of *Capsicum annum* at different observation periods.

References:

- Ait Rahou, Y., Ait-El-Mokhtar, M., Anli, M., Boutasknit, A., Ben-Laouane, R., Douira, A., ... & Meddich, A. (2021). Use of mycorrhizal fungi and compost for improving the growth and yield of tomato and its resistance to *Verticillium dahliae*. *Archives of Phytopathology and Plant Protection*, 54(13-14), 665-690.
- Al-Karaki, G. N. (2006). Nursery inoculation of tomato with arbuscular mycorrhizal fungi and subsequent performance under irrigation with saline water. *Scientia horticultrae*, 109(1), 1-7.
- Al-Karaki, G. N., McMichael, B. and Zak, J. (2004). Field response of wheat to arbuscular mycorrhizal fungi and drought stress. *Mycorrhiza*, 14:263-269.
- Anli, M., Symanczik, S., El Abbassi, A., Ait-El-Mokhtar, M., Boutasknit, A., Ben-Laouane, R., ... & Meddich, A. (2021). Use of arbuscular mycorrhizal fungus *Rhizoglyphus irregularis* and compost to improve growth and physiological responses of *Phoenix dactylifera* 'Boufgouss'. *Plant Biosystems-An International Journal*

Dealing with all Aspects of Plant Biology, 155(4), 763-771.

- Bona, E., Todeschini, V., Cantamessa, S., Cesaro, P., Copetta, A., Lingua, G., ... & Massa, N. (2018). Combined bacterial and mycorrhizal inocula improve tomato quality at reduced fertilization. *Scientia Horticulturae*, 234, 160-165.
- Charron G, Furlan V, Bernier-Cordou M, Doyon G (2001). Response of onion plants to arbuscular mycorrhizae. 1. Effects of inoculation method and phosphorus fertilization on biomass and bulb firmness. *Mycorrhiza* 11: 187-197.
- Chen, S., Zhao, H., Zou, C., Li, Y., Chen, Y., Wang, Z., Jiang, Y., Liu, A., Zhao, P., Wang, M. and Ahammed, G. J. (2017). Combined Inoculation with multiple arbuscular mycorrhizal fungi improves growth, nutrient uptake and photosynthesis in cucumber seedlings. *Frontiers Microbiology*, 8:1-11.
- Cooper, K. M. and Tinker, P. B. (1978). Translocation and transfer of nutrients in vesicular arbuscular mycorrhizas. Uptake and translocation of phosphorus, zinc and sulphur. *New Phytologist*, 81:43-52.
- Gerdemann, J. W. and Nicolson, T. H. (1963). Spores of mycorrhizal Eadogone. species extracted from soil by wet sieving and decanting. *Transaction of British Mycology Society*, 46: 235-244.
- Giri, B., Kapoor, R. and Mukerji, K. G. (2003). Influence of arbuscular mycorrhizal fungi and salinity on growth, biomass and mineral nutrition of *Acacia auriculiformis*. *Biology and Fertility of Soils*, 38: 170-175.
- Habte, M., Zhang, Y. C. and Schmitt, D. P. (1999). Effectiveness of *Glomus* species in protecting white clover against nematode damage. *Canadian Journal of Botany*, 77:135-139.
- Ibrahim, M. S., Faieza, A. A., Sapuan, S. M. and Tahir, S. M. (2011). Effect of filler loading and coupling agent on tensile and impact properties of polypropylene with oil palm ash composites. *Key Engineering Material*, 471-472:1130-1135.
- Menendez, J. A., Menendez, E. M., Iglesias, M. J., Garcia, A. and Pis, J. J. (1999). Modification of the surface chemistry of active carbons by means of microwave-induced treatments. *Carbon*, 37:1115-1121.
- Ortas, I., Kaya, Z., & Cakmak, I. (2001). Influence of arbuscular mycorrhizae inoculation on growth of maize and green pepper plants in phosphorus-and zinc-deficient soil. In *Plant Nutrition* (pp. 632-633). Springer, Dordrecht.
- Ortas, I., Sari, N., Akpinar, Ç., & Yetisir, H. (2011). Screening mycorrhiza species for plant growth, P and Zn uptake in pepper seedling grown under greenhouse conditions. *Scientia Horticulturae*, 128(2), 92-98.
- Phillips, J. M. and Hayman, D. S. (1970). Improved procedure for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Transactions British Mycological Society*, 55: 158-161.
- Plenchette, C., Fortm, J. A. and Furlan, V. (1983): Growth response of several plant species to mycorrhizae in a soil of moderate P fertility. I. Mycorrhizal dependency under field conditions. *Plant Soil*, 70: 199- 209.

Stutz, J.C., Coperman, R., Martin, C.A. and Morton, J. B. (2000). Patterns of species composition and distribution of arbuscular mycorrhizal fungi in arid regions of southwestern North America and Namibia, *South African Journal of Botany*, **78**: 237-245.

Thilagar, G. and Bagyaraj, D. J. (2013). Influence of different Arbuscular Mycorrhizal fungi on growth and yield of chilli. *Springer*, **85**(1):71-75.

UNDER PEER REVIEW