

## Effect Of Organic Management on Growth, Yield, and Economics of Greengram, Pigeon Pea, and Sunflower Under Rainfed Conditions

### Abstract

A field experiment was conducted during the harif season of 2021, to assess the effect of organic management on the growth, yield, and economics of greengram, pigeon pea, and sunflower under rainfed conditions. The experiment was laid out in a randomized complete block design with three treatment combinations viz., Organic (recommended 100 % N equivalent FYM), Integrated (recommended 25 % N equivalent FYM + 75 % N and 100 % P and K through chemical fertilizers) and Control (recommended 100 % NPK through chemical fertilizer) replicated nine times. Based on the results, it was found that the growth, yield attributes, and yield were significantly higher with organic treatment.

**Keywords:** Organic, Rainfed, Pigeon pea, Greengram, Sunflower

### 1. Introduction

Oilseeds are India's second most important agricultural product after grains. It accounts for 13% of the gross cropped area and ~~they contribute~~ around 3 % of the gross national product (GNP) and 10 % of all the agricultural commodities (DOD, 2021). Despite being one of the major producers of oilseeds, India, with its large population, is unable to supply the demand for edible oils. India is paying substantial foreign currency to import edible oils to meet domestic demand.

Sunflower (*Helianthus annuus* L.) is a popular oilseed crop in India due to its ability to adapt to a variety of agro-climatic zones and cropping patterns. Sunflower is grown on 2.25 lakh hectares in the country, with a yield of 2.28 lakh tonnes and a productivity of 1,011 kg ha<sup>-1</sup> (Indiastat, 2021). Karnataka is the country's most prolific producer. Telangana covers 7,000 hectares, with a yearly yield and productivity of 16,390 tonnes and 2,342 kg ha<sup>-1</sup>, respectively (Indiastat, 2021).

Pulses are a strong source of protein in the diet and can help satisfy the demands of a rapidly rising population. The output of pulses is exceedingly low, which has become a difficult challenge in light of our country's rapidly rising population. Pulses ~~contains~~ contain approximately 21-25% protein and it also contains a limited amount of essential amino acids such as tryptophan, methionine, lysine, and cysteine. (Tiwari and Singh, 2012). Pulses contain a significant quantity of lysine. ~~As a~~ Besides the nutritional benefit, pulses can potentially fix atmospheric nitrogen symbiotically in the soil through root nodules, which helps them maintain and improve soil fertility.

Greengram (*Vigna radiata* L.) popularly known as mungbean is grown on 51.3 lakh hectares in India, with a total yield of 30.85 lakh tonnes and a productivity of 601 kg ha<sup>-1</sup> (Indiastat, 2021). The top ~~greengram-green gram-producing~~ producing states are Rajasthan, Maharashtra, Andhra Pradesh, Gujarat, Bihar, and Karnataka. Greengram covers 75,000 ha in Telangana, with yield and productivity of 38,050 tonnes and 507 kg ha<sup>-1</sup>, respectively (Indiastat, 2021). It requires less water than other crops because of its shorter duration. In the harif and summer seasons, it is commonly planted as an intercrop, mixed crop, or single crop. Among grain legumes, it is one of India's most important conventional pulse crops. It is only second to chickpeas in terms of productivity. Its seed is tastier, more nutritious, easier to digest, and less flatulent than other pulses growing in that particular region.

Pigeon pea (*Cajanus cajan*) is India's second most important pulse crop after chickpea, commonly known as ~~redgram~~ red gram (Arhar or Tur) in local languages. It is grown on 4.72 million hectares in India, with a yield of 4.31 million tonnes and a productivity of 914 kg ha<sup>-1</sup> (Indiastat, 2021). Telangana has a total area of 3.25 lakh hectares with an annual production and productivity of 2.52 lakh tonnes and 775 kg ha<sup>-1</sup>, respectively (Indiastat, 2021). As it is an important nitrogen-fixing crop, it is widely grown for soil enrichment in rainfed areas. Its deep penetrating roots help move nutrients from deeper soil layers to the surface. Effective rhizobium nodulation can ~~an~~ average up to fix up to 160-200 kg nitrogen ha<sup>-1</sup> year<sup>-1</sup>.

Climate change and variability are a considerable threat to agricultural communities, particularly in India. This threat includes the likely increase ~~of~~ in temperature, extreme weather conditions, increased water stress and drought, and desertification.

Traditional crop production methods may be jeopardised by seasonal differences in weather events, either owing to a lack of water or a surplus of water, resulting in erosion. Soil stability will become increasingly important in order to store water in the soil profile, withstand extreme weather events, and reduce soil losses. These changes will bring new challenges to farmers. Farmers ~~require~~ ~~require~~ a new production system to assist them in adapting to these changing circumstances. Organic farming is one such alternative that has the potential to both mitigate and adapt to climate change, especially in rainfed agriculture. Organic agriculture is claimed to be the most sustainable approach ~~in food in food~~ production. It emphasizes recycling techniques and low external input and high output strategies. It is based on enhancing soil fertility and diversity at all levels and makes soils less susceptible to erosion (Niggli et al., 2007).

Scientific investigations are restricted in their capacity to reveal the productivity levels, stability, and profitability that may be obtained through pure organic sources of nutrient delivery in comparison to inorganic sources. It is critical to design robust feasible and compatible packages of nutrient management using organic sources for varied crops and cropping systems to fit local circumstances while being economically viable.

## 2. Materials and Methods

The field experiment was conducted in Gungal (around 43 km from Hyderabad) Research Farm (17.08°N latitude and 78.66°E longitude and 542 m above ~~msl~~MSL) of ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India, during ~~harif~~ Kharif 2021. The farm represents a semi-arid tropical region with a mean annual temperature of 25.58 °C and rainfall of 835.7 mm ~~was~~ received during the crop period ~~in of~~ 52 rainy days. The soil of the experimental field was clay loam in texture, with soil pH 6.51, low in organic carbon (0.43%), available nitrogen (179.0 kg/ha), high in available P<sub>2</sub>O<sub>5</sub> (24.7 kg/ha) and medium in available K<sub>2</sub>O (218.1 kg/ha). Experiment ~~The experiment~~ was laid out in a randomized complete block design with three treatment combinations viz., Organic (recommended 100 % N equivalent FYM), Integrated (recommended 25 % N equivalent FYM + 75 % N and 100 % P and K through chemical fertilizers) and Control (recommended 100 % NPK through chemical fertilizer) replicated nine times. In the plots under organic management, farmyard manure was thoroughly incorporated into 15 cm surface soil on the recommended N equivalent basis to all ~~the~~ three crops at two weeks before sowing of crops and the P requirement was supplemented by rock phosphate. The ~~initial~~ ~~initially~~ available K status of ~~the~~ experimental site was high hence no potassium was applied. In the plots under integrated management, 25% of ~~the~~ equivalent recommended N was applied through farmyard manure. The remaining 75% N and 100% P and K ~~is were~~ applied through chemical fertilizers. The plots under inorganic management ~~were~~ applied ~~with~~ ~~with~~ ~~recommended~~ ~~recommended~~ dose of chemical fertilizers (20:50 kg N and P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> for pigeon pea ~~and~~ ~~nd~~ ~~green~~ ~~green~~ ~~gram~~; 60:60:30 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup> for sunflower). The quantity of manure and fertilizer ~~applied at~~ ~~applied~~ field ~~are is~~ represented ~~at in~~ ~~Table~~ 1. The FYM used in the field were analysed for N, P, K and other micronutrients. On ~~an~~ average it contains 0.5% N, 0.27% P, 0.4% K, 452 ppm Fe, 228.7 ppm Mn, 143.1 ppm Zn and 27.9 ppm Cu. ~~Further~~ ~~Further~~ this FYM was treated with *Trichoderma viridae viridian* at 2.5 kg ha<sup>-1</sup> and incubated in the soil for about 20 days as ~~a~~ prophylactic measure against various ~~s soil~~ ~~bor~~ ~~soil~~ ~~borne~~ diseases. The sources of fertilizers for NPK were urea (46% N), SSP (16% P<sub>2</sub>O<sub>5</sub>, 12% S) and MOP (60% K<sub>2</sub>O). Varieties used in ~~the~~ experiment are WGG-37 (~~green~~ ~~green~~ ~~gram~~), PRG-158 (pigeon pea) and DRSH-1 (sunflower). Sowing was done ~~during at~~ ~~the~~ first fortnight of July using healthy, well-filled, and matured seeds of pigeon ~~pea~~, ~~green~~ ~~green~~ ~~gram~~ ~~gram~~, and sunflower at 15, 20, and 6 kg ha<sup>-1</sup> seed rate. For pigeon ~~pea~~, ~~green~~ ~~green~~ ~~gram~~ ~~gram~~, and sunflower, seeds were sown by hand dibbling @ two seeds hill<sup>-1</sup> at a depth of 3-5 cm, with 90 cm, 30 cm, and 60 cm between rows and 20 cm, 10 cm, and 30 cm between plants within the row, respectively. Thinning and gap filling were completed after 15 DAS, with one healthy seedling hill<sup>-1</sup> remaining. In conformity with organic standards, no chemical herbicides, insecticides, or fungicides were employed in the organically managed plots. Manual weeding was used once, followed by two ~~hoeing~~ ~~hoeing~~ using a manually driven wheel-hoe. As a preventative treatment, Azadirachtin (*Azadirachta indica* - based formulation) was applied against insect pests at 20-25 days intervals during crop growth. Crop-specific prescribed herbicides and pesticides were applied in the control plots to manage weeds, insect pests, and diseases. Recommended integrated pest management (IPM) modules were applied in the integrated management plots. Greengram was harvested first due to its short duration, followed by sunflower, and finally pigeon pea. Sunflower is harvested by cutting the heads after the green base of the head has

become yellow and eventually brown while pigeon pea and greengram and green gram are harvested by picking the pods when they are brown and dried. The data on growth and yield attributes were recorded from randomly selected five plants in each plot and seed and haulm/stalk yield was recorded from the plot and converted on a hectare basis.

Table 1: Application rates of organic manure and mineral fertilizers in different treatments

| Treatments | Crops      | Manure<br>(t ha <sup>-1</sup> ) | Chemical fertilizer<br>(kg ha <sup>-1</sup> ) |      |      | Rock phosphate<br>(kg ha <sup>-1</sup> ) |
|------------|------------|---------------------------------|---|------|------|--|
|            |            |                                 | N   | P    | K    |  |
| Organic    | Greengram  | 4.4                             | 0.0   | 0.0  | 0.0  | 167.0                                    |
|            | Pigeon pea | 4.4                             | 0.0   | 0.0  | 0.0  | 167.0                                    |
|            | Sunflower  | 13.3                            | 0.0   | 0.0  | 0.0  | 0.0                                      |
| Integrated | Greengram  | 3.3                             | 15.0  | 50.0 | 0.0  | 0.0                                      |
|            | Pigeon pea | 3.3                             | 15.0  | 50.0 | 0.0  | 0.0                                      |
|            | Sunflower  | 10.0                            | 45.0  | 60.0 | 30.0 | 0.0                                      |
| Inorganic  | Greengram  | 0.0                             | 20.0  | 50.0 | 0.0  | 0.0                                      |
|            | Pigeon pea | 0.0                             | 20.0  | 50.0 | 0.0  | 0.0                                      |
|            | Sunflower  | 0.0                             | 60.0  | 60.0 | 30.0 | 0.0                                      |

### 3. Results and Discussion

#### 3.1. Growth Parameters

##### 3.1.1. Plant height (cm)

In greengram green gram, application of organic treatment has recorded significantly highest plant height at harvest stage (41.6 cm) compared to rest of the treatments which was followed by integrated treatment (37.7 cm) and it was found be at par with control (37.1 cm). In pigeon pea under organic plots being on par with those under integrated plots (132.6 cm) recorded significantly higher plant height (137.6 cm) compared to control (127.1 cm). However, the integrated treatment is in turn at par with control. In sunflower, the application organic source of nutrient-nutrients results in a significantly highest plant height (152.3 cm) over the control and which is statistically at par with integrated treatment (143.9 cm). However, this integrated treatment was also statistically at par with the control (138.8 cm).

Significantly highest plant height was recorded with an organic source of nutrient-nutrients compared to integrated and control. Organically treated crops outperformed all other treatments because organic manure enhances cell division, cell elongation, and chlorophyll content of leaves, which improves photosynthetic activity and consequently plant height. While the plants with control were the smallest in stature, this might be owing to a lack of nutrients for crop growth. The present results were in accordance with the findings of Mishra et al. (2015), Mohbe et al. (2015) and Muhsin et al. (2021).

##### 3.1.2. Number of branches

In greengram green gram, the highest number of branches at harvest (8.9) was recorded with organic treatment and it was at par with integrated treatment (8.8) and significantly highest over control (8.6). However, integrated treatment is statistically similar with to the control. At harvest pigeon pea under the organic source of nutrient-nutrients being on par with those under an integrated source of nutrient-nutrients (14) recorded significantly the highest number of branches (14.3) compared to control (13.9). however, the integrated treatment where statistically at par with the control.

Number-The number of branches found be is significantly highest in the organic treatment this is because increased availability of nutrients due to enhanced microbial activity with the application of organic manures results in increasing-increased of cellular activity and cell division results in the greater number of branches. The similar-Similar results are found in Mishra et al. (2015), Rao et al. (2018) and Muhsin et al. (2021).

##### 3.1.3. Dry matter accumulation (kg ha<sup>-1</sup>)

In ~~greengram~~ green gram, the highest dry matter accumulation at harvest was recorded with organic treatment (2,463.3 kg ha<sup>-1</sup>) over control (2,201.2 kg ha<sup>-1</sup>) ~~and~~ however, this organic treatment was at par with integrated treatment (2,343.3 kg ha<sup>-1</sup>). At harvest, pigeon pea under the organic source of ~~nutrient-nutrients~~ being on par with those under an integrated source of ~~nutrient~~ nutrients (4,572.0 kg ha<sup>-1</sup>) recorded significantly the highest dry matter accumulation (4,842.5 kg ha<sup>-1</sup>) compared to control (4,326.5 kg ha<sup>-1</sup>). However, the integrated treatment was statistically at par with the control. ~~Similar-A similar~~ trend was observed with all other growth stages viz., pod formation and at harvest. In sunflower application, organic ~~source-sources~~ of ~~nutrient-nutrients~~ results resulted in significantly higher dry matter accumulation at harvest (4,666.4 Kg ha<sup>-1</sup>) over the control and which is statistically at par with integrated treatment (4,428.2 Kg ha<sup>-1</sup>). However, this integrated treatment was also statistically at par with the control (4,325.3 Kg ha<sup>-1</sup>).

The total dry matter production is significantly higher in organic plots which ~~is~~ are followed by integrated and control. This could be mainly due to higher plant height and higher leaf area maintained throughout the crop phase resulting in enhanced carbohydrate synthesis, which finally led to higher dry matter production. The results are in close conformity with Daniel et al. (2013), Ullasa et al. (2013) and Rao et al. (2018).

### 3.2. Yield attributes

#### 3.2.1. Number of pods plant<sup>-1</sup>

~~Greengram~~ Green gram has recorded a significantly higher number of pods plant<sup>-1</sup> in the plots where organic treatment was applied (22), which was on par with the pod number ~~resulted~~ resulting from the due application of integrated treatment (20). Lower The lower number of pods was with unsprayed treatment (19.7). whereas, Pigeon pea under organic plots ~~were~~ was recorded with a significantly highest number of pods plant<sup>-1</sup> (36.3) compared to crops under integrated plots (33.4) which is statistically at par. However, this integrated treatment is at par with control (32.4).

This might be due to an increase in the availability of nearly all plant necessary nutrients, which promotes vegetative growth and hence increases photosynthetic activity, as well as the translocation and storage of photosynthates in economic sinks. Our results ~~are in consonance~~ align with the findings of Mishra et al. (2015), Muhsin et al. (2021) and Rao et al. (2021).

#### 3.2.2. Capitulum diameter (cm)

Data revealed that the capitulum diameter of sunflower did not vary significantly due to the different production systems applied. Among all the treatments applied, organic treatment ~~was~~ recorded the highest capitulum diameter (15.1 cm) which is statistically at par with both integrated treatment (14.9 cm) and control (13.9 cm).

Higher capitulum diameter was recorded with the organic production system this is because organic manures are the ~~store~~ storehouse of several macro and micronutrients that are released during the process of mineralization by stimulating the activity of microorganisms to make the plant nutrients readily available to crops therefore a synthesis of more photosynthates may take place which might be resulted in better partitioning of photosynthates from source to sink and improved the yield attributes like capitulum diameter. This was supported by Chaitra and Sujith (2021) and Muhsin (2021).

#### 3.2.3. Number of filled seeds pod<sup>-1</sup>

There was no significant difference between the treatments. In ~~greengram~~ green gram, among different treatments tried, a higher number of filled Seeds Pod<sup>-1</sup> was noticed in the plots where organic treatment was applied (11.22), which was on par with seed number ~~resulted~~ resulting due application of both integrated treatment (11) and control (10.67). In pigeon pea, among different treatments tried, a higher number of filled seeds pod<sup>-1</sup> was noticed in the plots where organic treatment was applied (4.22), which was on par with seed number ~~resulted~~ resulting in the due application of both integrated treatment (4) and control (3.89).

In ~~sunflowers~~ sunflowers, ~~several~~ number of filled seeds head<sup>-1</sup> differed significantly among different production systems. Crops ~~under~~ under organic source of nutrient ~~recorded~~ higher number of filled seeds head<sup>-1</sup> (630.3) compared to an integrated source of nutrient and control. However, the integrated source of ~~nutrient-nutrients~~ (583.3) was on par with the control (545.4).

~~Combined~~ The combined effect of improved growth parameters (plant height, leaf area and dry matter accumulation) through efficient metabolic activity, increased photosynthetic rate and a steady supply of photosynthates from source to sink might have led to

~~the~~ maximum expression of yield attributing characters. ~~The similar~~Similar results are found with Chaithra and Sujith (2021) and Muhsin (2021).

#### 3.2.4. Test weight (g)

Application of different production systems on test weight among all ~~the~~ three crops ~~are is~~found to be non-significant. The highest test weight of ~~greengram-green gram~~seeds was recorded in organic treatment (30.2 g) which remains at par with both integrated treatment (30 g) and control (29.6 g) and the highest test weight of pigeon pea seeds was recorded in organic treatment (81.6 g) which being on par with both integrated treatment (80 g) and control (80.6 g). In sunflower, among all the treatments, the higher test weight was recorded in organic treatment (44.6 g) which ~~were was~~statistically at par with both integrated treatment (44.4 g) and control (43.9 g).

UNDER PEER REVIEW

Table 2: Effect of different production systems on Greengram, Pigeon pea and Sunflower growth at harvest

| Treatments  | Plant height (cm) |            |           | No. of branches |            | Dry matter accumulation (kg ha <sup>-1</sup> ) |            |           |
|-------------|-------------------|------------|-----------|-----------------|------------|--|------------|-----------|
|             | Greengram         | Pigeon pea | Sunflower | Greengram       | Pigeon pea | Greengram                                      | Pigeon pea | Sunflower |
| Organic     | 41.6              | 149        | 152.3     | 8.9             | 14.3       | 2463.3   | 4842.5     | 4666.4    |
| Integrated  | 37.7              | 142.9      | 143.9     | 8.6             | 14         | 2298.9   | 4572       | 4428.2    |
| Control     | 37.1              | 138.7      | 138.8     | 8.3             | 13.9       | 2162.2   | 4326.5     | 4325.3    |
| SEm±        | 1.1               | 3.3        | 3.9       | 0.2             | 0.4        | 97.3   | 129.7      | 109.6     |
| CD (P=0.05) | 3.1               | 9.9        | 11.6      | 0.6             | 1.1        | 291.7  | 388.8      | 329.0     |

Table 3: Effect of different production systems on yield attributes in Greengram, Pigeon pea and Sunflower

| Treatments  | Pods plant <sup>-1</sup> |            |           | Filled seeds pod <sup>-1</sup> /head <sup>-1</sup> |            |           | Test weight (g) |            |           |
|-------------|--------------------------|------------|-----------|--|------------|-----------|-----------------|------------|-----------|
|             | Greengram                | Pigeon pea | Sunflower | Greengram  | Pigeon pea | Sunflower | Greengram       | Pigeon pea | Sunflower |
| Organic     | 22                       | 36.3       | 15.5      | 11.22  | 4.2        | 630.3     | 30.2            | 81.6       | 44.6      |
| Integrated  | 20                       | 33.4       | 14.9      | 11   | 4          | 583.3     | 30              | 81         | 44.4      |
| Control     | 19.7                     | 32.4       | 13.8      | 10.67  | 3.9        | 545.4     | 29.6            | 80.6       | 43.9      |
| SEm±        | 0.7                      | 1.1        | 0.5       | 0.3  | 0.2        | 14.6      | 0.2             | 0.2        | 0.3       |
| CD (P=0.05) | 2.2                      | 3.4        | 1.5       | NS   | NS         | 43.9      | NS              | NS         | NS        |

Formatted Table

### 3.3. Yield

#### 3.3.1. Grain yield ( $\text{Kg ha}^{-1}$ )

The use of an organic source of nutrient resulted in a significantly highest grain yield of greengram-green gram ( $673 \text{ kg ha}^{-1}$ ) over control and remains at par with the integrated source of nutrient ( $661 \text{ kg ha}^{-1}$ ) and this integrated treatment was statistically at par with control ( $630 \text{ kg ha}^{-1}$ ). The use of organic treatment resulted in a significantly highest seed yield of Pigeon pea ( $383 \text{ kg ha}^{-1}$ ) compared to integrated treatment ( $358 \text{ kg ha}^{-1}$ ). However, which remains at par with ~~the~~ integrated treatment. Lowest-The lowest seed yield among the treatments was recorded with control ( $323 \text{ kg ha}^{-1}$ ). The higher seed yield in sunflower was recorded in organic treatment ( $944 \text{ kg ha}^{-1}$ ) which was higher than the remaining treatments which ~~was~~ however comparable with that resulted grain yield of integrated treatment ( $920 \text{ kg ha}^{-1}$ ) but significantly higher than with control ( $835 \text{ kg ha}^{-1}$ ).

Higher seed yield in organic treatment might be attributed to an increase in the availability of practically all plant necessary nutrients due to the transfer of photosynthates produced under the effect of the organic nutrient source. Furthermore, photosynthate translocation and accumulation in economic sinks improved yield characteristics, chlorophyll content, and nitrate reductase activity, resulting in higher seed production. ~~The similar~~ Similar results are found by Pradeepet al. (2018), Chongreet al. (2019) and Muhsin et al. (2021).

#### 3.3.2. Straw yield ( $\text{Kg ha}^{-1}$ )

Greengram under organic plots being on statistically par with those under integrated plots ( $1879 \text{ kg ha}^{-1}$ ) recorded significantly higher plant height ( $1908 \text{ kg ha}^{-1}$ ) compared to control ( $1797 \text{ kg ha}^{-1}$ ). However, the integrated treatment is in turn at par with control. Pigeon pea under organic plots being on par with those under integrated plots ( $2521 \text{ kg ha}^{-1}$ ) recorded significantly higher stalk yield ( $2647 \text{ kg ha}^{-1}$ ) compare to crops under control ( $2402 \text{ kg ha}^{-1}$ ). However, the crops under the integrated treatment will be statistically at par with the control. Among the different sources of nutrients tried, incorporation of organic treatment recorded the significantly higher stalk yield ( $2484 \text{ kg ha}^{-1}$ ) in sunflower which was however statistically at par with the integrated source of nutrient ( $2434 \text{ kg ha}^{-1}$ ) and the lowest stalk yield ~~were was~~ recorded under control ( $2216 \text{ kg ha}^{-1}$ ).

The higher stalk yield due to the application of organic treatment might be attributed to improved vegetative growth and greater dry matter production as a result of the availability of all plant nutrients and improved soil physical qualities. The present result was in accordance with the findings of Rekha et al. (2018), Pradeepet al. (2018) and Muhsin et al. (2021).

#### 3.3.3. Harvest Index (%)

In greengram-green gram, among the different sources of nutrients tried, incorporation of organic treatment recorded the higher harvest index (26.1 %) which is at par with both integrated source of nutrient (26.0 %) and control (26.0 %). Similarly, at pigeon pea, incorporation of organic treatment recorded ~~the~~ higher harvest index (12.6 %) which is at par with both integrated source of nutrient (12.5 %) and control (11.9 %) and ~~at~~ sunflower, incorporation of organic treatment recorded the higher harvest index (27.6 %) which were with integrated treatment (27.5 %) and statistically at par with control (27.4 %).

### 3.4. Economics

#### 3.4.1. Gross returns ( $\text{Rs. ha}^{-1}$ )

In greengram-green gram, application of an organic source of nutrients fetched the highest gross returns ( $51160 \text{ Rs ha}^{-1}$ ) among all the ~~treatment-treatment~~ in comparison. The next best treatment was integrated treatment which resulted in gross returns of  $50088 \text{ Rs ha}^{-1}$ . The lowest gross returns were recorded with control ( $47750 \text{ Rs ha}^{-1}$ ). In pigeon pea application of an organic source of nutrients fetched significantly highest gross returns ( $26818 \text{ Rs ha}^{-1}$ ) among all the treatment in comparison. The next best treatment was integrated treatment which resulted in gross returns of  $25159 \text{ Rs ha}^{-1}$ . The lowest gross returns were recorded with control ( $22772 \text{ Rs ha}^{-1}$ ). In sunflower application of an organic source of nutrients resulted the highest gross returns ( $58137 \text{ Rs ha}^{-1}$ ) among all the treatment in comparison. The next best treatment was integrated treatment which resulted in gross returns of  $56635 \text{ Rs ha}^{-1}$ . The lowest gross returns were recorded with control ( $51389 \text{ Rs ha}^{-1}$ ).

Since the organic produce grows purely under chemical free ~~condition~~ conditions, the quality of the ~~produce~~ product will be excellent compared to integrated and inorganic produce. So, to ~~compensate the~~ compensate for the high cost of

cultivation under organic due to high cost of FYM per kg of nutrient supply we would like add the premium of 15% for ~~greengram~~ green gram and pigeon pea organic produce and 30% for sunflower organic produce and sold at ~~fixed a fixed~~ premium price. Therefore, the gross returns of organically produced ~~greengram-green gram~~ were increased by 15% and results in the gross returns of 58546 ₹ ha<sup>-1</sup> compared to normal gross returns of 51160 Rs ha<sup>-1</sup>. Similarly, in pigeon pea and sunflower gross returns had increased to 30443 and 75210 ₹ ha<sup>-1</sup> respectively.

#### 3.4.2. Net returns (Rs ha<sup>-1</sup>)

Among the treatments, ~~the~~ application of ~~an~~ integrated source of ~~nutrient-nutrients~~ registered ~~highest~~ the highest net returns (26614 Rs ha<sup>-1</sup>) followed by inorganic source of nutrients (control) which gained a net return of 26135 Rs ha<sup>-1</sup>. The lowest net returns were noticed in the treatment where ~~an~~ organic source of ~~nutrient-nutrients was~~ applied (25075 Rs ha<sup>-1</sup>). Among the different treatments in ~~pigeon-pea~~ pigeon pea, ~~the~~ application of ~~an~~ integrated source of ~~nutrient-nutrients~~ ~~registered~~ registered the highest net returns (1885 Rs ha<sup>-1</sup>) followed by ~~the~~ inorganic source of nutrients (control) which gained a net return of 1357 Rs ha<sup>-1</sup>. The lowest net returns were noticed in the treatment where ~~an~~ organic source of ~~nutrient-nutrients was~~ applied (933 Rs ha<sup>-1</sup>). Among the different ~~treatment in sunflow~~ treatments ~~the lowerer~~, application of ~~an~~ integrated source of ~~nutrient-nutrients~~ registered the highest net returns (30789 Rs ha<sup>-1</sup>) followed by ~~the~~ inorganic source of nutrients (control) which gained a net return of 30403 Rs ha<sup>-1</sup>. The lowest net returns were noticed in the treatment where organic ~~source-sources~~ of ~~nutrient-nutrients were~~ applied (22502 Rs ha<sup>-1</sup>). ~~The similar~~ Similar results are found by Arbad and Sayad (2011), Choudhary et al. (2013) and Verma et al. (2018).

Organically produced ~~greengram-green gram~~ which ~~is~~ sold at a ~~15~~ 15% premium will have net returns of 32411 Rs ha<sup>-1</sup> compared to normal gross returns of 25075 Rs ha<sup>-1</sup>. ~~Similarly, Similarly,~~ ~~pigeon-pea~~ pigeon pea and sunflower net returns had increased to 4,506 and 39543 Rs ha<sup>-1</sup> respectively.

#### 3.4.3. ~~Benefit-cost~~ Benefit-cost (B:C) ratio

In ~~greengram~~ green gram, higher ~~B:C~~ B: C was recorded with the control plots (2.21) followed by integrated plots (2.13). Whereas, ~~lower a lower~~ B:C ratio was recorded with organic plots (1.96). In pigeon pea, higher B:C was recorded with the integrated plots (1.08) followed by control plots (1.06). Whereas, lower B:C ratio was recorded with organic plots (1.04). In sunflower, higher B:C was recorded with the control plots (2.45) which was followed by integrated plots (2.19). Whereas, lower B:C ratio was recorded with organic plots (1.63).

Organically produced ~~greengram-green gram~~ which sold at 15% premium, B: C ratio were increased to 2.24 compared to normally sold pigeon pea which is having 1.96 B: C ratio. Similarly, pigeon pea and sunflower have B: C ratio of 1.18 and 2.11 respectively. This was supported by Meena and Sharma (2013), Verma et al. (2018) and Somalraju et al. (2021).

UNDER PEER REVIEW




Table 4: Effect of different production systems on yield in Greengram, Pigeon pea and Sunflower

| Treatments  | Seed yield (kg ha <sup>-1</sup> ) |            |           | Stalk yield (kg ha <sup>-1</sup> ) |            |           | Harvest Index (%) |            |           |
|-------------|-----------------------------------|------------|-----------|------------------------------------|------------|-----------|-------------------|------------|-----------|
|             | Greengram                         | Pigeon pea | Sunflower | Greengram                          | Pigeon pea | Sunflower | Greengram         | Pigeon pea | Sunflower |
| Organic     | 673                               | 383        | 944       | 1908                               | 2647       | 2484      | 26.1              | 12.6       | 27.6      |
| Integrated  | 661                               | 358        | 920       | 1879                               | 2521       | 2434      | 26                | 12.5       | 27.5      |
| Control     | 630                               | 323        | 835       | 1797                               | 2402       | 2216      | 26                | 11.9       | 27.4      |
| SEM±        | 14                                | 9          | 25        | 36                                 | 61         | 65        | 0.5               | 0.3        | 0.8       |
| CD (P=0.05) | 42                                | 29         | 74        | 108                                | 184        | 196       | NS                | NS         | NS        |

Table 5: Effect of different production systems on economics in Greengram, Pigeon pea and Sunflower

| Treatments | Cost of cultivation (₹ ha <sup>-1</sup> ) |            |           | Gross returns (₹ ha <sup>-1</sup> ) |            |           | Net returns (₹ ha <sup>-1</sup> ) |            |           | Benefit Cost ratio |            |           |
|------------|---|------------|-----------|-------------------------------------|------------|-----------|-----------------------------------|------------|-----------|--------------------|------------|-----------|
|            | Greengram                                 | Pigeon pea | Sunflower | Greengram                           | Pigeon pea | Sunflower | Greengram                         | Pigeon pea | Sunflower | Greengram          | Pigeon pea | Sunflower |
| Organic    | 26085                                     | 25885      | 35635     | 51160                               | 26818      | 58137     | 25075                             | 933        | 22502     | 1.96               | 1.04       | 1.63      |
| Integrated | 23474                                     | 23274      | 25846     | 50088                               | 25159      | 56635     | 26614                             | 1885       | 30789     | 2.13               | 1.08       | 2.19      |
| Control    | 21615                                     | 21415      | 20986     | 47750                               | 22772      | 51389     | 26135                             | 1357       | 30403     | 2.21               | 1.06       | 2.45      |

Table 6: Effect of different production systems on economics in Greengram, Pigeon pea and Sunflower when sold at premium for organic product:

| Treatments | Cost of cultivation (₹ ha <sup>-1</sup> ) |            |           | Gross returns (₹ ha <sup>-1</sup> ) |            |           | Net returns (₹ ha <sup>-1</sup> ) |            |           | Benefit Cost ratio |            |           |
|------------|---|------------|-----------|-------------------------------------|------------|-----------|-----------------------------------|------------|-----------|--------------------|------------|-----------|
|            | Greengram                                 | Pigeon pea | Sunflower | Greengram                           | Pigeon pea | Sunflower | Greengram                         | Pigeon pea | Sunflower | Greengram          | Pigeon pea | Sunflower |
| Organic    | 26085                                     | 25885      | 35635     | 58546                               | 30443      | 75210     | 32411                             | 4506       | 39543     | 2.24               | 1.18       | 2.11      |
| Integrated | 23474                                     | 23274      | 25846     | 50088                               | 25159      | 56635     | 26614                             | 1885       | 30789     | 2.13               | 1.08       | 2.19      |
| Control    | 21615                                     | 21415      | 20986     | 47750                               | 22772      | 51389     | 26135                             | 1357       | 30403     | 2.21               | 1.06       | 2.45      |

Formatted Table

Formatted Table

#### 4. Conclusion

Climate change is affecting us severely all over the world and farming is one of the main victims. There is a need to find a solution on which scientists are working relentlessly. Organic farming could be one of the best possible solutions to minimize the impact of climate change on farming. There has been an increase in awareness about organic farming in our country which has resulted in enhancement in area under organic farming. Pulses and oilseeds have to be cultivated under organic management practices as it fetches a better price comparatively which ultimately benefits the farmers as well as our economy. This investigation was carried out to compare the impact of organic, inorganic and integrated management practices on growth, yield and economics of greengram, pigeon pea and sunflower under rainfed conditions. It is concluded that greengram, pigeon pea and sunflower under organic farming have performed well in terms of growth, productivity and economics compared to integrated as well as inorganic farming.

#### 5. References

- Arbad, B.K. and Ismail, S. Effect of integrated nutrient management on soybean (*Glycine max*)-safflower (*Carthamus tinctorius*) cropping system. *Indian Journal of Agronomy*. 2011;56(4):340-345.
- Chaithra, M. and M Sujith, G. Influence of Farmyard Manure and Jeevamrutha on Growth and Yield of Sunflower. 2021.
- Chongre, S., Mondal, R., Biswas, S., Munshi, A., Mondal, R. and Pramanick, M. Effect of Liquid Manure on Growth and Yield of Summer Green Gram (*Vigna radiata* L. Wilczek). *Current Journal of Applied Science and Technology*. 2019;1-7.
- Choudhary, H.R., Sharma, O.P., Singh, R.K., Singh, K., Kumar, R. and Yadav, L. Influence of organic manures and chemical fertilizer on nutrient uptake, yield and profitability of mungbean [*Vigna radiata* (L.) Wilczek]. *Madras Agric J*. 2013;100(1-3):747-750.
- Directorate of oilseeds development. Oilseeds gross national product. 2021. [oilseeds.dac.gov.in](http://oilseeds.dac.gov.in)
- Indiastat. Area, production and productivity of greengram, pigeonpea and sunflower. 2021. <http://www.indiastat.com>
- Leo Daniel Amalraj, E., Praveen Kumar, G., Mir Hassan Ahmed, S.K., Abdul, R. and Kishore, N. Microbiological analysis of panchagavya, vermicompost, and FYM and their effect on plant growth promotion of pigeon pea *Cajanus cajan* (L.) in India. *Organic Agriculture*. 2013;3(1):23-29.
- Meena, R.S. and Sharma, S.K. Effect of organic and inorganic sources of nutrients on yield attributes, yield and economics of greengram [*Vigna radiata* (L.) Wilczek]. *Annals of Agri Bio Research*. 2013;18(3):306-308.
- Mishra, S.K., Parihar, S. and Namdeo, K. Effect of organic and inorganic sources of nutrients on growth, yield and economics of pigeonpea. *Annals of Plant and Soil Research*. 2015;17(4):374-376.
- Mohbe, Sandeep., Mishra, U.S. and Pandey, R.C. A study on organic manure on green gram [*Phaseolus radiata* (L.)] under rainfed condition of Chitrakoot area. *Trends in Biosciences*. 2015;8(23):6551-6554.
- Muhsin, S.J., Ramadhan, M.N. and Nassir, A.J. Effect of organic manure and tillage depths on sunflower (*Helianthus annuus* L.) production. In *IOP Conference Series: Earth and Environmental Science*. 2021;735(1):012070.
- Niggli, U., Schmid, H. and Fliessbach, A. Organic Farming and Climate Change. *Technical Paper*, International Trade Centre, Geneva. 2007;27.

- Pradeep, S., Ullasa, M.Y., Kumar Naik, A.H., Ganapathi and Divya, M. Effect of different organic nutrient management practices on growth, yield of pigeon pea (*Cajanus cajan* L. Millsp.) and soil properties. *Research Journal of Agricultural Sciences*. 2018;9(2): 352-357.
- Rao, D., Sachan, C.P., Pashwan, V.R. and Sakpal, A.V. Effect of organic manure, biofertilizer, phosphorus and nitrogen on growth, seed yield, seed quality attributes of green gram, *Vigna radiata* (L.). 2021.
- Rao, G.S., Immanuel, R.R., Raj, T.S. and Pushpanathan, K.R. Organic foliar nutrition influence on productivity of greengram. 2018.
- Rekha, K., Pavaya, R.P., Neha, C. and Patel, S. Effect of FYM, Phosphorus and PSB on Growth, Yield and Quality of Greengram [*Vigna radiata* (L.) Wilczek] on Loamy Sand. *International Journal of Bio-resource and Stress Management*. 2018;9(2):220-223.
- Somalraju, S., Goyal, G., Gurjar, L.S., Chaturvedi, M. and Singh, R. Effect of organic and inorganic fertilizer on the growth and yield of green gram (*Vigna radiata* L.). *The Pharma Innovation Journal*. 2021;10(12):1959-1962.
- Tiwari, B.K. and Singh, N. *Pulse chemistry and technology*. Royal Society of Chemistry. 2012
- Ullasa, M.Y., Sheshadri, T., Geetha, K.N. and Shankaralingappa, B.C. Standardization of best management practices for yield maximization in sunflower (*Helianthus annuus* L.). *Helia*, 2013;36(58): 95-100.
- Verma, S., Singh, A., Pradhan, S.S., Singh, J.P. and Verma, S.K. Effects of organic formulations and synthetic fertilizer on the performance of pigeonpea in eastern region of Uttar Pradesh. *Bangladesh Journal of Botany*. 2018;47(3): 467-471.

UNDER PEER REVIEW

