

## Original Research Article

### **INFLUENCE OF NUTRIENT CONCENTRATION ON GROWTH, YIELD AND QUALITY OF SPINACH (*Spinacia oleracea* L.) IN HYDROPONIC SYSTEM**

#### **ABSTRACT**

The present experiment entitled “**Influence of nutrient concentration on Growth, Yield and Quality of Spinach (*Spinacia oleracea* L.) in Hydroponic system**” was carried out in Shade net, Research Field, Department of Horticulture, SHUATS, Prayagraj during March- April, 2020. The experiment was laid out in complete randomized design (CRD), with eight treatments, replicated thrice with nutrient field technique hydroponics system and variety Mulayam of Spinach, the treatments were T<sub>1</sub> (NPK 6.25 ml/plant), T<sub>2</sub> (NPK 8.33 ml/plant), T<sub>3</sub> (NPK 10.41 ml/plant), T<sub>4</sub> (NPK 12.5 ml/plant), T<sub>5</sub> (NPK 14.58 ml/plant), T<sub>6</sub> (NPK 16.66 ml/plant), T<sub>7</sub> (NPK 18.75 ml/plant) and T<sub>0</sub> (NPK 0 ml/plant). From the present experimental findings, it is found that structure with treatment T<sub>4</sub> was found best in terms of Growth and yield parameters of spinach in NFT hydroponic system followed by structure with T<sub>3</sub> and T<sub>1</sub>. Maximum gross return (1871.30Rs) and net return (746.30 Rs) and maximum benefit cost ratio (1.66) was also observed in treatment T<sub>4</sub>.

**Keywords:** Spinach, Nutrient film techniques, Hydroponic and NPK.

#### **INTRODUCTION**

Spinach (*Spinacia oleracea* L.) commonly called as Spinach (English) and Palak in India is an important leafy vegetable of which the leaves and

tender shoots are consumed fresh or processed. It belongs to the family Chenopodiaceae. It is native of Central Asia, most probably of Persia (Iran).

Soil cultivation is practiced since centuries

as it contains nutrients formed during natural decay. However, it may take several years of building soil up with soil amendments to obtain the desired texture and composition. A new technique called soil-less culture commonly known as hydroponics has been developed to further improve the crop productivity in lesser space and time by controlling the supply of water and nutrients. Hydroponics comes from two Greek words *hydro* means water and *ponos* means labour or working with water. The word hydroponics was coined by Professor William Gericke in the early 1930s. Hydroponics is the technique of growing plants in nutrient enriched water with or without the use of an inert medium (aggregate culture) such as gravel, vermiculite, rockwool, peat moss, saw dust, coir dust, coconut fibre etc. A modern definition of hydroponics: A system where plants are grown in growth media other than natural soil. In this study the principle of Nutrient film technique (NFT) was applied. Nutrient film technique (NFT) is a closed hydroponic system. It is commonly used to grow leafy vegetables productions. NFT was developed by Dr. Alen Cooper in the mid-1960s. In this system, water or a nutrient solution recirculates throughout the entire system and enters the growth through a pump without a time control (Domingues *et al.*, 2012).

The aim of the current research was to evaluate the effect of different concentrations of nutrients on quantitative and qualitative traits of spinach grown in floating system.

Various experimental findings outline that leafy greens (lettuce, spinach, parsley, celery etc) can be successfully and easily grown in hydroponic systems Nisha *et al.* (2018). Therefore keeping the above facts in view the present study entitled "Production of Lettuce in Nutrient film technique (NFT) of hydroponic systems" was conducted in Department of Horticulture, SHUATS, Prayagraj.

## MATERIALS AND METHODS

The present Experiment was conducted in Complete Randomized Design (CRD), with eight treatments, of NPK replicated thrice with Spinach variety (Mulayam), in the Shade net, Research field,

Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during March, 2020 to April, 2020. Total number of treatments were eight viz. T<sub>1</sub> (NPK 6.25 ml/Plant), T<sub>2</sub> (NPK 8.33 ml/Plant), T<sub>3</sub> (NPK 10.41 ml/Plant), T<sub>4</sub> (NPK 12.5 ml/Plant), T<sub>5</sub> (NPK 14.58 ml/Plant), T<sub>6</sub> (NPK 16.66 ml/Plant), T<sub>7</sub> (NPK 18.75 ml/Plant) and T<sub>0</sub> (Control (NPK 0 ml/Plant)).

### Climatic condition in the experimental site:

The area of Prayagraj district comes under subtropical belt in the south east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C- 48°C and seldom falls as low as 4°C- 5°C. The relative humidity ranges between 20 to 94 %. The average rainfall in this area is around 1013.4 mm annually. However, occasional precipitation is also not uncommon during winter months.

## RESULTS AND DISCUSSION

The present investigation entitled "Influence of nutrient concentration on Growth, Yield and Quality of Spinach (*Spinacia oleracea* L.) in Hydroponic system" was carried out during March, 2020 to April, 2020 in Shade net, Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. The results of the present investigation, regarding the influence of different concentration of NPK on growth, yield and quality of Spinach in (NFT) hydroponic system, have been discussed and interpreted in the light of previous research work done in India and abroad. The experiment was conducted in Complete Randomized Design with 8 treatments, and three replications.

The results of the experiment are summarized below:

Comment [u1]: Move it to last paragraph of the introduction

Comment [u2]: Delete

Comment [u3]: Delete

## A. Growth Parameters

In terms of Plant Height. Maximum 5, 8.81, 19.82, 26.41 and 28.59 cm at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub> (NPK 12.5 ml/plant) followed by T<sub>3</sub> (NPK 10.41 ml/plant) with 4.71, 8.16, 16.76, 26.41 and 25.4 cm, minimum 4.10, 7.44, 7.76, 7.99 and 8.81cm, was recorded in Control. Significantly maximum plant height was recorded in T<sub>4</sub> at 10, 20, 30, 40 and 50 DAP, which might be due to higher levels of N, P & K, which found suitable for spinach in hydroponic system helpful in cell elongation of leaves use to development of cell and rapid cell division and cell elongation in meristematic region of plant due to production of plant growth substance and this may be due to abundant supply of plant nutrients and water which led in the growth of Spinach. This finding correlates the findings of **Park et al., 1999** in Basil, **Oztekin et al. (2008)** in Spinach, **Adams, P. (2002)** in Hydroponic production of vegetables.

In terms of Plant spread. Maximum 4.63, 5.62, 21.1, 41.26 and 45.96 cm at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub> (NPK 12.5 ml/plant) followed by T<sub>3</sub> (NPK 10.41 ml/plant) with 4.10, 5.02, 18.93, 38.76 and 42.79 cm, minimum 3.46, 4.34, 6.04, 8.04, 10 cm, was recorded in Control. Significantly maximum plant spread was recorded in T<sub>4</sub> at 10, 20, 30, 40 and 50 DAP, which might be due to higher levels of N, P & K, which found suitable for Spinach in hydroponic system resulted enhanced photosynthetic and other metabolic activities which lead to increase in various plant metabolites responsible for cell division and elongation. This finding correlates the findings of **Park et al., 1999** in Basil, **Oztekin et al. (2008)** in Spinach, **Adams, P. (2002)** in Hydroponic production of vegetables.

In terms of Number of leaves/plant. Maximum 4.60, 6.83, 13.22, 21.55 and 23.33 leaves/plant at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub> (NPK 10.41 ml/plant) with 4.11, 6.02, 11.22, 19.56 and 21.44 leaves/plant and minimum 3.62, 4.44, 5.84, 6,4 and 7.44 leaves/plant was recorded in Control. Significantly maximum Number of leaves

per plant was recorded in T<sub>5</sub> at 10, 20, 30, 40 and 50 DAP, which might be due to higher levels of N, P & K, which found suitable for Spinach in hydroponic system resulted enhanced photosynthetic and other metabolic activities which lead to increase in various plant metabolites responsible for cell division and elongation. This finding correlates the findings of **Park et al., 1999** in Basil, **Oztekin et al. (2008)** in Spinach, **Adams, P. (2002)** in Hydroponic production of vegetables.

In terms of Leaf length. Maximum 5.98, 10.04, 21.39, 27.20 and 29.07 cm , at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub>(NPK 10.41 ml/plant) with 5.93, 8.86, 17.50, 24.82 and 25.50 cm, and minimum 4.63, 7.03, 7.87, 8.62 and 9.61 cm , was recorded in Control. Significantly maximum Leaf length was recorded in T<sub>4</sub> at 10, 20, 30, 40 and 50 DAP, which might be due to higher levels of N, P & K, which found suitable for Spinach in hydroponic system and enhanced photosynthetic and other metabolic activities which lead to increase in various plant metabolites responsible for cell division and elongation. This finding correlates the findings of **Park et al., 1999** in Basil, **Oztekin et al. (2008)** in Spinach, **Adams, P. (2002)** in Hydroponic production of vegetables.

In terms of Leaf width. Maximum 1.13, 3.38, 7.28, 8.49 and 8.96 cm, at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub> (NPK 10.41 ml/plant) with 1.01, 2.91, 5.49, 8.08 and 8.37 cm, and minimum 0.87, 1.19, 1.58, 1.83 and 2.08 cm , was recorded in Control. Significantly maximum Leaf width was recorded in T<sub>4</sub> at 10, 20, 30, 40 and 50 DAP, which might be due to higher levels of N, P & K, which found suitable for Spinach in hydroponic system and enhanced photosynthetic and other metabolic activities which lead to increase in various plant metabolites responsible for cell division and elongation. This finding correlates the findings of **Park et al., 1999** in Basil, **Oztekin et al. (2008)** in Spinach, **Adams, P. (2002)** in Hydroponic production of vegetables.

In terms of Root length. Maximum 7.58, 17.98, 32.29, 48.86 and 70.55 cm, at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub>(NPK 10.41ml/plant) with 7.19, 16.20, 27.67, 44.92 and 61.33 cm, and minimum 6.29, 13.02, 24.33, 38.86 and 54.67 cm, was recorded in Control. Significantly maximum Root length was recorded in T<sub>4</sub> at 10, 20, 30, 40 and 50 DAP, this may be due to large amount of nutrients directly available to the plants roots, which led in the higher water uptake of plants and ultimately growth of roots of Spinach. This finding correlates the findings of **Maneejantra et al. (2016)** in Spinach and **Sapkota et al. (2019)** in Lettuce.

In terms of Water used. Maximum 5.91, 12.21, 13.39, 17.97, and 18.46 litre, at 10, 20, 30, 40 and 50 DAP was recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub> (NPK 10.41 ml/plant) with 5.17, 11.85, 12.17, 16.43 and 17.56 litre, and minimum 4.07, 6.60, 7.92, 12.88 and 15.65 litre was recorded in Control.. This might be due higher dose of N, P and K in NFT hydroponic system in which plant growth, yield, and water consumption vary depending on the concentration of nutrient solution as well as the temperature during the growing season. When the temperature and nutrient concentration increased plant growth and water consumption also increased. This finding correlates the findings of **Zhang et al. (2016)** in water use efficiency of Tomato and **Oztekin et al., (2018)** in Spinach.

### B. Yield Parameters

In Average weight of plants. Maximum Average weight (62.48 g) were recorded in T<sub>4</sub> (NPK 12.5ml) after application of NPK, followed by T<sub>3</sub> (NPK 10.41ml) with (54.63 g) whereas minimum weight (3.89 g) was recorded in T<sub>0</sub> (Control).This might be due to N, P, and K application might be attributed to enhanced photosynthesis, accumulation of carbohydrates and favourable effect on vegetative growth which also increase the weight and size of plants. This finding correlates the findings of **Maneejantra et al. (2016)** in Spinach and **Akanbi et al. (2007)** in pepper.

In Yield/Structure (kg). Maximum Yield/Structure (1.50 kg) were recorded in T<sub>4</sub> (NPK12.5 ml) after application of NPK, followed by T<sub>3</sub> (NPK 10.41 ml) with (1.31 kg) whereas minimum yield (0.09 kg) was recorded in T<sub>0</sub> (Control).This might be due to the plant growth and final yield depends on the continued supply of food material and water. Since N, P and K help in the absorption of water and carbohydrates metabolism, its deficiency may cause poor growth and yield of plants. This finding correlates the findings of **Oztekin et al. (2018)** in Spinach and **Paradosi et al. (1987)** in Tomato.

### C. Quality parameters

The data on Dry matter percent of Spinach of different treatment combinations of NPK in NFT hydroponics system of Spinach was recorded and are presented in table 4.12. Critical analysis of data displayed in table clearly marked out the obvious difference among the treatments with respect to Dry matter content. Significant difference was observed due to different concentration of NPK for Dry matter content. Maximum dry matter percent (35.53 %) was recorded in T<sub>4</sub> (NPK 12.5ml) after application of NPK, followed by T<sub>3</sub> (NPK 10.41ml) with (33.3 %) whereas minimum dry matter content (24.90 %) was recorded in T<sub>0</sub> (Control).

The data on **organoleptic** evaluation parameter of spinach of different treatment combinations of NPK in NFT hydroponics system of spinach was recorded and are presented in table 4.13. Critical analysis of data displayed in table clearly marked out the obvious difference among the treatments with respect to **organoleptic** parameters.

In terms of Score for colour and appearance, maximum score was recorded in treatment T<sub>4</sub> (NPK 12.5 ml/Plant) with 8.60 followed by treatment T<sub>3</sub> (NPK 10.41 ml/Plant) with 7.12, and minimum score was observed in treatment T<sub>0</sub> (Control) with 2.14. In terms of Score

**Comment [u4]:** Replace with organoleptic

**Comment [u5]:** Replace with organoleptic

for Texture, maximum score was recorded in treatment T<sub>4</sub> (NPK 12.5 ml/Plant) with 8.55 followed by treatment T<sub>3</sub> (NPK 10.41 ml/Plant) with 7.32, and minimum score was observed in treatment T<sub>0</sub> (Control) with 2.93. In terms of Score for Flavour and taste maximum score was recorded in treatment T<sub>4</sub>(NPK 12.5 ml/Plant) with 8.71 followed by treatment T<sub>3</sub> (NPK 10.41 ml/Plant) with 8.04, and minimum score was observed in treatment T<sub>0</sub>(Control) with 3.13. In terms of Score for Overall acceptability maximum score was recorded in treatment T<sub>4</sub> (NPK 12.5 ml/Plant) with 8.89 followed by treatment T<sub>3</sub>(NPK 10.41 ml/Plant) with 7.77, and minimum score was observed in treatment T<sub>0</sub> (Control) with 2.34.

#### D. Economics

In terms of economics, maximum cost of cultivation (Rs. 1425.00) was observed in T<sub>7</sub> (NPK 18.75 ml/plant) followed by T<sub>6</sub> (NPK 16.66 ml/plant) (Rs. 1324.70), minimum cost of cultivation (Rs. 525.00) was observed in Control. In Gross return (Rs. 1871.30) is recorded in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub>(NPK 10.41 ml/plant) (Rs.1636.28), minimum Gross return (Rs. 116.45) was recorded in T<sub>0</sub> (NPK 0 ml/plant). In Net return (Rs.746.30) is found in T<sub>4</sub>(NPK 12.5 ml/plant) followed by T<sub>3</sub>(10.41 ml/plant) (Rs.611.60), minimum Net return (Rs. 87.00) was found in T<sub>7</sub> (NPK 18.75 ml/plant). In Benefit: Cost ratio (1.66) is observed in T<sub>4</sub> (12.5ml/plant) followed by T<sub>1</sub> (NPK 6.25 ml/plant) (1.63), minimum B:C ratio (0.22) was observed in T<sub>0</sub> (NPK 0 ml/plant).

Comment [u6]: Replace with was

**Table 1.** Performance of Spinach for growth parameters in NFT hydroponic system.

Treatment	Plant Height (cm)					Plant Spread (cm)					Number of Leaves/plant				
	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP
T <sub>0</sub>	4.10	7.44	7.76	7.99	8.81	3.46	4.34	6.04	8.04	10.00	3.62	4.44	5.89	6.44	7.44
T <sub>1</sub>	4.28	7.78	13.84	17.60	22.00	3.94	5.02	17.54	27.46	30.22	4.08	5.94	10.33	16.89	18.78
T <sub>2</sub>	4.41	7.59	13.10	14.96	16.61	3.64	4.41	13.60	21.28	23.14	4.00	5.68	8.78	13.44	15.00
T <sub>3</sub>	4.71	8.16	16.76	22.06	25.44	4.10	5.02	18.93	38.76	42.79	4.11	6.02	11.22	19.56	21.44
T <sub>4</sub>	5.00	8.81	19.82	26.41	28.59	4.63	5.62	21.11	41.26	45.96	4.60	6.83	13.22	21.55	23.33
T <sub>5</sub>	4.68	7.91	12.94	16.68	20.28	4.07	4.67	14.47	25.08	30.89	3.89	5.78	8.67	14.22	15.89
T <sub>6</sub>	4.74	7.49	15.47	17.51	21.03	4.03	4.80	14.09	22.80	25.08	3.89	5.33	8.11	13.22	14.11
T <sub>7</sub>	4.68	7.50	14.46	17.21	19.90	3.93	4.42	13.48	19.02	21.18	3.71	5.00	7.56	11.89	14.00
F-Test	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
C.D. at 5%	0.41	0.63	1.01	1.13	0.83	0.39	0.41	0.92	1.34	1.32	0.39	0.46	0.69	0.79	0.67
SE(d)	0.19	0.30	0.47	0.53	0.39	0.18	0.19	0.43	0.63	0.62	0.18	0.22	0.33	0.37	0.31
C.V.	5.09	4.61	4.05	3.69	2.35	5.66	4.84	3.55	3.02	2.65	5.63	4.72	4.31	3.09	2.37

**Table 2.** Performance of Spinach for growth parameters in NFT hydroponic system.

Treatment	Leaf Length (cm)					Leaf width (cm)					Root length (cm)				
	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP
T <sub>0</sub>	4.63	7.03	7.87	8.62	9.61	0.87	1.19	1.58	1.83	2.08	6.29	13.02	24.33	38.86	54.67
T <sub>1</sub>	5.42	8.74	16.44	18.20	21.68	0.98	2.82	5.17	7.88	8.06	4.93	12.37	20.50	31.10	45.14
T <sub>2</sub>	5.12	8.26	13.86	17.67	21.00	0.94	2.80	4.92	7.60	7.92	7.41	17.17	29.67	46.77	64.26
T <sub>3</sub>	5.43	8.86	17.50	24.82	25.50	1.01	2.91	5.49	8.08	8.37	7.19	16.20	27.67	44.92	61.33
T <sub>4</sub>	5.98	10.04	21.39	27.20	29.07	1.13	3.38	7.28	8.49	8.96	8.12	20.37	35.29	52.37	76.86
T <sub>5</sub>	5.10	8.61	14.53	18.39	21.73	0.94	2.83	4.99	7.23	7.71	7.58	17.98	32.29	48.86	70.55
T <sub>6</sub>	5.48	8.18	15.26	17.88	20.97	0.99	2.82	4.76	5.27	5.66	7.09	14.67	27.50	42.67	59.33
T <sub>7</sub>	4.97	8.09	13.51	15.64	17.47	0.88	2.03	4.34	4.69	4.84	6.95	14.17	26.00	41.44	57.33
F-Test	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
C.D. at 5%	0.53	0.61	0.96	1.06	1.04	0.094	0.241	0.398	0.306	0.241	0.54	1.12	1.80	1.93	2.31
SE(d)	0.25	0.29	0.45	0.50	0.49	0.044	0.113	0.186	0.143	0.113	0.26	0.53	0.84	0.90	1.08
C.V.	5.72	4.11	3.65	3.28	2.86	5.561	5.311	4.736	2.748	2.060	4.49	4.08	3.68	2.54	2.16

**Table 3.** Performance of Spinach for water used, Weight and yield parameters in NFT hydroponic system.

Treatment Symbol	Water used (liter)					Avg. weight of plant (g)	Yield/structure (kg)
	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP		
T <sub>0</sub>	4.07	6.6	7.92	12.88	15.65	3.89	0.09
T <sub>1</sub>	4.76	11.3	11.48	16.18	17.03	49.88	1.20
T <sub>2</sub>	4.69	9.18	10.4	14.52	16.23	41.86	1.00
T <sub>3</sub>	5.17	11.85	12.17	16.43	17.56	54.11	1.30
T <sub>4</sub>	5.91	12.21	13.34	17.97	18.46	57.74	1.39
T <sub>5</sub>	4.48	10.66	11	15.36	16.96	52.39	1.26
T <sub>6</sub>	4.18	9.55	10.87	14.89	16.78	47.27	1.14
T <sub>7</sub>	4.33	9.92	10.21	14.23	16.6	42.72	1.02
F-Test	S	S	S	S	S	S	S
C.D. at 5%	0.085	0.344	0.225	0.053	0.028	0.85	0.02
SE(d)	0.182	0.734	0.225	0.113	0.061	2.39	2.48
C.V.	2.215	4.141	2.526	0.421	0.206	1.82	0.05

**Table 4.** Performance of Spinach for Dry matter, Organoleptic evaluation (Accessed with Hedonic scale) and B: C ratio.

Treatment	Dry matter	Organoleptic Evaluation				B:C Ratio
		Score for colour and appearance	Score for texture	Score for Flavour and Taste	Score for Over all acceptability	
T <sub>0</sub>	24.90	2.14	2.93	3.13	2.34	0.22
T <sub>1</sub>	32.04	7.34	7.17	7.34	7.54	1.63
T <sub>2</sub>	30.94	6.92	6.22	6.45	6.29	1.44
T <sub>3</sub>	33.30	7.12	7.32	8.04	7.77	1.60
T <sub>4</sub>	35.53	8.60	8.55	8.71	8.89	1.66
T <sub>5</sub>	30.31	7.03	6.75	6.97	6.69	1.31
T <sub>6</sub>	29.44	6.89	6.76	6.56	6.31	1.18
T <sub>7</sub>	28.35	5.97	6.44	6.36	6.42	0.94
F-Test	S	S	S	S	S	
C.D. at 5%	0.50	0.14	0.12	0.15	0.13	
SE(d)	2.00	2.56	2.25	2.78	2.43	
C.V.	1.07	0.29	0.26	0.33	0.28	

## CONCLUSION

From the present investigation it is concluded that treatment T<sub>4</sub> found best in terms of Growth and yield parameters of Spinach in NFT hydroponic system followed by T<sub>3</sub> and T<sub>1</sub>. Minimum growth and yield was recorded in T<sub>0</sub>. Maximum gross return and net return was recorded in T<sub>4</sub> and maximum cost benefit ratio 1.66 was observed in treatment T<sub>4</sub> minimum gross and net return recorded in T<sub>0</sub> and T<sub>7</sub> and cost benefit ratio in T<sub>0</sub>. In hydroponics, the flow of nutrient and nutrient uptake is a vital criterion for good growth and yield. Nutrient deficit leads to stunted growth of plants and overly absorption of nutrients can lead to antagonistic consequence on plant growth. Therefore, an ideal nutrient help the plants in good growth and yield. As seen from the experiment, it is observed that the treatment T<sub>4</sub> is ideal nutrient for Spinach good growth and yield.

## REFERENCES

- A.Pardosi, F.tognoni and G.Bertero. (1987).** The influence of nutrient solution concentration on growth, mineral uptake and yield of Tomato plants grown in N.F.T. *Advances in Horticultural Science.*, 1 (1987):55-60.
- Adams, P. 2002.** Nutritional control in hydroponics.,p. 211–261. In: D. Savvas and H. Passam(eds.) *Hydroponic Production of Vegetables*
- Akanbi, W. B., Togu, A. O., Akinfasoye, J. O. and Tairu, F. M. (2007).** Physico-chemical properties of eggplant (*Solanum melongena* L.) fruit in response to nitrogen fertilizer and fruit size. *Agr. J.*, 2(1): 140-148.
- Anonymous (2016).** Statistical data on Horticultural Crops in India, 8-13.
- Carmassi, G., Incrocci, L., Maggini, R., Malorgio, F., Tognoni, F., and Pardossi, A., (2007).** An aggregated model for water requirements of greenhouse tomato grown in closed rockwool culture with saline water. *Agricultural water management*, 88(1):73-82.
- Della Costa, L. and Gianquinto, G., (2002).** Water Stress and Watertable Depth Influence Yield, Water Use Efficiency, and Nitrogen Recovery in Bell Pepper: Lysimeter Studies. *Aust. J. Agric. Res.*, 53: 201-210.
- Domingues, D.S., Takahashi, H.W., Camara, C.A.P. and Nixdorf, S.L. (2012).** Automated system developed to control pH and concentration of nutrient solution evaluated in hydroponic lettuce production. *Computers and Electronics in Agriculture*, 84,53-61.
- Hoyos, M., & Rodriguez-Delfin, A., (2007).** Effect of two nutrient solution formulae in the nutritional diagnosis and yield of tomato grown in soilless culture system. *Actahorticulturae*, 747:471-475.
- Hussain, A., Iqbal, K., Aziem, S., Mahato, P., and Negi. A. K., (2014).** A Review on The Science of Growing Crops Without Soil (Soilless Culture) A Novel Alternative For Growing Crops. *International Journal of Agriculture and Crop Sciences*, 7(11):833- 842.
- Mohammed Dahiru Toungos (2017)** The effects of different levels of inorganic fertilizer (NPK 15:15:15,) on growth and yield of sweet pepper (*capsicum annum*) in mubi, Nigeria *International Journal of Development Research* ,07(06): 13120-13124.

Comment [u7]: Insert comma

Comment [u8]: Insert s

Comment [u9]: Replace with was

Comment [u10]: Replace with was

- Maneejantra N, Tsukagoshi S, Lu N, Supaibulwatana K, Takagaki M (2016)** A Quantitative Analysis of Nutrient Requirements for Hydroponic Spinach (*Spinacia oleracea* L.) Production Under Artificial Light in a Plant Factory. *J Fertil Pestic* 7: 170.
- N.sharma,S.Acharya,K.Kumar,N. Singh and O.P Chaurasia (2018)** Hydroponics as an advanced technique for vegetable production:An Overview.*Journal of Soil and water conservation* 17(4):364-371.
- Oztekin, G. B., Uludag, T., and Tuzel, Y. (2018)**Growing spinach (*Spinacia oleracea* l.) in a floating system with different concentrations of nutrient solution., *Applied ecology and environmental research.*,**16**(3): 3333-3350.
- Park Kuen Woo., Park Kwang Woo and Suh EunJoo (1999).** Effect of different concentrations of nutrient solutions on the growth, yield and quality of basil. *Acta Hort*483.
- Sezen, S. M., Celikel, G., Yazar, A., Mendi, Y. Y., Sahinler, S., Tekin, S. and Gencil, B., (2006).** Effects of drip irrigation management and different soilless culture on yield and quality of tomato grown in a plastic house. *Pakistan Journal of Biological Sciences*, **9**: 766-771.
- Sharma, N., Acharya, S., Kumar, K., Singh, N. and Chaurasia, O.P. (2019)** Hydroponics as an Advanced Technique for Vegetable Production: An Overview. *Journal of Soil and Water Conservation*, 17, 364-371.
- Sundar, S., Sanjib, S., and Liu, Zhiming (2019)** Effects of Nutrient Composition and Lettuce Cultivar on Crop Production in Hydroponic Culture., *Horticulturae.*,**5**(72) 10.3390/horticulturae5040072.
- Sharma, N., Acharya, S., Kumar, K., Singh, N. and Chaurasia, O.P. (2019)** Hydroponics as an Advanced Technique for Vegetable Production: An Overview. *Journal of Soil and Water Conservation*, 17, 364-371.
- Pengfei Zhang, Masateru Senge and Yanyan Dai (2016).** Effects of salinity stress on growth, Yield, fruit quality and water Use efficiency of tomato under Hydroponics system (2016) *Reviews in Agricultural Sciences*, 4:46-55