

Original Research Article

Study of Jamun (*Syzygium cumini* Skeels.) Genotypes Based on Shelf life and Interspecific Crossability

ABSTRACT

An experiment was conducted during 2019 to study the jamun genotypes based on shelf life of fruit. Shelf life studies were conducted on eight jamun genotypes and results indicated that, the maximum shelf life of the Jamun fruit was ~~5~~ five days under normal room temperature after harvest at ripened stage. Genotypes Dhoopdal, Selection-45 and Savadatti had better shelf life compare to other genotypes. In crossing, out of 100 crossed flowers, only ~~5~~ five crossed flowers were set and produced fully mature fruits.

Key-~~Key~~-words: Jamun genotypes, Shelf life, Inter-specific crossing

Introduction

Jamun, botanically called *Syzygium cumini* Skeels, belongs to the family ~~myrtaceae~~ *Myrtaceae* (Sharma *et al.*, 2012). Jamun is a multipurpose tree of both food and medicinal values. Fruits are used for table purpose and for the preparation of different variety of products (~~Juice~~ juice, ~~Squash~~ squash, ~~Jam~~ jam *etc.*). Tree ~~grows is~~ tall and ~~is~~ evergreen, long lived and also has got ornamental value; ~~Inflorescences~~ inflorescences are borne in leaf axils of branch lets (panicles). Flowers are bisexual and light yellow in colour. Jamun is a cross pollinated tree. Fruits are oblong and round in shape, deep purple or bluish, juicy with sweet pulp with single seed. The most common type is Rajamun, which ~~had~~ has large sized, oblong shaped fruits of deep purple colour with juicy sweet pulp. All parts of the tree such as fruits, leaves, seeds, and bark are used in Indian medicine system like Ayurveda, Homeopathy, Sidda and Unani (~~AYUSH~~) *etc.* Different parts of the jamun were also reported for antioxidant, anti-inflammatory, anti-microbial, anti-diarrheal, gastro protective and anti-ulcerogenic properties. Before the discovery of insulin, in the treatment of diabetes *S. cumini* was used either alone or in combination with other hypoglycemic plants even in Europe (~~Helmstadter, et al., 2008~~).

The fruits of jamun can be harvested after fully mature and ripen stage, but the harvesting is difficult due to large tree size. The fruits get damaged easily while harvesting which also weakens the storability. Jamun (*S. cumini*) has very low shelf life, whereas rose apple (*S. jambos*) has good keeping quality. It also produces fruits twice a year whereas jamun produces fruits only once in a year. ~~Keeping In view of~~ these positive characters, ~~the in view~~ interspecific crossing is done in between these *Syzygium* species. ~~So~~ Therefore, crop improvement work in jamun is through selection of seedling trees based on high shelf life and other promising traits ~~is needed~~. The study ~~hwas~~ ~~the continued with the help of~~ following objectives: to study fruit quality based on shelf life in jamun collections and to assess crossability among the *Syzygium jambos* with *S. cumini*.

Comment [A1]: The objective of interspecific crossing remains to be mentioned.
Quote the species studied and place of study.

Comment [A2]: Authors could choose keywords that do not appear in the title, to increase the search possibilities for the manuscript.

Formatted: Font: Italic

Formatted: Font: Italic

Comment [A3]: Put meaning in the first quote of the acronym.

Formatted: Font: Italic

Comment [A4]: Is this the entire paragraph reference?

Comment [A5]: Include the species author in the first quote.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Not Italic

Material and Methods

The experiment entitled “Study of Jamun (*Syzygium cumini* Skeels.) Genotypes Based on Shelf life and Interspecific Crossability” was conducted at the experimental plot of division of fruit crops at ICAR-Indian Institute of Horticultural Research, Bengaluru, during June 2019. The accessions for the study were selected from jamun germplasm block of ICAR-IIHR. The study on number of days, the fruit remained good marketable and fit for consumption under room temperature was recorded. For this, eight genotypes were used for shelf life analysis of fruits. To study crossability percentage of *Syzygium* sp. especially rose apple with jamun pollen, 100 selected fully mature flowers were tagged, bagged and hand pollinated with jamun pollen in morning between 10am –and 12pm. Emasculated and bagged stigma of opening flowers were dusted with pollen of jamun genotype that is Huruli chikkanahalli. Selectively 100 Rose apple flowers were pollinated a day before flower opening and bagged. After 20-25 days fruit set was recorded and percentage of fruit set was calculated.

Comment [A6]: There is no need to repeat the manuscript title. It might just be: “The present study was conducted...”

Comment [A7]:

Comment [A8]: Quote all genotypes used in the study.

Formatted: Font: Not Bold

Formatted: Font: Not Italic

Results and Discussion

The results of the study pertaining to the Study of Jamun (*Syzygium cumini* Skeels.) Genotypes on Shelf life and Interspecific Crossing were presented in Table 1. Although jamun fruits have got high neutraucitical properties, the fruits have maximum shelf life of only 5 days at room temperature. Fruits stored under normal temperature condition weight of the fruits were decreasing day to day (Table 1). The genotypes Dhoopdal sel-20 (10.01 g), Selection-45 (10.08 g) and Savadatti (11.00 g) has maintained highest fruit weight in fifth day of storage, compare to other genotypes. It may be due to loss of water in storage. A similar result was reported by Dalvadi *et al.*, (2017) in jamun significantly reduced physiological loss in weight under room storage. TSS of the stored fruits was increasing from first day to fifth day in almost all genotypes, especially in Dhoopdal sel-20 (22.2 °B) and AJG-85 (25.2 °B) were showing drastic increase (Table 1). It may be due to respiration process of the fruit under storage, utilisation stored carbohydrates conversion into sugars. Acidity of the stored fruits was decreasing towards ripening up to three days and increasing of acidity was observed towards senescence stage in Selection-45 (1.34%) and Savadatti (1.00%).

Comment [A9]: Same as previous comment.

Comment [A10]: Suggestion: “Under normal temperature conditions, fruits stored weight of the fruits were decreasing day to day.”

Comment [A11]: Put meaning in the first quote of the acronym.

The data of crossability was presented in Table 2. Hybridization/Crossing is done in between *Syzygium jambos* × *Syzygium cumini* to see the fruit set percentage in different species of same genus of *Syzygium*. The Rose apple was flowering twice in a year (October and March) and Jamun flowering in Feb/March, hence the crossing is done in during March and by using Emasculatation and Hand pollination method. Crossing is was done in 100 emasculated flowers, fruit set will occur occurred 2-3 weeks after pollination a. And the fruits will matured within 50-60 days. Out of 100 crossed flowers, 5-five flowers set to matured ripened (golden yellow coloured) fruits (Table 2). Inter-specific crosses between *Syzygium jambos* and *S. cumini* was attempted in 100 emasculated flowers, of fruit set observed 2-3 weeks after pollination. And the fruits matured within 50-60 days. Out of 100 crossed flowers, only 5 flowers matured ripened (golden yellow coloured) fruits. This may be due to low cross compatibility between rose apple and jamun. And pre-fruit set flower drop was more in *Syzygium* sp. Kataoka *et al.*, (2003) in kiwi and Noguchi *et al.*, (2002) in strawberry.

Comment [A12]: These data should be in the Material and Methods section.

Comment [A13]: Futures are not used in results, as these are data that have already been obtained.

Comment [A14]: It would be interesting to add a photo of the fruit generated from the hybridization.

Comment [A15]: Did these fruits have seeds? Are the seeds viable?

Comment [A16]: Repeated excerpt.

Comment [A17]: Check the year of publication, as the references are 2009.

Comment [A18]: No reference to strawberry?

Formatted: Font: Italic

Conclusion

Jamun fruits treat as new emerging fruit crop for future due to having high neutraucitical properties. Although, the fruits have maximum shelf life of only 5-five days at room temperature. So, this study on shelf life and interspecific crossing of jamun collections will helps, to further development of genotypes Dhoopdal, Selection-45 and Savadatti to improve shelf life in jamun.

Comment [A19]: No need to put result data on conclusion. You might write for example: "...the fruits have a short shelf life at room temperature."

References

- Dalvadi, A. C., Dodiya, K. D., Bhad, D. M., Chaudhari, T. and Trambadiya, R., 2017, Effect of chemicals on shelf life of (*Syzygium cuminii* Skeels) CV. *Local. Agric. Sci. Digest.* **38** (1): 68-70.
- Helmstadter, A., 2008, *Syzygium Cumini* (L.) Skeels (Myrtaceae), 2008, a Against diabetes 125 years of research. *Pharm Sci.* **63**(2):91-101.
- Kataoka, K. Beppu, K. Fukuda, T. and Mabuchi, S., 2003, Evaluation of characteristics of Actinidia interspecific hybrid 'Kosui'. *Acta. Hort.* 610.
- Noguchi, Y., Morishita, M. and Muro, T. 2009, The possibility of using decaploid interspecific hybrids (*Fragaria* × *ananassa* × *F. nilgerrensis*) as a parent for a new strawberry. *Acta. Hort.* 842.
- Sharma, S., Mehta, B. K., Mehta, D., Nagar, H. and Mishra, A., 2012, A review on pharmacological activity of *Syzygium cumini* extracts using different solvent and their effective doses. *Int. Res. J. Pharm.* **3**(1): 54-58.

Comment [A20]: Check the year of publication, as the text is 2002.

Table 1. Shelf life study of jamun genotypes

Genotypes	Weight (g)					TSS (°B)					Acidity (%)				
	Days					Days					Days				
	1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
Dhoopdal sel-20	11.80	11.30	10.70	10.30	10.0	21.00	22.50	24.00	25.70	22.20	1.30	1.10	0.80	0.90	1.20
Selection-45	12.10	11.45	11.22	10.75	10.08	21.00	21.50	22.00	24.80	21.60	1.10	0.90	1.30	1.42	1.34
Selection-58	7.80	7.52	7.37	7.04	6.80	17.00	17.70	19.20	19.80	18.30	0.80	0.60	0.65	0.77	1.00
Savadatti	12.20	11.84	11.73	11.40	11.00	18.00	19.00	19.50	18.00	18.00	1.40	1.20	1.28	0.98	1.30
Kaithnal	7.20	6.94	6.44	6.23	5.92	20.00	23.50	26.00	24.50	22.00	1.50	1.30	1.50	1.20	1.40
AJG-85	8.20	7.90	7.51	7.11	6.79	21.00	21.50	23.70	25.90	25.20	1.70	1.62	1.50	1.60	1.90
Dharwad-3a	1.30	1.13	0.89	0.59	0.55	28.00	28.40	29.00	28.30	26.00	2.40	2.30	2.12	2.32	**
Madhya Pradesh -5	0.90	0.60	0.56	0.43	0.40	16.00	17.50	19.00	18.00	--	3.10	3.20	3.41	3.20	**

Comment [A21]: Put the meaning of the acronym in the legend.

(--: not suitable for TSS estimation)
 (**: not suitable for Acidity estimation).

Table 2. Crossability percentage in between *Syzygium* species

<i>S. jambos</i>	No. of flowers crossed	Fruit set (%)	Control (%)
X <i>S. cumini</i>	100	5	3

Formatted: Font: Italic

Comment [A22]: This control was not explained in the results.

Formatted: Font: Italic

Formatted: Font: Italic

UNDER PEER REVIEW

