

## Original Research Article

### Effect of land configuration and moisture regimes on productivity of hybrid maize (*Zea mays* L.) in summer season.

#### Abstract

A field experiment was conducted on “Effect of land configuration and moisture regimes on productivity of hybrid maize (*Zea mays* L.) in summer season.” during 2018-19 at Agronomy Research Farm, ANDUA, and T, Narendra Nagar (Kumarganj), Ayodhya (U.P.). The experiment comprised of 12 treatment combinations and laid out in split plot design, replicated four times. The Experiment consisted of three land configuration viz. flat bed planting (P<sub>1</sub>), ridge planting (P<sub>2</sub>) and paired row planting on raised bed (P<sub>3</sub>) were kept in main plots and four moisture regimes viz. 0.5 IW/CPE ratio, 0.7 IW/CPE ratios, 0.9 IW/CPE ratios and 1.1 IW/CPE ratios were allotted in sub plots. The experimental results revealed that growth parameters, such as plant height, leaf area index, dry matter accumulation, days to 50 percent ear emergence and days to 50 percent maturity and yield, its attributes such as number of cobs plant<sup>-1</sup>, number grains row<sup>-1</sup>, number of grains cob<sup>-1</sup>, cob length (cm), cob girth (cm), grains weight cob<sup>-1</sup>(g), cob weight (g), shelling percentage (%), grain yield (q ha<sup>-1</sup>), stover yield (q ha<sup>-1</sup>), and biological yield (q ha<sup>-1</sup>) were observed significantly maximum except test weight and harvest index under paired row planting on raised bed followed by ridge planting and flat bed planting. In case of moisture regimes, the same trends were observed more with the irrigation at 1.1 IW/CPE ratios as compared to other moisture regimes.

**Keywords-** Maize, Land configuration, Moisture regimes, Yield, Growth. Leaf area index

#### Introduction

Maize (*Zea mays* L.) is one of the most important cereal crops in the world. It belongs to family *Poaceae*, broadly used in industries besides serving as human food and animal feed. Maize is called “queen of cereal” because it has got very high yield potential and wide adaptability under various agro climatic conditions than any other cereal crops (Singh, 2013). The world leading producer of maize in descending order are: United States of America (357.267 million tons), China (215.00 million tons), Brazil (95.00 million tons), Argentina (33.80 million tons), Ukraine (30.94 million tons), India (25.00 million tons), Mexico (23.27million tons), and Indonesia (19.00 million tons) (Anonymous, 2017). In India, maize is the third most important food grain crop after wheat and rice. In India area under maize cultivation in both the seasons (*kharif*, 7.7 mha) and *rabi*, 1.6 mha) with a production of 19.5 and 7.6 mt respectively. (Anonymous, 2017)

India produces about 2% of the world’s maize production out of which Karnataka is leading producer of maize about 16% of total maize production followed by Telangana and Bihar which together contribute about 20% of maize production. Maharastra, Madhya Pradesh, Tamilnadu, Andhra Pradesh, Rajasthan and Uttar Pradesh are other maize producing states of India. About 71%

Deleted: &

Deleted: with

Deleted: ratio

Deleted: ratio

Deleted: ratio

Comment [H1]: Too long sentence. Recast.

Deleted: ratio

Deleted: :

Deleted: crop

Formatted: Font: Italic

Deleted: maize

Deleted: ity

Comment [H2]: This reference is not listed. Update.

Deleted: ,

Deleted: &

Deleted: india

of maize in India is produced in the *kharif* season. It is generally grown in Karnataka, Madhya Pradesh, Tamilnadu, Maharashtra, Telangana, Uttar Pradesh and Rajasthan etc. Bihar, Andhra Pradesh and Tamilnadu are major states which produces maize in *rabi* season out of which Tamilnadu is a leading state to share 40% of production.

Deleted: &

Deleted: &

Deleted: produced

Land configuration plays a major role in minimizing soil erosion and improving water and nutrient use efficiency of field crops. Easy and uniform germination as well as growth and development of plant are provided by manipulation of sowing methods Chiroma *et al.* (2008) and also increases availability of nutrients to crops. The superiority of ridges and furrow system could be described to proper drainage of excess water coupled with adequate aeration at the time of irrigation or heavy rainfall. Parihar *et al.* 2009 reported that ridges and furrow method of sowing improved grain as well as stover yield of maize over the flat bed method of sowing. Water stress can affect growth, development and physiological processes of maize plants, which reduce biomass yield Payero *et al.* (2008). Water requirement of crop mostly depends on Evapo-transpiration, which mainly depends on climate. The amount of water lost by evapo-transpiration is estimated from climatological data and when ET reaches a particular level irrigation is scheduled. The amount of irrigation is given to be equal to ET or fraction of ET. IW/CPE approach is well known amount of irrigation water applied when cumulative pan evaporation reaches predetermine level. ET by a full crop cover in closely associated with the evaporation from an open pan, Dasten (1967). Parihar *et al.*, (2009) suggested relatively more practical meteorological approach of the ratio between a fixed amount of irrigation water and CPE as a basis of irrigation scheduling. It was found that irrigation maize crop at an IW/CPE ratio of 1.0 was significantly superior to 0.6 IW/CPE but at par with 0.8 IW/CPE regarding plant dry weight (g) at harvest stage, number of kernels/cob, kernel weight (g)/cob, kernel yield (kg ha<sup>-1</sup>) and stover yield (kg ha<sup>-1</sup>) Malla Reddy *et al.*, (2012).

Deleted: 2009,

Deleted: . Evapotranspiration

Deleted: ly

Deleted: in

Comment [H3]: Never start a sentence with an abbreviation or figure.

Comment [H4]: Never start a sentence with an abbreviation or figure.

Comment [H5]: Clear objective of this research is not stated.

Comment [H6]: This reference is not listed. Update.

## Material and methods

The experiment was conducted at Agronomy Research Farm of Acharya Narendra Deva University of Agriculture and Technology. Kumarganj, Ayodhya (U.P.). The farm situated on Ayodhya, Raibareily road at the distance of 42 km from Ayodhya district head quarter. Geographically, this experimental site falls under semi-arid sub tropical climate of Indo-genetic plains having alluvial plains (IGP) having alluvial calcareous soil and is located at 26° 47" N latititude and 82° 12" E longitude and an altitude 113 meters above mean sea level. Soil of experimental field was slightly alkaline in reaction (8.0 pH), low in organic carbon 0.32 per cent and low in available nitrogen (180 kg ha<sup>-1</sup>), medium in Phosphorus (15. 25 kg ha<sup>-1</sup>) and potassium

Deleted: research

Deleted: &

Deleted: Latitude

(270.0 kg ha<sup>-1</sup>). Kanchan (K-25) variety of maize was used for sowing at 25 kg ha<sup>-1</sup>. The crop was sown at 07 April 2019 on a distance of 60 cm in row and 20 cm apart from plant.

Deleted: @

Deleted: with keeping the

Deleted: was

The experiment comprised of twelve treatment combinations with three land configuration *viz.* flat bed planting (P<sub>1</sub>), ridge planting (P<sub>2</sub>) and paired row planting on raised bed (P<sub>3</sub>) were kept in main plots and four moisture regimes *viz.* 0.5 IW/CPE ratio, 0.7 IW/CPE ratios, 0.9 IW/CPE ratio and 1.1 IW/CPE ratio were allotted in sub plots. It was conducted in split plot design with four replications. Five plant were selected randomly from each plot for the observation of growth character and yield and its attributes *viz.*, as plant height (cm), Leaf area index, dry matter accumulation (gm<sup>-2</sup>), number of cobs plant<sup>-1</sup>, number of rows cob<sup>-1</sup>, number of grains cob<sup>-1</sup>, number of grains row<sup>-1</sup>, grain weight cob<sup>-1</sup>, girth of cob, length of cob, weight of cob (g) and shelling percentage, grain yield (qha<sup>-1</sup>), stover yield (qha<sup>-1</sup>), and biological yield (qha<sup>-1</sup>), at 30, 60 and at harvest stage.

Deleted: ratio

### Shelling percentage

Five sample cobs taken from each plot for grain weight per cob were also used for this purpose. Those five sample cobs were weighed and grains were separated. Shelling percent was taken used the following formula.

$$\text{Shelling percent} = \frac{\text{Grain weight per cob}}{\text{Cob weight}} \times 100$$

### Harvest Index (%)

Harvest index is defined as the ratio of economic yield to total biological yield (Donald, 1962) and expressed in percentage. The harvest index for maize was worked out as indicated below.

Comment [H7]: This reference is not listed. Update.

$$\text{Harvest index \%} = \frac{\text{Economic yield (kg/ha)}}{\text{Biological yield (kg/ha)}} \times 100$$

## Result and Discussion

Deleted: discussion

### Growth parameter

The growth parameters *viz.*, plant height, leaf area index, and dry matter accumulation were significantly affected by land configuration at successive growth stages. Crop sown on paired row planting on raised bed (L<sub>3</sub>) produced higher plant height, leaf area index and dry matter accumulation at all the stages, which was significant superior over ridge planting (L<sub>2</sub>) and flat bed planting (L<sub>1</sub>) presented in Table 1. This might be due to increases soil aeration, nutrient and

moisture availability to the plant as compared to other land configuration. Similar result was reported by Kumar (2008) and Joshi *et al.* (2018).

Deleted: reported

In case of moisture regimes, increasing moisture supply increased the plant height significantly at all growth stages, except at 30 DAS. Maximum plant height at 60 DAS and at harvest stage was observed under 0.9 IW/CPE ratio which was at par with 1.1 IW/CPE ratio and significantly superior over 0.7 IW/CPE ratio, and 0.5 IW/CPE ratio. Moisture regimes significantly influenced leaf area index at 60 DAS, while it was found non-significant at 30 DAS and at harvest stage. The maximum leaf area index was recorded under irrigation at 0.9 IW/CPE ratios as compared to the 1.1 IW/CPE ratios, 0.7 IW/CPE ratios and 0.5 IW/CPE ratios. This might be due to rapid growth caused by maintenance of adequate and continuous water supply to the crop which maintained good establishment of the roots and various metabolic processes which performed higher nutrient mobility and uptake contributed to rapid cell division and cell elongation which resulted in higher plant height and higher leaf area under the treatment. Minimum plant height and leaf area index was obtained under 0.5 IW/CPE ratio at all the stages due to poor root growth caused by moisture deficit. The results are in close proximity to those of Hussaini *et al.* (2002), Meena *et al.* (2017) and Kumar *et al.* (2011).

Deleted: with

Deleted: the

Deleted: and

Deleted: were influenced

Deleted: on

Deleted: ratio

Deleted: ratio

Deleted: ratio

Deleted: ratio

Comment [H8]: Too long sentence. Recast.

Comment [H9]: This reference is not listed. Update.

Deleted: due to

Deleted: were

Deleted: ratio

Deleted: for

Deleted: resulted

Deleted: ratio

Dry matter accumulation ( $\text{gm}^{-2}$ ) was influenced significantly by different moisture regimes at 60 DAS and at harvest stages except 30 DAS as presented in (Table 1) the maximum dry matter accumulation was recorded with the irrigation at 0.9 IW/CPE ratios. This might be due to increased plant height, leaf area index, girth of stem and increase in uptake of nutrient viz., N, P, and K through adequate moisture supply. All these contributed to cell turgidity and opened leaves which increased the photosynthetic activity of crops resulting in higher dry matter accumulation. Initially, it increased slowly and thereafter rapidly till the harvest stage due to variable conditions. The lowest dry matter accumulation was obtained under 0.5 IW/CPE ratios. This might be due to lowest number of irrigations were given which results reduced in plant height, leaf area and nutrient uptake when lead to reduced the photosynthetic activity which ultimately reflected in lowest dry matter accumulation. Similar finding were reported by Singh (2001), Hussaini *et al.* (2002) and Kumar *et al.* (2011).

### Yield attributes

Yield attributes likes number of cobs  $\text{plant}^{-1}$ , number of rows  $\text{cob}^{-1}$ , number of grains  $\text{cob}^{-1}$ , number of grains  $\text{row}^{-1}$ , grain weight  $\text{cob}^{-1}$ , girth of cob, length of cob, weight of cob (g) and shelling percentage were significantly influenced due to land configuration that as presented in

Deleted: is

Table 2 except test weight. Significantly higher values were recorded in paired row planting on raised bed as compared to ridge planting and flat bed planting. Flat bed planting restricted the crop growth and induced early ear emergence. This might be due to the low aeration and low nutrient to crop, which resulted in reduced value of all attribute, also reported by several others Tanveer *et al.* (2014) and Singh *et al.* (2015).

Deleted: was

Comment [H10]: This reference is not listed. Update.

In case of moisture all the attributes *viz.*, number of cobs plant<sup>-1</sup>, number of rows cob<sup>-1</sup>, number of grains cob<sup>-1</sup>, number of grains row<sup>-1</sup>, grain weight cob<sup>-1</sup>, girth of cob, length of cob, weight of cob and shelling percentage were significantly influenced by different moisture regimes. Highest values of all the yield attributes were recorded under 0.9 IW/CPE ratios which were at par with 1.1 IW/CPE ratios. This might be due to favorable vegetative growth and development because it received adequate water during entire period of plant growth. As per need which increased the all growth parameters and increased photosynthetic activity of leaves, beside translocation of photosynthesis from source to sink also increased under wettest regime through higher uptake of potassium which led to better yield attributes. Lowest values of yield attributes was found in 0.5 IW/CPE ratio because plant were unable to extract more water and nutrient under moisture deficit condition which resulted in poor growth and yield attributes. Similar finding was also reported by Talu *et al.* (1998) and Reddy *et al.* (2012).

Deleted: due to

Deleted: ratio

Deleted: was

Deleted: ratio

Deleted: photosynthates

## Yield

Grain and stover yield of maize were affected significantly by land configuration as presented in (Table 3). Crop sown on paired row planting raised bed recorded significantly higher grain yield as compared to ridge planting and flat bed planting. A similar trend was also recorded in straw yield of maize. This might be due to poor growth and translocation of photosynthates from source to sink. All the growth and yield attributes which determined the grain and stover yield of maize crop, were adversely influenced when the sowing on flat bed planting. Significant reduction in grain and stover yield of maize in flat bed method has also been reported by several Aggarwal and Goswami (2002), Mishra *et al.* (2009) and Joshi *et al.* (2018).

Deleted: data

Deleted: flat

Deleted: .(

Grain and straw yield was influenced significantly by different moisture regimes have been presented in Table 3. The maximum grain yield was obtained under irrigation at 0.9 IW/CPE ratios. This might be due to optimum moisture availability as per demand of crop, which contributed to better growth parameters and yield attributes. Productivity of crop collectively determined by vigour of vegetative growth and yield attributes. Irrigation at 0.5 IW/CPE ratio recorded lowest grain yield due to poor moisture supply during crop growth period. Poor moisture supply during

Deleted: ratio

critical stages reduced the yield attributes and resulted in poor grain and stover yield. Similar finding were reported by Talu *et al.* (1998), Singh (2001), Hussaini *et al.* (2002) and Manna *et al.* (2018).

The biological yield was influenced significant by land configuration. The maximum biological yield was obtained under treatment paired row planting on raise bed (L<sub>3</sub>). This might also be increased due to better soil environment. Similar result reported by Mehta *et al.* (2010) and Meena *et al.* (2015).

The biological yield was influenced significantly by different moisture regimes. The maximum biological yield was obtained under irrigation at 0.9 IW/CPE [ratios](#). This might be due to adequate supply of water, which contributed to increasing in dry matter accumulation. Better vegetative growth coupled with high yield attributes resulted into higher biological yield. The minimum biological yield recorded under 0.5 IW/CPE [ratios](#) this might be due to both poor growth and yield attributes. Similar finding were reported by Meena *et al.* (2015).

**Comment [H11]:** This reference is not listed. Update.

**Deleted:** ratio

**Deleted:** ratio





**Table -3. Grain & Stover yield, biological yield and harvest index as influenced by land configuration and moisture regimes of maize in summer season.**

Formatted: Width: 8.27", Height: 11.69"

Treatments	Grain yield (q ha <sup>-1</sup> )	Biological yield (q ha <sup>-1</sup> )	Stover yield (q ha <sup>-1</sup> )	Harvest index (%)
<b>A- Land configuration</b>				
Flat bed planting	33.50	85.27	51.77	39.20
Ridge planting	37.00	93.94	56.94	39.30
Paired row planting	43.20	108.57	65.37	39.70
Sem±	0.785	1.900	1.206	0.859
CD at 5%	1.839	4.450	2.824	N.S
<b>B- Moisture regimes</b>				
6 cm irrigation at 0.5 IW/CPE ratio	28.10	72.57	44.47	38.70
6 cm irrigation at 0.7 IW/CPE ratio	33.40	85.16	51.76	39.20
6 cm irrigation at 0.9 IW/CPE ratio	45.70	114.76	69.06	39.80
6 cm irrigation at 1.1 IW/ CPE ratio	44.40	111.22	66.82	39.90
Sem±	0.651	1.645	1.006	0.626
CD at 5%	1.367	3.455	2.112	N.S

Formatted Table

**Reference:**

Aggarwal, P. and Goswami, B. (2002). Bed planting system for increasing water use efficiency of wheat (*Triticum aestivum* L.) grown in inceptisol. *Indian J Agric Sci* **73**: 422-425.

Anonymous (2017). www. Agricoop. Nic.in & http://eands.dacnet.nic.in.

Chiroma, A.M.; Alhassan, A.B. and Khan, B. (2008).Yield and water use efficiency of millet as affected by land configuration treatments. *J Sustainable Agric.* **32**(2):321-333.

Dastane, N.G. (1967). A practical manual for water use research. *Poona navabharat prakashans.* **13**(16):46-50.

Hussaini, M.A.; Ogunlela, V.B.; Ramalan, A.A. and Falaki, A. M. (2002). Growth and development of maize (*Zea mays*) in response to different level of N.P and irrigation. *Crop Research.* **22** (2) : 139-149.

Comment [H12]: Spellings not correct. Check inside the text.

- Joshi, J.R.; Patel, V.M.; Barad, H.L.; Macwan, S.M.; and Ehsas, J. (2018). Effect of land configuration and fertilizer management practices on growth, yield and yield attributes and economics of Summer maize (*Zea mays* L.) under South Gujrat condition. *Int. J. Curr. Microbiol. App. Sci.* **7**(1):1148-1155.
- Kumar, A. (2008). Growth, yield and water use efficiency of different maize (*Zea mays* L.) based cropping systems under varying planting methods and irrigation levels. *Indian J Agric Sci.* **78**: 254-57.
- Kumar, H.K.; Ramachandrappa, B.K.; Nanjappa, H.V. and Mudalagiriappa. (2011). Effect of phenophase based irrigation schedules on growth, yield and quality of baby corn (*Zea mays* L.) *Agricultural Sciences.* **2**(3): 267-272.
- Manna, T.; Saha, G.; Saha, A.; Dutta, D. and Nanda, Nanda. (2018). Contribution of micro climate towards yield attributing factors and yield of summer baby corn (*Zea mays* L.) under different irrigations and mulches. *Int.J.Curr.Microbiol. App. Sci.* **7**(7): 1542-1552.
- Meena, R.K.; Tiwari, R.C. and Meena, V.D.; Charpota, J.L.; and Meena, A.K. (2017). Effect of Irrigation Management and Plant Population on the Performance of Summer baby corn (*Zea mays* L.). *Int J. Curr. Microbiol. App. Sci.* **6**(7): 2274- 2282.
- Meena, R.L.; Idnani, L.K.; Kumar, A.; Khanna, M.; Shukla, L. and Choudhary, R.L. (2015). Water economization in rabi maize (*Zea mays* L.) to enhance productivity through land configuration and irrigation scheduling in the Indo-Gangetic plains of India. *Journal of Soil and Water Conservation.* **14**(1): 49-55.
- Mishra, P.K.; Reddy, V.; Ch and U. Kumar, S. (2009). An evaluation of furrows for managing soil and water loss from shallow alfisols under simulated rainfall. *Soil Use and Management*, **24**: 171- 180.
- Parihar, C.M.; Rana, K.S and Parihar, M.D. (2009). Crop productivity, quality and nutrient uptake of maize (*Zea mays* L.) cropping system as influenced by configuration and direct and residual effect of nutrient management. *Indian J. agric. Sci.* **79**(11):927-930.
- Payero, J.O.; Tarkalson, D.D.; Irmak,S.; Davison, D. and Petersen, J.L. (2008). Effect of irrigation amounts applied with subsurface drip irrigation on corn evapotranspiration, yield, water use efficiency, and dry matter production in a semi arid climate. *Agric. Water Manage.***95**: 895-908.
- Reddy, M. M.; Padmaja, B. and Reddy, V.V.D. (2012). Response of Maize (*Zea mays* L.) to irrigation scheduling and nitrogen doses under no till condition in rice fallows. *J. Res. ANGRAU.*, **40**(1):6-12.
- Singh, J. and Vashist, K.K. (2015). Effect of planting methods, mulching and irrigation regimes on maize productivity. *Agric Res J.* **52**: 23-27
- Singh, S.S.D. (2001). Effect of irrigation regimes and nitrogen levels on growth yield and quality of baby corn (*Zea mays* L.). *Madras Agricultural Journal.* **88** (7-9): 367-370.

**Comment [H13]:** This reference was not cited. Delete.

Tanveer, M.; Ehsanullah, Anjum, S.A.; Zahid, H.; Rehman, A. and Sajjad. (2014). Growth and development of maize (*Zea mays* L.) in response to different planting methods. *J Agric Res* **52**: 511-522.

Tulu, L.; Ramachandrappa, B.K. and Nanjappa, H.V. (1998). Response of maize (*Zea mays* L.) to moisture stress at different growth stages in alfisols during summer. *Mysore Journal of Agricultural Sciences*. **32**: 201-207.