

## Original Research Article

# Impact of Fall Army Worm (FAW) (*Spodoptera frugiperda*) on the Yield of Maize under different cropping patterns of Western Agro Climatic Zone of Tamil Nadu

### ABSTRACT

The invasive pest, Fall Army ~~Worm-worm~~ (FAW), is a serious ~~one~~ because of its widespread outbreak ~~on-in~~ maize growing regions. Since its occurrence across the world, the main adoption technique to manage the pest is the spraying of chemical pesticides. This leads to negative environmental impacts. Thus, the present investigation was undergone during the Kharif and Rabi seasons of 2019 and 2020 to study the influence of different dates of sowing and intercropping of legumes on the level of infestation of fall armyworm on maize crop. Three dates of sowing were adopted in each season and intercrops like cowpea, soybean, sunn\_hemp, and tephrosia were included beside sole maize with and without insecticide spray. The results showed that the maize planted during July and October ~~month~~ has reported less FAW infestation ~~of FAW~~ and provided a higher yield than the rest of the planting taken. The cowpea and sunn\_hemp intercropped maize has performed well in terms of yield and the leaf damage was also found to be lower apart from the insecticide applied maize crop.

*Keywords: Fall armyworm, Maize yield, Intercropping, Dates of sowing*

### 1. INTRODUCTION

Until 2016, the pest, Fall Army ~~W~~worm (FAW), *Spodoptera frugiperda* (J.E. Smith) (*Lepidoptera: Noctuidae*) has been a major ~~issue-pest of maize and rice~~ in America. In 2016, the FAW was reported in the African continent for the first time outside to its natural habitat and the reason for the invasion is mysterious. But it is believed that the haplotype that emerged from Florida and the Caribbean existed in Africa (Huesing *et al.*, 2018). Moreover, the other reason for FAW to easily invade the continent is because of its prolific reproducing ability and skill-capacity to migrate long distances. The FAW larva caused severe destruction to its host by consuming foliage while the FAW adult ~~takes-took~~ care of dispersion. The early instar larva causes skeletonized leaves and windowed whorls, while the matured larva feeds on maize cob resulting in a fall in yield and quality (Capinera *et al.*, 2017). FAW can cause a yield reduction of about 34 to 38% (Nelly *et al.*, 2021).

The pest FAW has caused severerious devastation to the maize crop in India since its outbreak in 2018. After its first report at the College of Agriculture, Shivamogga, Karnataka, it has rapidly spread across the nation (reference). The chief maize cultivating states contribute 79% of the country's maize production and they are Uttar Pradesh, Bihar, Madhya Pradesh, Maharashtra, Rajasthan, Gujarat, Karnataka, Andhra Pradesh, and Tamil Nadu. Most of the maize produced in the country is used for poultry and livestock feed and ~~for~~ industrial purposes. The input of Tamil Nadu to the nation's maize yield is almost 7.25% from the major growing districts like Perambalur, Ariyalur, Cuddalore, Dindigul, and Tirupur. Of the 3.55 lakh acres of maize cultivated, the pest FAW damaged 2.2 lakh acres during 2018 in Tamil Nadu.

Comment [WU1]: In which crop

The foremost strategy adopted across the globe to manage FAW is the application of chemical insecticides. Though this technique reduces the pest damage caused and decreases the spread of the pest, its constant use may lead to the development of insecticide resistance in FAW (reference). Further, this method is not useful for FAW management as the excreta of this pest are heavy, which blocks the pesticide and may not reach the whorl where the larvae are feeding (Day *et al.*, 2017). Hence, the alternative technique for the management of FAW should be is to implement an environment-friendly, and moreover cost effective. It was reported that integrated approach which includes push-pull technology, use of botanical extracts, intercropping, natural enemies and parasitoids, utilizing locally available materials, etc effectively reduces the incidence of FAW (Khatiri *et al.* 2020; Keerthi *et al.* 2020). Keeping this in view, the present investigation has been carried out to find a management option that deliberates ecological values.

## 2. MATERIAL AND METHODS

The trial was carried out in Eastern Block Farm at Tamil Nadu Agricultural University, Coimbatore from 2019 to 2020. The investigational field is situated at 11°N and 76°57'E longitude and an altitude of 426.7 m above MSL in the Western Agro-Climatic Zone (ACZ) of Tamil Nadu. The soil of the experimental site was sandy clay loam in texture. The climatic condition of the site was hot semi-arid under the Koppen climate classification. The wet season of the site lasts from June to November.

Maize was sown at three dates of sowing in four seasons viz., Kharif, 2019 (D<sub>1</sub> - June 15th, D<sub>2</sub> - July 15th, and D<sub>3</sub> - August 15th), Rabi, 2019 (D<sub>1</sub> - September 15th, D<sub>2</sub> - October 15th and D<sub>3</sub> - November 15th), Late Rabi, 2020-2021 (D<sub>1</sub> - December 15th, D<sub>2</sub> - January 15th and D<sub>3</sub> - February 15th) and **Summer, 2021** (D<sub>1</sub> - March 15th, D<sub>2</sub> - April 15th and D<sub>3</sub> - May 15th) and were taken in the main plot. Along with sole maize without insecticide spray (S<sub>1</sub>) and sole maize with recommended insecticidal spray (S<sub>2</sub>), the intercrops were taken in subplot viz., maize + cowpea (S<sub>3</sub>), maize + soybean (S<sub>4</sub>), maize + sunn hemp (S<sub>5</sub>), and maize + tephrosia (S<sub>6</sub>). The split-plot design was adopted and replicated thrice.

The fields were ploughed with a cultivator and rotovator to get a fine, clod-free tilth. After forming ridges and furrows, the area was delineated into plots, and treatments were allocated. Neem cake was applied two weeks before the sowing and fertilizers were applied as per recommendation given by TNAU (135: 62.5: 50 kg NPK/ha). Maize was planted at a spacing of 60 x 25 cm while the intercrops were sown in between the maize rows. The after cultivation for maize was done as per the recommended practices while the intercrops were not undergone any such operations like maize. The intercrops were brown manured at 50 days after sowing by spraying 2, 4-D herbicide. The chemical pesticides for treatment S<sub>2</sub> (Sole maize with recommended insecticide spray) were sprayed at 20 and 40 DAS using Azadirachtin and Emamectin Benzoate, respectively.

The damage caused by FAW on maize foliage was noted down before tasseling stage by using Davis *et al.* (1992) scoring scale and the yield of maize was recorded for different dates of sowing.

## 3. RESULTS AND DISCUSSION

The influence of fall armyworm leaf damage on yield of maize at different dates of sowing and intercropping is presented in Figure 1 to 6.

Among the various treatments adopted, the sole maize with recommended insecticide spray (S<sub>2</sub>) has produced a better yield than the others irrespective of the sowing dates of sowing. This is due to the quick action and higher toxicity of insecticides that are sprayed at the whorls of maize where the larvae reside and this was in accordance with the findings of Pitre (1986). The next greatest yield was found in maize intercropped with cowpea (S<sub>3</sub>) followed by sunn\_hemp (S<sub>5</sub>) in all three plantings of Kharif and Rabi season. This is because the cowpea and sunnhemp intercrops possesses the ability to minimize can minimize the masses of FAW, which also delays its time of infestation in succeeding maize

Formatted: Space After: 12 pt

Comment [WU2]: Please check the date, whether it is Late rabi 2019-20; and summer 2020.

Comment [WU3]: Summer 2021 results missing in result and discussion

Formatted: Highlight


Formatted: Highlight

Comment [WU4]: Otherwise there wont be a continuity in the work.


Comment [WU5]: Wheteher it was 0-4 scale

growing season. This was similar to the results of Meagher *et al.* (2004), who reported that the FAW larvae took more time to develop in cowpea and sunn\_hemp crops and also the pupae weight was found to be 20 to 25% lower than the other treatments implemented.


Of the three sowings taken in Kharif and Rabi season, the maize planting done during July (7200 kg/ha) and October (7015 kg/ha) month has produced better yield compared to the other two plantings in both seasons, which is because of the less leaf damage of FAW in July (1.47) and October (1.18) month. The minimal leaf damage during these two plantings might be due to the copious amount of rainfall received during its initial stages of vegetative growth stages. This effective rain might have provided better growing condition for the maize crop during its initial stages and have reduced the spread and damage of FAW. This result was similar to the outcomes of Harrison *et al.* (2019), which stated that the first effective rains provide better growing conditions for maize, making use of more heat units at the beginning of the cropping season.




**Fig 1. Influence of FAW leaf damage on the yield of maize at first date of sowing (June 15<sup>th</sup>) during Kharif, 2019**




**Fig 2. Influence of FAW leaf damage on the yield of maize at second date of sowing (July 15<sup>th</sup>) during Kharif, 2019**




**Fig 3. Influence of FAW leaf damage on the yield of maize at third date of sowing (August 15<sup>th</sup>) during Kharif, 2019**



**Fig 4. Influence of FAW leaf damage on the yield of maize at first date of sowing (September 15<sup>th</sup>) during Rabi, 2019**



**Fig 5. Influence of FAW leaf damage on the yield of maize at second date of sowing (October 15<sup>th</sup>) during Rabi, 2019**



**Fig 6. Influence of FAW leaf damage on the yield of maize at third date of sowing (November 15<sup>th</sup>) during Rabi, 2019**

#### 4. CONCLUSION

It ~~is was evident obvious~~ from the above results that the maize crop which received better growing condition can withstand the damage caused by FAW ~~pest~~ and those with cowpea and sunn\_hemp intercropping performed well next to the chemical treatment.

#### REFERENCES

1. Capinera JL. Fall Armyworm, *Spodoptera frugiperda* (J. E. Smith) (Insecta: Lepidoptera: Noctuidae). Publication EENY098. Florida (FL): UF/IFAS Extension Service, University of Florida. 2017.
2. Davis FM, Williams WP. Visual rating scales for screening whorl-stage corn for resistance to fall armyworm. Mississippi Agricultural & Forestry Experiment Station. Technical Bulletin 186, Mississippi State University, USA. 1992.
3. Day R, Abrahams P, Bateman M, Beale T, Clotey V, Cock M, Gomez J. Fall armyworm: impacts and implications for Africa. *Outlooks on Pest Management*. 2017;28(5):196-201.
4. Harrison RD, Thierfelder C, Baudron F, Chinwada P, Midega C, Schaffner U, Van den Berg J. Agro-ecological options for fall armyworm (*Spodoptera frugiperda* JE Smith) management: Providing low-cost, smallholder friendly solutions to an invasive pest. *J. Environ. Manag.*, 2019;243:318-330.
5. Huesing JE, Prasanna BM, McGrath D, Chinwada P, Jepson P, Capinera JL. Integrated pest management of fall armyworm in Africa: an introduction. In: Prasanna B.M, Huesing JE, Eddy R, Peschke VM (Eds), *Fall armyworm in Africa. A Guide for integrated pest management*. CIMMYT, Mexico, CDMX. 2018.

[Keerthi MC, Sravika A, Mahesha HS, Gupta A, Bhargavi HA, Ahmed S. Performance of the native predatory bug, \*Eocanthecona furcellata\* \(Wolff\)\(Hemiptera: Pentatomidae\), on the fall](#)

[armyworm, \*Spodoptera frugiperda\* \(JE. Smith\)\(Lepidoptera: Noctuidae\), and its limitation under field condition. Egyptian Journal of Biological Pest Control. 2020 Dec;30:1-4.](#)

[Khatri S, Pakuwal P, Khanal S. Integrated pest management of fall armyworm infestations in maize fields in Nepal: A review. Archives of Agriculture and Environmental Science. 2020 Dec 25;5\(4\):583-91.](#)

6. Meagher RL, Nagoshi RN, Stuhl C, Mitchell ER. Larval development of fall armyworm (Lepidoptera: Noctuidae) on different cover crop plants. Florida Entomologist. 2004;87 (4).

7. Nelly N, Hamid H, Lina EC, Yunisman. The use of several maize varieties by farmers and the infestation of *Spodoptera frugiperda* (Noctuidae: Lepidoptera). IOP Conf. Series: Earth and Environmental Science 2021;662.

8. Pitre H. Chemical control of the fall armyworm (Lepidoptera: Noctuidae). Florida Entomol. 1986;69(3):570–578.

UNDER PEER REVIEW