

Original Research Article

INFLUENCE OF SALINITY WITH DIFFERENT $\text{Cl}^-:\text{SO}_4^{2-}$ RATIOS ON WHEAT (*Triticum aestivum* L.) GROWTH

Abstract

A significant threat to sustainable agriculture is the continuous increase in salt-affected areas. Salt stress causes land degradation, ecological imbalance, environmental pollution and reduces crop production. It changes different physical and chemical processes in the plant body, such as seed germination and different nutrients and water uptake. Wheat is the main cereal crop and primary diet for one-third of humans. It can tolerate the salinity effect to some extent, but its production reduces significantly at a higher level of salinity. Salt-affected soils have more salts that are easily solubilized in water and exchangeable sodium than the average soil. These salts are ionized and produce different cations (Mg^{2+} , Ca^{2+} , and Na^+) and anions (Cl^- , CO_3^{2-} , SO_4^{2-} , and HCO_3^-). Sulfate salinity is also toxic for plant growth as chloride salinity. This research aims to assess the effect of salinity with various $\text{Cl}^-:\text{SO}_4^{2-}$ ratios on wheat growth. A pot trial was conducted on wheat by creating salinity with different levels of chloride and sulphate (4:1, 3:1, 2:1, 1:1, 1:2, 1:3, and 1:4). The experimental design was a complete randomized design (CRD). Different physicochemical parameters of the soil before and after crop harvesting was determined. Different growth parameters will also be determined. Data were analyzed using statistical software. Based on these results, it was determined that both types of salinity reduced the wheat growth significantly, but chloride type salinity has a more injurious effect than sulphate salinity. It was concluded that chloride ion has more toxic effects than sulphate ion for wheat growth and development.

Keywords: Wheat, salinity, chloride, sulfate, abiotic stress

1. Introduction

These days, water shortages and environmental degradation are significant issues. Salinization and urbanization pose a danger to the long-term viability of agriculture (Gould *et al.*, 2021). Salinity may lead to a decrease in agricultural productivity, an imbalance in the ecosystem, and changes in physiochemical soil characteristics (Machado and Serralheiro,

2017). The annual increase in salinized area is estimated at 10 % each year by factors like high surface evaporation, low precipitation, breakdown of the rocks, poor cultural activities, and irrigation with water having more salts (Kumar and Sharma, 2020). A saline-sodic soil has electrical conductivity (EC_e) of saturated extract more than 4 dS m^{-1} and 15 exchangeable sodium percentage (ESP) at 25°C . The production of most of the crops is reduced at this EC_e and ESP (Munns, 2005). Consequently, the Plant undergoes morphological, biochemical, and physiological modifications (Wang *et al.*, 2021). Seed germination is inhibited by soluble salts, which restrict seeds from taking water from the soil due to osmotic action (Tavares *et al.*, 2021).

Salinity and sodicity result in a decrease in crop production, ecological imbalance, and change in soil physiochemical properties, which ultimately causes low income, soil erosion, and environmental pollution (Gould *et al.*, 2021).

More than half of the world's irrigated land and over a quarter of the world's cultivated land are suffering from salt stress. About 14 % of Pakistan's irrigated land has been salinized, and salinity is responsible for 64 % of the country's output losses. There are now only 23 million hectares (23 million acres) of land used for farming (Sanower and Sultan, 2019; Syed *et al.*, 2021). Almost 3 percent of irrigated land in Punjab, 18 percent in Sindh, and 2 percent in the NWFP suffer from severe surface salinity (Syed *et al.*, 2021). Pakistan's geographic size is estimated to be 79.61 million square miles, however only roughly 4.5 million square miles is salt affected (Tavares *et al.*, 2021).

Chlorine and sulphur both are essential nutrients for plants but due to their higher concentration and mobility in irrigated water caused toxicity as a micronutrient it is crucial to regulate the osmotic pressure, photosynthesis, and enzyme action (Reich *et al.*, 2015; Geilfus, 2018). Sulphur is essential to produce several biomolecules, including methionine, cysteine, chlorophyll, proteins, and the oil content of seeds, among others. Understanding how sulphur interacts with other macro components is critical to maximise crop yields (Aziz *et al.*, 2019; Geilfus, 2018). There is an adverse association between salinity and germination rate of seeds. When there is an increase in salinity and sodicity, the germination rate can be reduced and delayed (Yan *et al.*, 2019). Salinity causes to reduce the osmotic potential, protein and nucleic acid metabolism, uses of stored food by seeds (Helal and Mengel, 1979, Machado and Serralheiro, 2017, Yan *et al.*, 2019). It alters the enzymatic activates during germination process (Tavares *et al.*, 2021). It not only delays the germination but also affects the

vegetative and propagative growth of the Plant by creating osmotic and oxidative stresses as well as deficiency and toxicity in them (Benedetti *et al.*, 2020).

Salt-affected soils have high concentration of cations (Ca^{2+} , Mg^{2+} and Na^+) and anions (Cl^- , CO_3^{2-} , SO_4^{2-} , and HCO_3^-) (Geilfus, 2018). Chlorine and sulphur toxicity is widespread in salt-affected soil because it is the most common anion in salt-affected soil (Reich *et al.*, 2015; Tavares *et al.*, 2021). High concentration of Cl^- ion in chloroplast or cytosol tissue limits the growth of many glycophytic plants (Geilfus, 2018). It also retards the growth of root and tuber in pea, potato and avocado (Duan *et al.*, 2013). Chlorine (Cl^-) has an antagonistic impact on numerous nutrients, such as nitrate absorption by Brassica rapa, which is reduced due to increased soil Cl^- concentrations (Reich *et al.*, 2017). When combined with phosphate and sulphate, it has an antagonistic impact on plant cells, decreasing their absorption and transport (Massa *et al.*, 2009). But excessive concentration of sulphur causes toxicity in plants. In many plant species, it is more harmful than chloride salinity (Ranault *et al.*, 2001).

Plants' responses to salt-affected soils have been studied by incorporating a combination of all salts that may be present (Hu *et al.*, 2018; Jixiang, 2012; Yan *et al.*, 2019; Zhang and Zhao, 2011, Zhao *et al.*, 2014). According to published research, Cl^- reduces germination in three Mediterranean salt marsh species more severely than SO_4^{2-} (Zhao *et al.*, 2014). In salt-stressed areas of Argentina, the germination of an inbred species of salt-affected plants is more severely inhibited by SO_4^{2-} than by Cl^- (Sosa *et al.*, 2005). When it comes to whether ion is more hazardous to plants, there is a lot of confusion. This research will examine the wheat's ability to withstand Cl^- and SO_4^{2-} salinity.

The objective of this study was:

- To assess the influence of salinity with different ratios of Cl^- and SO_4^{2-} on wheat growth.
- To determine which anion (Cl^- or SO_4^{2-}) is more toxic for wheat growth

2. Methodology

2.1. Growth condition and Experimental Layout

A pot trial was conducted in the wire house of Institute of Soil and Environmental Sciences (ISES), University of Agriculture, Faisalabad (UAF) was conducted during the year 2018-19. The test crop was wheat (Galaxy 2013).

Soil for the experiment taken from farm area of ISES. For the soil preparation, it was first air-dried in a shade to remove the water contents. Then it was grounded and to remove the material other than soil (stones, plant debris) it was sieved through 2 mm sieve. This prepared soil was brought in the lab for different analyses like saturation percentage, texture, carbonate, bicarbonate, chloride, sulphate, calcium, magnesium, pH, EC and SAR before the start of the experiment (Richards, 1954). According to the treatments, salinity ($EC = 10 \text{ dS m}^{-1}$) was created artificially using Na_2SO_4 and NaCl salts.

2.2. Development of Salinity and Crop Growth

Soil was weighed, and a quadratic equation was used to calculate the amount of salts per the above treatments. The amount of water for dissolving the salts was equal to SP of the soil. Then soil was saturated by applying different salt solutions according to their treatments. After one month, incubated soil was dried and filled in pots (4.5 kg capacity).

At field capacity, 5 seeds of wheat were sown in each pot (24 pots). Recommended amounts of N, P and K for wheat were applied using urea, DAP and SOP as their sources. After germination, 3 plants were kept in each pot. Canal water was used to irrigate the plants as per crop requirement. At maturity, crop growth parameters were recorded. From each pot, soil samples were obtained with stainless steel tube and were analysed for chemical properties. The plant material was analyzed for N, P, K, Cl^- and SO_4^{2-} .

Comment [A1]: Any specific concentration of ions present in water. Please specify if yes

2.3. Plant and Grain Samples Preparation

The plant shoots and grains samples were also taken to analyse different parameters (N, P, K, Cl^- and SO_4^{2-}).

2.3.1. Grinding

For the preparation of plant and grain samples, grinding of these samples were done with the help of electric grinder (small in case of grains grinding) to get the material in powder form. After grinding, the digestion of plant material was done.

2.3.2. Di-acid Digestion

Di-acid digestion was used to prepare the Plant or grain sample for all other analysis except N. For di-acid digestion process, the mixture of two acids (hydrochloric acid (HNO_3) and perchloric acid (HClO_4) at the ratio of 2:1 respectively) were used. 0.5 g plant sample or grain sample was taken in flask. 5 ml of di-acid was used for each sample and keep overnight.

After 24 hours, these flasks were put on the hot plates at 300°C until the color of inside material became white or white fumes started to come out. After that, put out these flasks from the hot plate and cooled them. After cooling, the material inside these flasks was filtrated with Whatman filter paper number 40. 50 ml volume was made in Volumatic flask with DI water of each sample. Now the sample was prepared for further analysis.

2.4. Kjeldahl Method for N Determination in Plant

Kjeldahl method for N determination has following two steps;

2.4.1. Digestion

To prepare the sample for N analysis, digestion of that sample was done. 0.5 g grinded plant or grain samples were kept in digestion flasks or tubes. 5 g digestion mixture ($K_2SO_4 + CuSO_4 \cdot 5H_2O + Se$ at the ratio of 100: 10: 1 respectively) and 15 to 20 ml concentrated H_2SO_4 (0.1 N) were added in these flasks. Put these flasks in digestion block and heat them at 400 °C for 3 to 4 hours until the transparent or light yellowish colored material was observed in these flasks. After that, put out these flasks from the digestion block and cooled them. After cooling, the material inside these flasks was filtrated with Whatman filter paper number 40. 50 ml volume was made in Volumatic flask with DI water of each sample. Now the sample was prepared for further analysis.

2.4.2. Distillation

After the digestion of the sample, the next step for the N determination was the distillation. For distillation process, 10 ml of digested filtrate was kept in beaker. Added 10 ml of 4 % boric acid (H_3BO_3) in this beaker. 2 to 3 drops of indicator (methyl red) was added. The color of the sample became purple. After that, placed the beaker on the distillation unit for 3 min during which the color was changed from purple to light yellow. After 3 min, take the breaker out from the distillation unit and titrate it against the 0.1 N H_2SO_4 until the sample's color again becomes purple. The volume of 0.1 N H_2SO_4 used for titration was noticed, which was used to calculate N concentration in this sample.

The N concentration (%) was determined with following formula;

$$N (\%) = (V_0 - B) * 0.1 * V_1 * 14.01 * 100 / Wt. * V_2 \times 1000$$

Where;

V_0 = vol. of H_2SO_4 used for titration of sample

B = vol. of H_2SO_4 used for titration of blank

V_1 = total vol. of digest

W_t = weight of sample

14.01 = N atomic weight

2.5. Determination of P in Plants

For P determination, 10 ml of the filtrate sample was put into a flask (100 ml) after di-acid digestion. 10 ml of reagent ammonium-vanadomolybdate (ammonium heptamolybdate $[(NH_4)_6 Mo_7 O_{24} \cdot 4H_2O]$ + ammonium vanadate (NH_4VO_3) + nitric acid (HNO_3) was added in this flask and made the volume with DI water. The standards were formed by potassium dihydrogen phosphate salt, having the concentration of P (0.5, 1, 1.5, 2 and 2.5 ppm). 10 ml of each standard was also taken in flask (100 ml) and used reagent (10 ml). Samples, standards, and blank were kept for an hour to develop their color. After an hour, the absorbance of all the samples, standards and blank was read on spectrophotometer set at the wavelength of 410 nm (Chapman and Pratt, 1961). A standardized curve was designed against the spectrophotometer reading and respective P concentration in standards. This curve was used to find out the unknown concentration of P in the samples. The following formula was used to determine the P concentration in the sample.

$$P (\%) = (\text{Conc. of P (ppm) from curve}) * (V_0 / V_1 * W_t * 100)$$

Where;

V_0 = Total ml of digested sample

V_1 = Volume of the sample (ml) used for P determination

W_t = Plant material weight used for digestion (g)

2.6. Determination of K in Plants

For K determination in plants, Filtrate sample after di-acid digestion was used. Standards of K (10, 20, 40, 60, 80 and 100 ppm) were prepared by using KCl salt. The samples and the standards were run on flame photometer and emissions of these samples were noted. A standardized curve was designed against the flame photometer reading and respective K concentration in standards. This curve was used to find out the unknown concentration of K

in the samples. The following formula was used to determine the K concentration in the sample.

$$K \text{ (ppm)} = (\text{Conc. of K (ppm) from curve}) * (V/W_t)$$

Where;

V = Total ml of digested sample.

W_t = Plant material weight used for digestion (g).

2.7. Determination of SO_4 in Plants and Grains

For the determination of SO_4 in plants and grains material, the extraction process was done first before using the turbidimetric method (discussed on the above portion). For extraction, 1g of sample (plant material and grain) was taken in evaporating basin. 10 ml of magnesium nitrate [$Mg(NO_3)_2 \cdot 6H_2O$] was applied on this sample. Then, placed this sample on hot plate at $180^\circ C$. When the sample was completely dried, the hot plate temperature was raised at $280^\circ C$. At this temperature, the color of the sample changed from brown into yellow. Then, sample was kept in a muffle furnace at $450^\circ C$ overnight. Sample had put out from the furnace and cooled. Then, 10 ml of HCl was used in this sample and boiled the sample for 3 min. After boiling, 10 ml of DI water was used in it. Then sample was shifted in 100 ml Volumetric flask. It was up to the marked with DI water and then filtered through whatman filter paper number 541. The Extraction process was completed here, further the turbidimetric method was used for SO_4 determination in plant and grains.

2.8. Statistical analysis

Complete Randomize Design (CRD) with three replications was used. Data from the experiment was analyzed using ANOVA, while the significance of treatment means was separated by applying the LSD test at a 5 % level of significance (Steel and Torrie, 1996).

3. Results

3. Growth Parameters

3.1. Plant Height

Data showed that the treatments significantly ($P < 0.05$). Maximum plant height (98 cm) was noted with T1 and minimum with T2 (95.667 cm). When the ratio between Cl^- and SO_4^{2-} was decreased from T2 to T8, the wheat plant height was increased but remained less than T1 (Control). The percentage decrease in plant height as compared to T1 was 23.8%, 23.8, 23.8

%, 20.4 %, 13.6 %, 13.6 % and 6.8 % in treatments, from T2 to T8 respectively. The comparison between the treatments showed that T2, T3, and T4 treatments were statistically similar. T5, T6, T7 and T8 treatments were in between the response of treatment T1 and treatments T5, T6, T7 and T8. The effect of treatment (T1) was different than the other treatments (Figure (a)).

3.2. Yield

The data disclosed that the impact of treatments on wheat yield was significant at $P = 0.000$. Wheat yield was maximum with treatment T1 (27.25 g) while minimum with T2 (17.42 g). When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the yield of wheat was increased but remained less than T1 (Control). The percentage decrease in yield as compared to T1 was 36 %, 34.12 %, 34.75 %, 34.61 %, 29.17 %, 26 % and 25.81 % in treatments, from T2 to T8, respectively. The comparison between the treatments showed that the response of treatments (T3, T4, and T5) and (T7 and T8) on yield was relatively similar (Figure (b)).

3.3. 100-Grains Weight

Data was analyzed statistically, and it showed that the treatment's impact on the 100-grain weight was significant ($P = 0.000$). The T1 treatment has the maximum 100 grains weight (6.2033 g) while the minimum (3.2033g) was with T2. When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the grain's weight of wheat was increased but remained less than T1 (Control). The percentage decrease in gains weight as compared to T1 was 44.7 %, 41.59 %, 34.54 %, 33.9 %, 33.9 %, 26.2 %, and 17.9 % in treatments, from T2 to T8, respectively. The comparison between the treatments showed that the response of T2 and T3 treatments on wheat grain weight was relatively similar, and the T7 and T8 response was also relatively similar (Figure (b)).

3.4 Plant Nitrogen (%)

Data were analyzed statistically, and it showed that the treatment's impact on Plant N was significant ($P = 0.000$). The maximum concentration of N (0.9523%) in Plant was T1 and minimum N concentration (0.4587%) was T2. When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of N increased but remained less than T1 (Control). The percentage decrease in N concentration as compared to T1 was 51.83%, 50.43%, 49.7%, 48.96%, 46.26%, 42.45% and 40.17% in treatments, from T2 to T8 respectively. Treatments comparison showed that response of treatments (T7 and T8) on N concentration was relatively similar. All the other treatments relative response was different from each other's (Figure (d)).

3.5. Plant Phosphorus

Data was analysed statistically, and it showed that the treatments impact on plant P was significant ($P = 0.000$). Maximum concentration of P (0.0887%) was with T1 and minimum (0.0537%) with T2. The ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of P increased but remained less than T1 (Control). The percentage decrease in P concentration as compared to T1 was 39.45%, 37.65%, 37.65%, 34.27%, 32.35, 30.1% and 28.97% in treatments, from T2 to T8 respectively. The treatments comparison showed that response of all the treatments remained different from each other (Figure (e)).

3.6. Plant Potassium (%)

The data showed that the treatment's impact on Plant K was significant ($P = 0.000$). The T1 has the maximum concentration of K (0.9057 %), while treatment T2 has a minimum K concentration (0.5643 %). When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of K was not changed significantly, and it remained less than T1 (Control). The percentage decrease in K concentration as compared to T1 was 37.69 %, 36.7 %, 36.36 %, 36.81%, 35.51 %, 35.07 % and 35 % in treatments, From T2 to T8 respectively. The comparison between the treatments showed that the response of treatments (T2, T3, T4, T5, T6, T7, and T8) on K concentration was relatively similar (Figure (f)).

3.7. Soil Chloride (ppm)

Data were analyzed statistically, and it showed that the impact of the treatment on soil chloride was significant ($P = 0.000$). The maximum concentration of Cl in soil (45.66 ppm) was with T2, and treatment (T1) had a minimum Cl concentration in soil (7.2 ppm). When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of Cl also decreased but remained more than T1 (Control). The percentage increase in Cl concentration as compared to T1 was 84.23 %, 83.12 %, 77.5 %, 72.65 %, 67.76 %, 62.75 % and 59.24 % in treatments, from T2 to T8 respectively. The comparison between the treatments showed that the response of treatments (T2 and T3) was relatively similar while all treatments' relative response was different from each other's (Figure (g)).

3.8. Chloride in Plant (ppm)

Data was analysed statistically and it showed that the treatments impact on chloride concentration in wheat plant was significant ($P = 0.000$). The treatment (T2) have the maximum concentration of Cl in plant (233.67 ppm) while treatment (T1) have minimum Cl concentration in Plant (94.66 ppm). When the ratio between Cl^- and SO_4^{2-} was decreased

from treatment (T2) to treatment (T8), the concentration of Cl also decreased but remained more than T1 (Control). The percentage increase in Cl concentration as compared to T1 was 59.48 %, 52.66 %, 41.8 %, 37.99 %, 33.48 %, 25.06 % and 23.03 % in treatments, from T2 to T8 respectively. The comparison between the treatments presented that response of all the treatments was relatively different from each other's (Figure (h)).

3.9. Chloride in Wheat Grain

Data was analysed statistical and it showed that the treatments impact on soil chloride was significant ($P = 0.000$). The treatment (T2) have the maximum concentration of Cl in wheat grain (195.33 ppm) while treatment (T1) have minimum Cl concentration in wheat grain (73.66 ppm). When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of Cl also decreased but remained more than T1 (Control). The percentage increase in Cl concentration as compared to T1 was 62.28 %, 60.18 %, 58.84 %, 56.91 %, 53.66 %, 45.29 % 42.14 % in treatments, from T2 to T8 respectively. The comparison between the treatments showed that response of the treatments (T2 and T3) on Cl concentration in wheat grain was relatively similar. T7 and T8 also showed the relatively similar response on Cl concentration in wheat grain (Figure (i)).

3.10. Sulphate in Plant (ppm)

Data was analysed statistical and it showed that the treatments impact on sulphate concentration in wheat plant was significant ($P = 0.000$). The treatment (T8) has the maximum concentration of SO_4 (1256.9 ppm) while treatment (T2) has minimum SO_4 concentration (576.28 ppm). When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of SO_4 increased in wheat plant. The percentage decrease in SO_4 concentration as compared to T1 was 14.52%, 9.7% and 3.06% in Treatments, from T2 to T4 respectively. The percentage increase in SO_4 concentration as compared to T1 was 13.82 %, 30.25 %, 39.16 % and 46.36 % in treatments, from T5 to T8 respectively. The comparison between the treatments showed that response of all the treatments was relatively different from each other (Figure (j)).

3.11. Sulphate in Wheat Grains (ppm)

Data was analysed statistical and it showed that the treatments impact on sulphate concentration in wheat plant was significant ($P = 0.000$). The treatment (T8) has the maximum concentration of SO_4 (583.09 ppm) while treatment (T2) has minimum SO_4 concentration (376.43 ppm) in wheat grains. When the ratio between Cl^- and SO_4^{2-} was decreased from treatment (T2) to treatment (T8), the concentration of SO_4 increased in wheat

grain. The percentage decrease in SO_4 concentration as compared to T1 was 7.08%, 14.09% and 14.86% in treatments, from T2 to T4 respectively. The percentage increase in SO_4 concentration as compared to T1 was 7.49 %, 15.16 %, 22.62 % and 24.17 % in treatments, from T5 to T8 respectively. The comparison between the treatments showed that response of the treatments (T2 and T3) on SO_4 concentration in wheat grain was relatively similar. T7 and T8 also showed the relatively similar response on SO_4 concentration in wheat grain (Figure (k)).

4. Discussion

4.1. Physiological Parameters

The major problem of an irrigated area is soil salinity. The agricultural potential of arid and semiarid regions like in Pakistan is decreasing due to continues increase in salinity. The annual rainfall of these regains is low and net evaporation rate is high. Different types of salts (NaCl , Na_2SO_4 , NaHCO_3 and Na_2CO_3), mostly dominated with NaCl and Na_2SO_4 are accumulated on soil surface and causes the soil salinity (Blaylock *et al.*, 1994). The effects of salts on the growth and production of the Plant are investigated using single salt (NaCl) but in salinized area, different types of cations and anions are present (Song *et al.*, 2005; Hanif *et al.*, 2017, Qadir *et al.*, 2000). There are some studies, those investigate the response of different plants using mixture of all possible salts that may occur in salt-affected soils (Sosa *et al.*, 2005; Yang *et al.*, 2007; Zhang and Mu, 2009; Li *et al.*, 2010; Guo *et al.*, 2011; Lin *et al.*, 2014; Zhao *et al.*, 2014; Zhang *et al.*, 2015). There is lot of contradiction that which ion (Cl^- or SO_4^{2-}) is more toxic for plant growth and development under saline conditions. The present study examined the wheat tolerance against Cl^- and SO_4^{2-} salinity.

The parameters like respiration rate, photosynthetic rate, evaporation and evapotranspiration were affected by both types of salinities (chloride and sulphate salinity). The decrease in these parameters was due to specific ion toxicity, osmotic effect and deficiency of some nutrients. Under salinity stress, the moun of some ions (Na^+ , Ca^{2+} , Cl^- and SO_4^{2-}) were increased. This increase was toxic for plant development because more concentration of these ions inhibits the function of many enzymes during photosynthesis. Moreover, the relatively more numbers of these ions in soil solution were reduced the uptake of many important nutrients (like NH_4^+ , NO_3^- , K^+ , HPO_4^{2-} and $\text{H}_2\text{PO}_4^{2-}$) so, the deficiency of these nutrients were occurred during the salt stress. Salinity also create osmotic and oxidative stress. The osmotic

stress severely reduced the roots' uptake of major nutrients and water (Munns and Tester, 2008). These stresses reduced the stomatal conductance, photosynthesis, respiration and evapotranspiration of the wheat plant (Bie *et al.*, 2004; Akbarimoghaddam *et al.*, 2011; Hussain *et al.*, 2013; Chen *et al.*, 2016). When we compared between the two types of salinities (chloride and sulphate salinity), It was found that the effect of chloride salinity (treatments having the dominant ion was chloride) have a more negative affect on wheat physiological parameters than sulphate salinity (treatments have dominant ion was sulphate). These results favour the results of Geilfus (2018) as Cl^- ion was more mobile than SO_4^{2-} ion, the uptake of Cl^- ion was also more than SO_4^{2-} ion. The plasma membrane was more permeable for Cl^- ion than SO_4^{2-} ion. Cl^- also have an antagonistic effect with many plant nutrients (N, P and SO_4). So, it used their pathways as its concentration in soil solution was increased. When the uptake of Cl^- was increased in xylem cell, it caused an injury of leaf cells of the wheat plant. This damage of leaf cells ultimately reduced the process of photosynthesis. Slabu and his colleagues (2009) found that the field bean showed the signs of chlorosis (yellowing of the leaves) or necrosis (leaves burring) due to higher storage of Cl^- ion in chloroplast. Tavakkoli and his co-workers (2010) found that the higher concentration of Cl^- ion in chloroplast was related with the reduction or degradation of chlorophyll contents in Plant. This accumulation of Cl^- ion in chloroplast and reduction in chlorophyll contents causes several problems for photosynthesis. Higher concentration of Cl^- also caused a photo inhibitory damage of photosystem II reaction centre (Foyer *et al.*, 1994). Bose and his colleagues (2017) found that due to excessive concentration of Cl^- in chloroplast, the electron acceptors were destroyed during the photosynthesis process, which ultimately enhanced ROS production. The process of photosynthesis is an energy producing process used during respiration and other growth functions. Moreover, the chloride salinity affected the respiration and electron transport pathway during respiration (Jacoby *et al.*, 2016). Overall, the physiological parameters were more damaged by chloride salinity than sulphate salinity in my research.

4.2. Growth Parameters

Weight of the spike and stalk, plant height, number of spikes and grains weight were affected by both types of salinities (chloride and sulphate salinity). The decrease in these parameters was due to osmotic effect, specific ion toxicity and deficiency of some nutrients (Afzal *et al.*, 2006; Iqbal and Ashraf, 2007; Khan *et al.*, 2009). Under salinity stress, the concentration of certain ions (Na^+ , Ca^{2+} , Cl^- and SO_4^{2-}) was increased. This increase was toxic for plant

growth because higher amount of these ions inhibits the function of many enzymes during photosynthesis. Moreover, the higher concentrations of these ions in soil solution reduced the uptake of many important nutrients (NH_4 , NO_3 , K^+ , HPO_4^{2-} and $\text{H}_2\text{PO}_4^{2-}$). So, the deficiency of these nutrients were occurred during the salt stress. Salinity also create osmotic and oxidative stress. These stresses reduced the photosynthesis process in the wheat plant (Bie *et al*, 2004; Hussain *et al*, 2013; Alom *et al*, 2016). When we compared between the two types of salinities (chloride and sulphate salinity), it was found that the effect of chloride salinity (treatments having the dominant ion was chloride) have a more negative affect on wheat growth parameters than sulphate salinity (treatments have dominant ion was sulphate) (Minhas *et al*, 2007). Moreover, they found that with salinity level at the same electrolyte concentration, SO_4^{2-} had less impact than Cl^- on the yield of rice (*Oryza sativa* L.). Munns and Rawson (1999) found that chloride salinity had more damaging effect on wheat growth during reproductive phase. They stressed the wheat plant by applying different concentration of NaCl (100 to 170 mM). These treatments showed significant effect on the number of spikelet on per spike, emergence of the spike was delayed, and the fertility of the seeds reduced. This all resulted in poor grain yield of wheat. The possible reason behind this result was the plasma membrane was more permeable for Cl^- ion than SO_4^{2-} ion. Cl^- also have an antagonistic effect with many plant nutrients (N, P and SO_4) so, it used their pathways as its concentration in soil solution was increased. When the uptake of Cl^- was increased in xylem cell, it caused an injury of leaf cells of the wheat plant. This damage of leaf cells ultimately reduced the process of photosynthesis. As photosynthesis is an energy producing process, which was used during respiration and other plant growth functions, all the growth processes were affected by chloride salinity.

4.3. Plant Nutrients

Nitrogen (N) concentration was decreased in wheat plant under salinity stress. As the ratio between Cl^- : SO_4^{2-} increased (4:1, 3:1), the concentration of nitrogen in Plant decreased more. An antagonistic effect between Cl^- and NO_3^- was reported in many studied. This effect based on the nitrate transporters inhibition. These transporters unable to differentiate the NO_3^- over Cl^- especially when the Cl^- concentration was enhanced in soil solution (Wang *et al.*, 2012). So, Cl^- ion was taken up due to its higher concentration by these transporters instead of NO_3^- . While SO_4^{2-} did not have such an antagonistic effect with nitrogen, the concentration of N was more affected by chloride salinity than the sulphate salinity. It was concluded that chloride salinity has more negative effect for plant nitrogen than sulphate salinity.

Phosphorus (P) was reduced in wheat plant under salinity. As the ratio between Cl^- : SO_4^{2-} increased (4:1, 3:1), the concentration of nitrogen in Plant decreased more. An antagonistic effect between Cl^- and H_2PO_4^- or HPO_4^{2-} was reported in many studied. This effect was based on either the competition between these anions to occupy the binding site at phosphate transporter protein or due to more concentration of Cl^- in soil solution was displaced the phosphate from binding site. While SO_4^{2-} did not have such kind of antagonistic effect with phosphate so, the concentration of P was more affected by chloride salinity than the sulphate salinity. It was concluded that chloride salinity has more negative effect for plant phosphorus than sulphate salinity.

Potassium (K) concentration decreased in wheat plant under salinity. The concentration of K was decreased in all the treatment except control in almost same trend. It was because Plant take up K in K^+ ion so, various ratio of Cl^- : SO_4^{2-} at same concentration of Na^+ . There was a direct relation between K^+ and Na^+ and because all the treatments, the reduction in K concentration showed a similar trend in all the treatments except control. Munns and his colleagues (2006) find out that, when the Na^+ concentration enhanced, the K^+ concentration was decreased in wheat. This reduction in K^+ concentration in wheat caused to reduce its growth.

Overall, it was concluded that the chloride ion has more toxic effect than sulphate ion for wheat development. These results favored the work done by Minhas and his co-workers (2007), who concluded that the impact of Cl^- on rice yield was more than SO_4^{2-} when the electrolyte concentration was kept content. SO_4^{2-} salinity has more toxic effect than Cl^- salinity in *Brassica rapa* and also Cl^- salt resulted in less decrease of pepper plant growth than SO_4^{2-} salt (Navarro *et al*, 2003; Reich *et al*. 2018). These results also contradict the conclusion of work done by Datta and his co-workers (1995) who found that the Cl^- was less damaging than SO_4^{2-} to barley yield.

5. Conclusion

Agriculture lands are at great threat due to increase in salt affected areas. It not only disturb the soil, but also change chemical and physical processes going on in plant bodies. So, this experiment was done to check the effect of salinity with different Cl^- : SO_4^{2-} ratios on the growth parameters of wheat. It was concluded that the both types of salinities (chloride and sulphate) significantly reduced the growth of the wheat. Maximum growth reduction was observed in T2 and T3 (treatments with 4:1 and 3:1 of Cl^- and SO_4^{2-} ratios). Chloride salinity

was more injurious than sulphate salinity for wheat growth. Overall, it was concluded that the chloride ion has more toxic effect than sulphate ion for wheat growth.

Abbreviations:

Cl⁻= Chloride ion

CRD= Complete Randomized Design

DI= De-Ionized

dS m⁻¹= deciSiemens per meter

EC_e = Electrical Conductivity

ESP= Exchangeable Sodium Percentage

g= gram

K= Potassium

LSD= Least Significant Difference

ml= milliliter

N= Nitrogen

Na₂SO₄= Sodium Sulfate

NaCl= Sodium Chloride

nm= nanometer

NWFP= North West Frontier Province

P= Phosphorus

ppm= parts per million

SO₄²⁻= Sulfate ion

SP= Saturation Percentage

Declaration:

Ethics approval and consent to participate: not applicable

Consent for publication: Not applicable

Availability of data and material: The author will share data on demand.

References

- Afzal I, Basra SMA, Farooq M, Nawaz A. 2006 Alleviation of salinity stress in spring wheat by hormonal priming with ABA, salicylic acid and ascorbic acid. *Int. J. Agric. Biol.* 8: 23–28.
- Akbarimoghaddam H, Galavi M, Ghanbari A, Panjehkeh N. 2011. Salinity effects on seed germination and seedling growth of bread wheat cultivars. *Trakia J. Sci.* 9: 43-50.
- Alom, R., Hasan MA, Islam MR, Wang QF. 2016. Germination characters and early seedling growth of wheat (*Triticum aestivum* L.) genotypes under salt stress conditions. *J. Crop Sci. Biotech.* 19: 383-392.
- Aziz A, Ashraf M, Sikandar S, Asif M, Akhtar N, Shahzad SM, Wasaya A, Raza A, Babar BH. 2019. Optimizing sulfur for improving salt tolerance of sunflower (*Helianthus annuus* L.). *Soil Environ.* 38: 222-233.
- Benedetti L, Scherner A, Cuchiara CC, Moraes ÍL, Avila LA, Deuner S. 2020. Soybean plant osmotic and oxidative stress as affected by herbicide and salinity levels in soil. *Planta Daninha.* 38.
- Bie Z, Ito T, Shinohara Y. 2004. Effects of sodium sulfate and sodium bicarbonate on the growth, gas exchange and mineral composition of lettuce. *Scientia Horticulturae*, 99: 215-224.
- Blaylock AD. 1994. Soil salinity, salt tolerance and growth potential of horticultural and landscape plants. Co-operative Extension Service, University of Wyoming, Department of Plant, Soil and Insect Sciences, College of Agriculture, Laramie, Wyoming.
- Bose J, Munns R, Shabala S, Gilliam M, Pogson B, Tyerman SD. 2017. Chloroplast function and ion regulation in plants growing on saline soils: lessons from halophytes. *J. Exp. Bot.* 68: 3129-3143.
- Chapman HD, Pratt PF. 1961. Methods of analysis for soils. Plants and waters. 182-186.

- Chen W, Guo C, Hussain S, Zhu B, Deng F, Xue Y, Geng M, Wu L. 2016. Role of xylo-oligosaccharides in protection against salinity-induced adversities in Chinese cabbage. *Environ. Sci. Pollut. Res.* 23: 1254-1264.
- Datta KS, Kumar A, Varma SK, Angrish R. 1995. Differentiation of chloride and sulphate salinity on the basis of ionic distribution in genetically diverse cultivars of wheat. *J. Plant Nut.* 18: 2199-2212.
- Duan L, Dietrich D, Ng CH, Yeen Chan PM, Bhalerao R, Bennett MJ and Dinneny JR. 2013. Endodermal ABA signaling promotes lateral root quiescence during salt stress in *Arabidopsis* seedlings. *Plant Cell.* 25(1): 324-341.
- Foyer CH, Lelandais M and Kunert KJ. 1994. Photooxidative stress in plants.
- Geilfus CM. 2018. Review on the significance of chlorine for crop yield and quality. *Plant Sci.* 270: 114-122.
- Geilfus CM. 2018. Chloride: from nutrient to toxicant. *Plant Cell Physiol.* 59(5): 877-886.
- Gould I, Waegemaeker J de, Tzemi D, Wright I, Pearson S, Ruto E, Karrasch L, Christensen LS, Aronsson H, Eich-Greatorex S and Bosworth G. 2021. Salinization threats to agriculture across the North Sea region. *Fut. Sust. Agric. Saline Environ.* 71-92.
- Guo R, Zhou J, Hao W, Gong D, Zhong X, Gu F, Liu Q, Xia X, Tian J, Li H. 2011. Germination, growth, photosynthesis and ionic balance in *Setaria viridis* seedlings subjected to saline and alkaline stress. *Can. J. Plant Sci.* 91: 1077-1088.
- Hanif Z, Naem M, Ali HH, Tanveer A, Javaid MM, Peerzada AM, Chauhan BS. 2017. Effect of environmental factors on germination of *Salsola foetida*: potential species for rehabilitation of degraded rangelands. *Rangel. Ecol. Manag.* 70: 638-643.
- Helal HM, Mengel K. 1979. Nitrogen metabolism of young barley plants as affected by NaCl-salinity and potassium. *Plant and Soil.* 51(4).
- Hu H, Liu H, Liu F. 2018. Seed germination of hemp (*Cannabis sativa* L.) cultivars responds differently to the stress of salt type and concentration. *Ind. Crops Prod.* 123: 254-261.
- Hussain S, Khaliq A, Matloob A, Wahid MA., Afzal I. 2013. Germination and growth response of three wheat cultivars to NaCl salinity. *Soil Environ.* 32: 36-43.

- Iqbal M, Ashraf M. 2007. Seed treatment with auxins modulates growth and ion partitioning in salt-stressed wheat plants. *J. Integr. Plant Biol.* 49: 1003–1015.
- Jacoby RP, Che-Othman MH, Millar AH, Taylor NL. 2016. Analysis of the sodium chloride-dependent respiratory kinetics of wheat mitochondria reveals differential effects on phosphorylating and non-phosphorylating electron transport pathways. *Plant Cell Environ.* 39: 823-833.
- Jixiang Lin. 2012. Salinity-alkalinity tolerance in wheat: Seed germination, early seedling growth, ion relations and solute accumulation. *African J. Agric. Res.* 7(3).
- Khan HA, Pervez MA, Ayub CM, Ziaf K, Balal RM, M.A. Shahid, N. Akhtar. 2009. Hormonal priming alleviates salt stress in hot pepper (*Capsicum annuum* L.). *Soil Environ.* 28:130–135.
- Kumar P, Sharma PK. 2020. Soil Salinity and Food Security in India. *Frontiers in Sustainable Food Systems.* 4-174.
- Li R, Shi F, Fukuda K, Yang Y. 2010. Effects of salt and alkali stresses on germination, growth, photosynthesis and ion accumulation in alfalfa (*Medicago sativa* L.). *Soil Sci. Plant Nut.* 56: 725-733.
- Lin J, Mu C, Wang Y, Li Z, Li X. 2014. Physiological adaptive mechanisms of *Leymus chinensis* during germination and early seedling stages under saline and alkaline conditions. *J. Animal Plant Sci.* 24: 904-912.
- Machado RMA, Serralheiro RP. 2017. Soil salinity: Effect on vegetable crop growth. management practices to prevent and mitigate soil salinization. *Horticulturae.* 3(2): 30.
- Massa D, Mattson NS, Lieth HJ. 2009. Effects of saline root environment (NaCl) on nitrate and potassium uptake kinetics for rose plants: A Michaelis-Menten modelling approach. *Plant and Soil.* 318: 1–2.
- Minhas PS, Dubey SK, D. Sharma R. 2007. Effects on soil and paddy–wheat crops irrigated with waters containing residual alkalinity. *Soil use and manage.* 23: 254- 261.
- Munns R, James RA, Läuchli A. 2006. Approaches to increasing the salt tolerance of wheat and other cereals. *J. Exp. Bot.* 57: 1025-1043.

- Munns R, Tester M. 2008. Mechanisms of salinity tolerance. *Ann. Rev. Plant Biol.* 59: 651-681.
- Munns, R. 2005. Genes and salt tolerance: bringing them together. *New phytol.* 167(3): 645-663.
- Munns, R, Rawson HM. 1999. Effect of salinity on salt accumulation and reproductive development in the apical meristem of wheat and barley. *Aust. J. Plant Physiol.* 26: 459-464.
- Navarro JM, Garrido C, Carvajal M, Martinez V. 2002. Yield and fruit quality of pepper plants under sulphate and chloride salinity. *J. Horti. Sci. Biotech. Physiol. Plant.* 92: 696-717.
- Qadir M, Ghafoor A, Murtaza G. 2000. Amelioration strategies for saline soils: A review. *Land Degradation and Development.* 11(6).
- Reich M, Aghajanzadeh T, Helm J, Parmar S, Hawkesford MJ, de Kok LJ. 2017. Chloride and sulfate salinity differently affect biomass, mineral nutrient composition and expression of sulfate transport and assimilation genes in *Brassica rapa*. *Plant and Soil.* 411: 1-2.
- Reich M, Aghajanzadeh T, Stuiver CEE, Koralewska A, de Kok LJ. 2015. Impact of sulfate salinity on the uptake and metabolism of sulfur in Chinese cabbage. In *Molecular physiology and ecophysiology of sulphur*. Springer, Cham. pp. 227-238.
- Reich M, Aghajanzadeh TA, Parmar S, Hawkesford MJ, De Kok LJ. 2018. Calcium ameliorates the toxicity of sulfate salinity in *Brassica rapa*. *J. Plant Physiol.* 231: 1-8.
- Renault S, Croser C, Franklin JA, Zwiazek JJ. 2001 Effects of NaCl and Na₂SO₄ on red-osier dogwood (*Cornus stolonifera Michx*) seedlings. *Plant and Soil.* 233(2).
- Richards LA. 1954 Diagnosis and Improvement of Saline and Alkali Soils. *Soil Sci.* 78(2).
- Sanower Hossain M, Sultan Ahmad Shah J. 2019. Present scenario of global salt affected soils, its management and importance of salinity research article information. *Int. Res. J. Biol. Sci. Perspective.*
- Slabu C, Zörb C, Steffens D, Schubert S. 2009. Is salt stress of faba bean (*Vicia faba*) caused by Na⁺ or Cl⁻ toxicity?. *J. Plant Nutr. Soil Sci.* 172(5): 644-51.

- Song J, Feng G, Tian C, Zhang F. 2005. Strategies for adaptation of *Suaeda physophora*, *Haloxylon ammodendron* and *Haloxylon persicum* to a saline environment during seed germination stage. *Ann. Bot.* 96: 399-405.
- Sosa L, Llanes A, Reinoso H, Reginato M, Luna V. 2005. Osmotic and specific ion effects on the germination of *Prosopis strombulifera*. *Ann. Bot.* 96(2).
- Steel RGD, Torrie JH. 1996. Principles and procedures of statistics-a biometrical approach (3rd ed). McGraw Hill Book Company Inc. New York, USA.
- Syed A, Sarwar G, Shah SH, Muhammad S. 2021. Soil Salinity Research in 21st Century in Pakistan: Its Impact on Availability of Plant Nutrients, Growth and Yield of Crops. *Communications in Soil Science and Plant Analysis.* 52(3): 183-200.
- Tavakkoli E, Rengasamy P, McDonald GK. 2010. High concentrations of Na⁺ and Cl⁻ ions in soil solution have simultaneous detrimental effects on growth of faba bean under salinity stress. *J. exp. Bot.* 61: 4449-4459.
- Tavares DS, Fernandes TEK, Rita YL, Rocha DC, Sant'Anna-Santos BF, Gomes MP. 2021. Germinative metabolism and seedling growth of cowpea (*Vigna unguiculata*) under salt and osmotic stress. *South African J. Bot.* 139.
- Wang Q, Lu X, Chen X, Malik WA, Wang D, Zhao L. 2021. Transcriptome analysis of upland cotton revealed novel pathways to scavenge reactive oxygen species (ROS) responding to Na₂SO₄ tolerance. *Scientific Reports.* 11(1).
- Wang YY, Hsu PK, Tsay YF. 2012. Uptake, allocation and signaling of nitrate. *Trends Plant Sci.* 17: 458–467.
- Yan QQ, Zhang JS, Li XX, Wang YT. 2019. Effects of salinity stress on seed germination and root growth of seedlings in island cotton. *Acta Agronomica Sinica China.* 45(1).
- Yang C, Chong J, Li C, Kim C, Shi D, Wang D. 2007. Osmotic adjustment and ion balance traits of an alkali resistant halophyte *Kochia sieversiana* during adaptation to salt and alkali conditions. *Plant Soil.* 294: 263-276.
- Zhang H, Zhang G, Lü X, Zhou D, Han X. 2015. Salt tolerance during seed germination and early seedling stages of 12 halophytes. *Plant Soil.* 388: 229-241.

- Zhang H, Zhao Y. 2011. Effects of different neutral and alkaline salinities on seed germination and early seedling growth of maize (*Zea mays* L.). African J. Agric. Res. 6(15).
- Zhang JT, Mu CS. 2009. Effects of saline and alkaline stresses on the germination, growth, photosynthesis, ionic balance and anti-oxidant system in an alkali-tolerant leguminous forage *Lathyrus quinquenervius*. Soil Sci. Plant Nut. 55: 685-697.
- Zhao Y, Lu Z, He L. 2014. Effects of saline-alkaline stress on seed germination and seedling growth of *Sorghum bicolor* (L.) Moench. Appl. Biochem. Biotechnol. 173(7).
- Zhao Y, Lu Z, He L. 2014. Effects of saline-alkaline stress on seed germination and seedling growth of *Sorghum bicolor* (L.) Moench. Appl. Biochem. Biotech. 173: 1680-1691.

UNDER PEER REVIEW

Table 1. Characteristics of Experimental Soil.

Characteristics	Unit	Results
Texture		Sandy clay loam
EC_e	dS m ⁻¹	2.2
TSS	mmol _c L ⁻¹	22
pH_s		7.48
CO₃⁻	mmol _c L ⁻¹	0
HCO₃⁻	mmol _c L ⁻¹	6
Cl⁻	mmol _c L ⁻¹	11
SO₄²⁻	mmol _c L ⁻¹	5
Ca+Mg²⁺	mmol _c L ⁻¹	14
Na⁺	mmol _c L ⁻¹	8
SAR		2.65
Saturation percentage	%	34




Figure 1. T1= Control, T2=4:1 of Cl⁻: SO₄²⁻, T3=3:1 of Cl⁻: SO₄²⁻, T4=2:1 of Cl⁻: SO₄²⁻, T5=1:1 of Cl⁻: SO₄²⁻, T6=1:2 of Cl⁻: SO₄²⁻, T7=1:3 of Cl⁻: SO₄²⁻ and T8=1:4 of Cl⁻: SO₄²⁻. Mean sharing the same letter(s) do not differ significantly at P ≤ 0.05. (a) Effect of treatments on plant height. (b) Effect of treatments on wheat yield. (c) Effect of treatments on 100 grain weight of wheat crop. (d) Effect of treatments on N concentration in wheat straw. (e) Effect of treatments on P concentration in wheat. (f) Effect of treatments on the concentration of K in wheat. (g) Effect of treatment on the concentration of Cl in soil. (h) Effect of treatments on the concentration of Cl in wheat plant. (i) Effect of treatments on the concentration of Cl in wheat grains. (j) Effect of treatments on the concentration of SO₄ in wheat plant. (k) Effect of treatments on the concentration of SO₄ wheat grain.