

Original Research Article

Effect of different sources of silicon on the growth, yield and Si uptake in aerobic rice

ABSTRACT

A field experiment was conducted to study the different sources of silicon (Si) on plant growth and yield. The experiment was laid in randomized complete design (RCBD) with four treatments and four replications. The experiment results revealed that There was an increase in plant height and the number of tillers plant⁻¹ with the application of different sources of Si over RDF (T₁). At harvest RHB @ 4 t ha⁻¹ recorded significantly higher plant height (106.05±1.23 cm) and the number of tillers hill⁻¹ (15.25±2.75). Yield attributes like number of panicles hill⁻¹ (13.23±0.49), Panicle length 23.43±0.29 cm, test weight (23.55±0.05 g), straw (7.15±0.59 t ha⁻¹), grain (3.60±0.16 t ha⁻¹) and total biomass yield (10.75±1.01 t ha⁻¹) was recorded in treatment receiving SA @ 4 mL L⁻¹ (T₃). Whereas, higher Si content and uptake in both straw and grain was significantly higher in the recorded in the with the application of the RHB @ 4 t ha⁻¹. Thus, combined application of external Si sources and along with recommended dose of fertilizers found to increase the growth, yield and Si uptake in aerobic rice.

Keywords: Silicon, Rice, Diatomaceous earth, Silicic acid, Rice husk biochar

INTRODUCTION

In Karnataka, rice is grown in an area of 1.19 mha with an annual production of 3011 kg ha⁻¹ [1]. Rice cultivation is the most water consuming system and utilizes about 60 % of total available irrigation water. Aerobic rice system in a new term given by International Rice Research Institute (IRRI) for drought-tolerant, input-responsive, lodging-resistant and weed-competitive rice varieties grown under non-flooded conditions in non-puddled and unsaturated (aerobic) soil, which is responsive to nutrient supply, can be rainfed or irrigated and tolerates (occasional) flooding [2]. Maximization of aerobic rice yield could be achieved by the balanced use of fertilizers particularly

Comment [U1]: *The method was not clearly stated in the abstract section.*

Comment [U2]: *The introduction section should be more focused on the previous works done on this topic.*

major nutrients viz., nitrogen (N), phosphorus (P) and potassium (K) in optimum quantity. However, various abiotic and biotic factors are the main challenges and threats to aerobic rice production. Inclusion of the silicon (Si) fertilizers along with the recommended dose of fertilizers (RDF) will not only increase the production and productivity of aerobic rice [3-4] but also resistance to abiotic and biotic stress [5]. Hence, a field experiment was conducted with three different sources of Si to evaluate its effect on yield and uptake.

Rice is a typical Si accumulator plant that accumulates up to 10 % Si their aboveground biomass more than the major nutrients [6]. Most of the traditional rice fields are deficient in plant-available Si (PASi) [7]. Although various Si fertilizers have been reported for the higher growth and yield in crops such as diatomaceous earth [7], silicic acid [8], CaSiO_3 [4] and rice husk biochar [9].

However, the comparative study of different sources of Si on aerobic rice is lacking. In this context, a comparative study of these Si sources on the growth, yield and Si uptake in aerobic rice was taken.

MATERIAL AND METHODS

A field experiment was conducted at C- Block, ZARS, V. C. Farm, Mandya during *Summer*, 2018. The latitude, longitude and altitude of Mandya is 19° N, 76°E, 695 m above MSL respectively. The soil of experimental soil is neutral in reaction with sandy loam in texture. Acetic acid and calcium chloride extractable silicon were medium in range. Available nitrogen and potassium content of the soil is medium in range whereas available phosphorus is very high. Secondary and micronutrient is higher than the critical limit (Table 1).

The experiment was carried out following randomized complete block design (RCBD) with four treatments and four replications. The source of Si used in this study was concentrated soluble silicic acid (SA), diatomaceous earth (DE) and rice husk biochar (RHB). The composition of DE and RHB is presented in Table 2 and concentrated soluble silicic acid (SA), obtained from ReXil Agro BV, Chennai, India, which contains 2 per cent Si as soluble H_4SiO_4 (Table 3). The entire dose of P

and K were applied as per the treatments and N was applied in two split doses. The entire dose of DE and RHB was applied as basal before sowing whereas the SA was sprayed at 15 days of interval.

Treatment details:

T₁: RDF alone

T₂: T₁+ DE @ 300 kg ha⁻¹,

T₃: T₁ + SA @ 4 mL L⁻¹ and

T₄: T₁ + RHB @ 4 t ha⁻¹

Note: RDF- Recommended Dose of Fertilizers

DE- Diatomaceous earth

SA – Silicic Acid

RHB – Rice Husk Biochar

Five plants were randomly labelled and recorded for plant height and number of tillers at 30, 60, 90 DAT and at harvest. The height was measured from the base of the fully opened leaf or tip of the panicle, whichever is the longest. Mean of the height and tillers recorded from five plants were reported. The SPAD value and photosynthetic characteristics were measured with a SPAD-502 chlorophyll meter (Minolta, Osaka, Japan). The upper third of the rice flag leaves was used to measure the SPAD value at 60 and 90 DAS. Grains and straw yield from the corresponding net plot was sundried and the weight of grain and straw per net plot was computed and then expressed as ton per hectare.

Grain and straw samples were collected from the field after the harvest of the crop and washed with deionised water and were dried in an oven at 70 °C, powdered and analysed for Si content (Ma and Takahashi, 2002). The Si uptake by the crop was computed using Si content and expressed as kg ha⁻¹ using the following formula.

$$\text{Si uptake (kg ha}^{-1}\text{)} = (\text{Si content (\%)} \times \text{biomass (kg ha}^{-1}\text{)})/100$$

Data obtained were analysed using one way ANOVA at a 5 per cent level of significance as per the procedure outlined by [10]. Pearson's correlation and regression analysis was computed using MS-excel and SPSS 20.0.

RESULTS AND DISCUSSION

Plant height and number of tillers hill⁻¹

There was a significant ($p < 0.05$) improvement was observed on plant height with the application of DE @ 300 kg ha⁻¹ (T₂), SA @ 4 mL L⁻¹ (T₃) and RHB @ 4 t ha⁻¹ (T₄) over RDF alone (T₁) at 30, 60, 90 DAS and at harvest (Table 4). Application of RHB @ 4 t ha⁻¹ (11.66±0.55 cm) recorded significantly ($p < 0.05$) higher plant height at 30 DAS whereas, DE @ 300 kg ha⁻¹ (10.82±0.74 cm) showed on par results with the application of RHB @ 4 t ha⁻¹ and SA @ 4 mL L⁻¹ (10.04±0.13 cm) which was on par with the RDF (T₁) (9.18±0.51 cm). However, the effect of plant height at 60 and 90 DAS was recorded statistically similar among treatment receiving DE @ 300 kg ha⁻¹, SA @ 4 mL L⁻¹ and RHB @ 4 t ha⁻¹. Treatment receiving RHB @ 4 t ha⁻¹ (106.05±1.23 cm) and DE @ 300 kg ha⁻¹ (104.80±3.68 cm) recorded significantly similar results whereas, SA @ 4 mL L⁻¹ (101.75±2.68) was recorded statistically, on par with the RDF (T₁). The data presented in the Table 5 showed that treatments receiving external sources of silica *i.e.*, DE @ 300 kg ha⁻¹ (T₂), SA @ 4 mL L⁻¹ (T₃) and RHB @ 4 t ha⁻¹ (T₄) were recorded significantly ($p < 0.05$) higher number of tillers at 60, 90 DAS and at harvest over the RDF (T₁). At 60 DAS application of DE @ 300 kg ha⁻¹ (T₂), SA @ 4 mL L⁻¹ (T₃) and RHB @ 4 t ha⁻¹ (T₄) recorded statistically similar results (8.10±0.42, 8.15±0.97 and 8.20±0.63, respectively). At 90 DAS highest no of tillers was recorded in treatment receiving SA @ 4 mL L⁻¹ (11.70±1.45) and the remaining treatments were on par with each other. However, application of DE @ 300 kg ha⁻¹ (T₂), and RHB @ 4 t ha⁻¹ (T₄) recorded significantly ($p < 0.05$) higher no of tillers (14.00±2.18 and 15.25±2.75, respectively) at harvest.

There was no definite trend in the increase of plant height and number of tillers with the application of different sources of silicon. There was an increase in the plant height and number of

Comment [U3]: The data obtained should be presented in graphical form.

tillers per plant with the application of silicon sources over RDF (T_1). This can be attributed to a sufficient supply of nutrients and the beneficial effect of silicon released from the different silicon sources and thereby improvements in the nutrient use efficiency by crop. Rice husk biochar might have had a higher advantage due to its substantially higher amount of Si content (Table 2).

Si application makes the leaves and stems more erect, thus reducing self-shading and increasing photosynthesis rate, which contributes to an increase in plant height [11]. The beneficial effect of Si on the rice plant height and number of tillers are well known which has been reported by several others like Malav *et al.* (2015), Swe *et al.* (2021) and Cuong *et al.* (2017)] in rice [12-14]. Absorbed silicon is located in the leaf area in rice and this decreased the cuticle transpiration and decreases plant elongation [15]. Silicon improved plant height, inter-node length and fresh weight in rice [16]. A similar review was presented by Mauad *et al.* (2003) and Singh *et al.* (2006) for rice crops and reported an increase in the number of tillers due to the application of silicon [17-18].

SPAD Values

The addition of different sources of silicon caused a significant increase in SPAD values over RDF (T_1) in 60 and 90 DAS (Table 5). Application of SA @ 4 mL L⁻¹ (50.8±5.41) recorded significantly ($p < 0.05$) higher SPAD value at 60 DAS whereas, RHB @ 4 t ha⁻¹ (45.1±4.85) showed on par results with the application of DE @ 300 kg ha⁻¹ (47.0±6.57). At 90 DAS, SA @ 4 mL L⁻¹ (42.8±1.10) recorded significantly higher SPAD value followed by RHB @ 4 t ha⁻¹ (40.7±2.31). Application of DE @ 300 kg ha⁻¹ (36.7±2.37) recorded on par with the RDF (T_1) (38.5±2.33). Sivaranjani *et al.*, (2020) reported increase in SPAD reading with the application of silicic acid [19]. Similarly song *et al.* (2014), Haddad *et al.* (2018) and Yogendra *et al.* (2017) Increase in SPAD value and chlorophyll content by silicon fertilizers application over RDF (T_1) [4,20-21]. Rani and Narayanan (1994) reported that higher SPAD value with the application of Si along with the N could

be due to higher photosynthetic activity, better utilization of light and translocation of assimilated product to sink [22].

Yield attributes

The yield attributes were increased with the application of different sources of Si along with the RDF (Table 6). The yield attributes like number of panicles per hill, panicle length, test weight, straw, grain and total biomass yield were recorded lower in the treatment receiving RDF alone (T_1). RHB @ 4 t ha⁻¹ (14.23±0.49) recorded number of panicles per hill, whereas, higher panicle length (23.82±0.39 cm) and test weight (23.55±0.05 g) were in treatment receiving SA @ 4 mL L⁻¹ (T_3).

The straw yield was recorded significantly ($p<0.05$) higher with the application of SA @ 4 mL L⁻¹ (7.15±0.59 t ha⁻¹) whereas DE @ 300 kg ha⁻¹ (6.60±0.61 t ha⁻¹), and RHB @ 4 t ha⁻¹ (6.45±0.19 t ha⁻¹) are on par with each other but significantly ($p<0.05$) higher compared to RDF (5.90±0.58 t ha⁻¹). The total biomass yield followed the same trend as the straw yield. Application of Si sources recorded an empirical increase in the grain yield but statistically on par with the RDF (T_1) (Table 6).

The increase in the crop growth and yield attributes due to the external supply of Si in the present study are in agreement with several other workers. Accumulation of Si in the rice plant reduces the transpiration rate, thus increasing water use efficiency by the crop and improving the dry matter production [23]. The Si deposited on the leaf surface forms a protective barrier against invasion of pests and diseases as well as prevention of water losses through transpiration, imparting drought resistance [24]. Application of foliar silicic acid increased soybean yield by providing Si directly to the foliage [8]. These cumulative effects of Si on rice might have contributed to enhanced rice yield in the study. The beneficial effects of Si application *viz.*, reducing mutual shading by improving leaf erectness, decreasing susceptibility to lodging, decreased incidence of abiotic and biotic stresses, improving structural support and biomass [25] and improving nutrient uptake [26].

Chen *et al.* (2011) stated that silicon application increased grain yield by an increase in spikelet number, filled spikelet percentage and 1000-seed weight [27]. Gong *et al.* (2021) reported foliar application of nano-silica fertilisers increased the grain yield over RDF (T₁) [28]. Agostinho *et al.* (2017) reported a higher yield in rice receiving foliar Si over slag [29]. The beneficial effects of applied Si in rice had been reported in India (Singh *et al.*, 2006; Prakash *et al.*, 2011 and Malav *et al.*, 2017)[12, 18, 30], Thailand [31] and Korea [32].

Si content and uptake

The data pertaining to Si content and uptake as influenced by different sources of Si were presented in Table 7. Si content of rice straw ($4.26 \pm 0.11\%$) was significantly higher in treatment with RHB @ 4 t ha⁻¹(T₄). DE @ 300 kg ha⁻¹ ($3.31 \pm 0.51\%$) and SA @ 4 mL L⁻¹ ($2.92 \pm 0.31\%$) recorded empirically higher value but was statistically on par with the treatment receiving RDF (T₁) ($2.64 \pm 0.26\%$). Significantly higher Si uptake (275.06 ± 34.07 kg ha⁻¹) was recorded in RHB @ 4 t ha⁻¹. Si uptake in DE @ 300 kg ha⁻¹ (218.74 ± 52.97 kg ha⁻¹) was statistically on par with SA @ 4 mL L⁻¹ (208.30 ± 16.84 kg ha⁻¹). Lower Si uptake (154.49 ± 13.65 kg ha⁻¹) was recorded in RDF(T₁). Application of external sources of silicon recorded a numerical increase in the nutrient content and their uptake by the rice grain but was found to be non-significant. The total Si uptake by rice grain was also increased with the application of silicon sources over RDF *viz.*, RHB @ 4 t ha⁻¹ ($315.05 \pm 40.02\%$) and recorded higher Si content followed by DE @ 300 kg ha⁻¹ ($259.17 \pm 53.38\%$) and SA @ 4 mL L⁻¹ ($247.50 \pm 25.47\%$). A positive response of grain and straw Si content and uptake to Si fertilizer was observed. Similar results were reported by Sandhya *et al.* (2018) and Shwetakumari *et al.* (2020) [7-8]. The deficiency or sufficiency of Si in the soil is primarily determined by the rate of its replenishment in soil solution and its uptake during plant growth. Rice removes large quantities of Si (approximately 500 kg Si ha⁻¹ yr⁻¹) more than essential nutrients (N, P and K), therefore, continuous cropping without external supplementation of Si can lead to reduced plant available Si in the soil [33].

Conclusion

Comment [U4]: The conclusion section was explained too long

This may be concluded from the results that the application of different sources of Si along with the RDF increased plant growth parameters and yield over RDF (T₁). The higher plant height (106.05±1.23 cm) and the number of tillers per plant (15.25±2.75) were recorded in the RHB @ 4 t ha⁻¹. Higher straw (7.15±0.59 t ha⁻¹) and grain yield (3.60±0.16 t ha⁻¹) were recorded with the application of SA @ 4 mL L⁻¹ (T₃). Whereas, the Si content (straw: 4.26±0.11% and grain:1.15±0.11%) and uptake (Straw: 275.06±34.07 kg ha⁻¹, grain: 40.17±7.93 kg ha⁻¹ and total: 315.05±40.02 kg ha⁻¹) in straw and grain was recorded in the RHB @ 4 t ha⁻¹ (T₄). Application of DE, SA and RHB as Si sources increased the straw and grain yield over RDF. This improvement in growth, yield and Si uptake of rice with the addition of Si reduces abiotic and biotic stress. Therefore, external Si supplementation in aerobic rice can be effectively used for sustainable rice production. Further research is needed to find out the optimum levels of Si, N, P and K fertilizers for other locations with a different package of practices and rice varieties.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCE

1. Economic Survey of Karnataka. Planning, Programme Monitoring and Statistics Department, Government of Karnataka, 2020.

2. Bouman BAM, Tuang TP. Field water management to save water and increase its productivity in irrigated lowland rice. *Agricultural Water Management*, 2001;49 (1):11-30.
3. Sandhya K, Prakash NB. Effect of different sources of silicon fertilizer on dissolved silicon (DSi) in soil solution and its bioavailability for rice crop in three contrasted soils. *The Mysore Journal of Agricultural Sciences*. 2015;50(2):473-476.
4. Yogendra ND, Kumara BH, Chandrashekar N, Prakash NB, Anantha MS, Shashidhar HE. Real-time nitrogen management in aerobic rice by adopting leaf color chart (LCC) as influenced by silicon. *Journal of Plant Nutrition*. 2017;40(9):1277-1286.
5. Majumdar S, Prakash NB. Quantification of amorphous silicon by optimizing the 1% Na₂CO₃ method from intensively cultivated rice and sugarcane soils in a tropical climate. *Silicon*.2020;12:2989-3003.
6. Majumdar S, Prakash NB. Quantification and identification of the phytoliths in surface and sub surface soils of intensively cultivated rice and sugarcane crop sites. *The Mysore Journal of Agricultural Sciences*. 2017; 51(3):689-691.
7. Sandhya K, Prakash NB, 2018. Diatomaceous earth as source of silicon on the growth and yield of rice in contrasted soils of southern India. *Journal of Soil Science Plant Nutrition*. 2018;**18**(2):344-360.
8. Shwethakumari U, Pallavi T, Prakash NB. Influence of foliar silicic acid application on soybean (*Glycine max* L.) varieties grown across two distinct rainfall years. *Plants*. 2021;10:1162.
9. SHETTY R, PRAKASH NB. Effect of different biochars on acid soil and growth parameters of rice plants under aluminium toxicity. *Scientific Reports*. 2020;**10**:12249.
10. RANGASWAMY, R., 2010, A textbook of agricultural statistics. 2nd edition.
11. Yoshida S, Naveser SA, Ramirez EA. Effect of silicon on nitrogen supply on some leaf characters of rice plant. *Plant Soil*. 1969;**31**:48-56.
12. Malav JK, Patel KC, Sajid M, Ramani VP. Effect of silicon levels on growth, yield attributes and yield of rice in *typic ustochrepts* soils. *Ecology, Environment and Conservation*. 2015;21:AS205-AS208.
13. Swe MM, Mar SS, Naing TT, Zar T, Ngwe K. Effect of silicon application on growth, yield and uptake of rice (*Oryza sativa* L.) in two different soils. *Open Access Library Journal*. 2021;8: e7937
14. Cuong TX, Ullah H, Datta A, Hanh TC. Effects of silicon based fertilizer on growth, yield and nutrient uptake of rice in tropical zone of Vietnam. *Rice Science*. 2007;24(5): 283i290.

15. Datnoff LE, Snyder GH, Korndorfer GH. The use of silicon for irrigated disease management: reducing fungicide application and enhancing host plant resistance. *Studies in plant science: Silicon in agriculture*, Elsevier Science B. V, Amsterdam, the Netherlands. 2001; 171-182.
16. Fallah. Study of silicon and nitrogen effects on some physiological characters of rice. *International Journal of Agriculture and Crop Science*. 2012;4(5):238-241.
17. Mauad M, Carlos ACC, Helio GF, Juliano CC. Nitrogen and silicon fertilization of upland rice. *Scientia Agricola*, 2003;60:761-765.
18. Singh KK, Singh K, Singh R, Singh Y, Singh CS. Response of nitrogen and silicon levels on growth, yield and nutrient uptake of rice (*Oryza sativa* L.) *Oryza*. 2006;43(3):220-223.
19. Sivaranjani C, Chithra L, Baskar M, Thamizh Vendan R, Subrahmaniyan K. Influence of silicon and nitrogen on chlorophyll content of rice var. TKM-13 in Entisol. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(1): 2245-2249
20. Song A, Li P, Fan F, Li Z, Liang Y. The effect of silicon on photosynthesis and expression of its relevant genes in rice (*Oryza sativa* L.) under high zinc stress. *PLoS One*. 2014;9(11): e113782
21. Haddad C, Arkoun M, Jamois F, Schwarzenberg A, Yvin JC, Etienne P, Laine P. Silicon promotes growth of *Brassica napus* L. and delays leaf senescence induced by nitrogen starvation. *Frontiers in Plant Science*. 2018; 9:516.
22. Rani YA, Narayanan A. Role of silicon in plant growth. *Annual Review of Plant SIM Physiology and Plant Molecular biology*. 1994;1:243-262.
23. Marschner H. *Mineral nutrition of higher plants* (2nd Edition). Academic Press. 1995;899.
24. Paye W, Tubana B, Harrell D, Babu T, Kanke Y, Datnoff L. Determination of critical soil silicon levels for rice production in Louisiana using different extraction procedures. *Communications Soil Science and Plant Analysis*. 2018;49:1-12.
25. Meharg C, Meharg AA. Silicon the silver bullet for mitigating biotic and abiotic stress and improving grain quality, in rice? *Environmental and Experimental botany*. 2015;120:8-17.
26. Pati S, Pal B, Badole S, Hazra GC, Mandal B. Effect of silicon fertilization on growth, yield and nutrient uptake of rice. *Communications Soil Science and Plant Analysis*. 2016;47:284-290.
27. Chen JX, Tu NM, Yi ZX, Zhu, HL. Effect of silicon fertilizer on nitrogen utilization efficiency of super early rice. *Cereal Crop Research*. 2011;1:6.
28. Gong D, Zhang X, Yao JP, Dai G, Yug, Zhu Q, Gao Q, ZHENG W. Synergistic effects of bast fiber seedling film and nano silicon fertilizer to increase the lodging resistance and yield of rice. *Scientific report*. 2021;11:12788.

29. Agostinho FB, Tubana, BS, Martins MS, Datnoff, L. E. Effect of different silicon sources on yield and silicon uptake of rice grown under varying phosphorus rates. *Plants*. 2017;35(6):1-17
30. Prakash NB, Chandrashekar N, Mahendra C, Patil SU, Thippeshappa GN, Laane HM. Effect of foliar spray of soluble silicic acid on growth and yield parameters of wetland rice in hilly and coastal zone soils of Karnataka, South India. *Journal of Plant Nutrition*. 2011;34,1883-1893.
31. Ullah H, Luc PD, Gautam A, Datta A. Growth, yield and silicon uptake of rice (*Oryza sativa*) as influenced by dose and timing of silicon application under water-deficit stress. *Archives in Agronomy and Soil Science*. 2018;64:318-330.
32. Kim YH, Khan AL, Shinwari ZK, Kim DH, Waqas M, Kamran M, Lee IJ. Silicon treatment to rice (*Oryza sativa* L. cv 'gopumbyeo') plants during different growth periods and its effects on growth and grain yield. *Pakistan Journal of Botany*. 2012;44(3):891-897
33. Meunier JD, Guntzer F, Kirman S, Keller C. Terrestrial plant-Si and environmental changes. *Mineral Management*. 2008;72:263-267.

Table 1: Physico-chemical properties of soil of V. C. Farm, Mandya

Parameter	Content
pH (1:2.5 water)	7.1
EC (dSm ⁻¹) (1:2.5 water)	0.22
Organic carbon (g kg ⁻¹)	11.70
Particle size distribution (%)	
Sand	76.54
Silt	6.56
Clay	16.90
Textural class	Sandy loam
0.01M CaCl ₂ – Si (mg kg ⁻¹)	41.98
0.5M Acetic acid – Si (mg kg ⁻¹)	73.82
Available N (kg ha ⁻¹)	340.48

Available P ₂ O ₅ (kg ha ⁻¹)	230.31
Available K ₂ O (kg ha ⁻¹)	348.09
Exch. Ca (c mol (p ⁺) kg ⁻¹)	4.75
Exch. Mg (c mol (p ⁺) kg ⁻¹)	2.25
Micronutrients (mg kg ⁻¹)	
Zn	3.18
Mn	20.40
Fe	89.52
Cu	3.75

Table 2: Composition of diatomaceous earth (DE) and rice husk biochar (RHB)

Properties	DE	RHB
pH (1:2.5 water)	9.21	7.39
EC (dSm ⁻¹) (1:2.5 water)	0.72	1.62
Cation exchange capacity (c mol (p ⁺) kg ⁻¹)	52.00	38.63
Nutrient		
N	0.03	0.78
P	0.02	0.24

K	0.40	0.96
Si	30.00	31.00
Ca	2.70	0.36
Mg	3.25	0.31
S	0.17	0.05
Al ₂ O ₃	15.30	n.d
Mg kg⁻¹		
Fe	2.00	0.08
Mn	0.02	0.055
B	6.00	8.36
Zn	19.00	63.00
Cu	20.00	31.00
Mo	0.10	n.d
Se	1.30	n.d
Cd	0.50	n.d

DE – Diatomite; RHB – Rice hush biochars n.d – not determined

Table 3: Composition and pH of foliar silicic acid material

Composition	Content (%)
Si as soluble H ₄ SiO ₄ (%)	2.0
K as KCl (%)	1.2

B as H ₃ BO ₃ (%)	0.8
HCl (%)	1.0
Demiwater (%)	47.00
PEG* ₄₀₀ (%)	48.00

pH

pH of raw material	0.88
pH of 4 ml L ⁻¹ solution	6.00

*PEG- Poly ethylene glycol

UNDER PEER REVIEW

Table 4: Effect of different sources of Si on plant height of aerobic rice at different interval

Treatments	Plant height (cm)			
	30DAS	60 DAS	90 DAS	At harvest
T₁: RDF alone	9.18±0.50 (C)	39.28±1.38(B)	54.50±2.74(B)	95.80±3.25(B)
T₂:T₁ + DE @ 300 kg ha⁻¹	10.82±0.74(AB)	43.63±0.68(A)	62.15±3.10(A)	104.80±3.68(A)
T₃: T₁ + SA @ 4 mL L⁻¹	10.04±0.13(BC)	43.88±1.88(A)	61.25±2.93(A)	101.75±2.68(AB)
T₄: T₁ + RHB@ 4 t ha⁻¹	11.66±0.55(A)	44.15±1.04(A)	62.05±2.61(A)	106.05±1.23(A)
S.Em±	0.26	0.66	1.43	1.43
C. D. @ 5%	0.83	2.11	4.57	4.57

±Values indicated standard deviation

Mean value having same alphabets do not differ significantly at $p \leq 0.05$

Table 5: Effect of different sources of Si on number of tillers per hill and SPAD value of aerobic rice at different interval

Treatments	No. of tillers hill ⁻¹			SPAD value	
	60 DAS	90 DAS	At harvest	60 DAS	90 DAS
T₁: RDF alone	5.95±0.70(B)	8.80±0.5(B)	10.50±1.00(B)	41.4±1.76(A)	35.98±1.72(C)
T₂:T₁ + DE @ 300 kg ha⁻¹	8.10±0.42(A)	10.80±1.23(AB)	14.00±2.18(A)	45.4±4.51(A)	38.5±2.33(BC)
T₃: T₁ + SA @ 4 mL L⁻¹	8.15±0.97(A)	11.70±1.45(A)	13.75±2.14(AB)	50.8±5.41(A)	42.8±1.10(A)
T₄: T₁ + RHB@ 4 t ha⁻¹	8.20±0.63(A)	10.30±0.70(AB)	14.25±2.75(A)	47.8±4.59(A)	40.7±2.31(AB)
S.Em±	0.35	0.52	0.89	3.72	1.36
C. D. @ 5%	1.12	1.66	2.85		

±Values indicated standard deviation

Mean value having same alphabets do not differ significantly at p≤0.05

Table 6: Effect of different sources of Si on yield parameters in aerobic rice

Treatments	Number of panicles hill⁻¹	Panicle length (cm)	Test wt. (g)	Straw (t ha⁻¹)	Grain (t ha⁻¹)	Total biomass (t ha⁻¹)
T₁: RDF alone	8.43±0.50 (B)	22.81±0.85(B)	21.98±0.01(B)	5.90±0.58(B)	3.28±0.13(A)	9.18±1.23(A)
T₂:T₁ + DE @ 300 kg ha⁻¹	13.32±0.58(A)	23.43±0.29(A)	23.18±0.03(A)	6.60±0.61(AB)	3.50±0.18(A)	10.10±1.26(AB)
T₃: T₁ + SA @ 4 mL L⁻¹	12.43±1.00(AB)	23.82±0.39(A)	23.55±0.05(A)	7.15±0.59(A)	3.60±0.16(A)	10.75±1.01(A)
T₄: T₁ + RHB@ 4 t ha⁻¹	13.23±0.49(A)	23.24±0.47(A)	22.93±0.05(A)	6.45±0.19(AB)	3.55±0.21(A)	10.00±1.65(B)
S.Em±	0.89	0.52	0.28	0.27	0.09	0.90
C. D. @ 5%	12.11	1.60	0.28	0.83	NS	NS

±Values indicated standard deviation

Mean value having same alphabets do not differ significantly at p≤0.05

UNDER PEER REVIEW

Table 7: Effect of different sources of Si on Si content (%) and uptake (kg ha⁻¹) in straw and grain at harvest in aerobic rice

Treatments	Silicon (%)		Si uptake (kg ha ⁻¹)		
	Straw	Grain	Straw	Grain	Total
T₁: RDF alone	2.64±0.26(B)	1.06±0.17(A)	154.49±13.65(B)	35.20±10.63(A)	189.74±19.16(C)
T₂:T₁ + DE @ 300 kg ha⁻¹	3.31±0.51(B)	1.14±0.13(A)	218.74±52.97(AB)	40.52±9.77(A)	259.17±53.38(AB)
T₃: T₁ + SA @ 4 mL L⁻¹	2.92±0.31(B)	1.08±0.10(A)	208.30±16.84(AB)	39.01±9.10(A)	247.50±25.47(B)
T₄: T₁ + RHB@ 4 t ha⁻¹	4.26±0.11(A)	1.15±0.11(A)	275.06±34.07(A)	40.17±7.93(A)	315.05±40.02(A)
S.Em±	0.23	0.09	23.55	24.30	17.69
C. D. @ 5%	0.52	NS	51.87	NS	56.95

±Values indicates standard deviation

Mean value having same alphabets do not differ significantly at p<0.05.