

Original Research Article

Effect of seed pelleting with *Rhizobium* and nutrition management on growth parameters of cowpea [*Vigna unguiculata* (L.) Walp.]

Abstract

An experiment was laid out in a Randomized Block Design (Factorial) with different pelleting material i.e. no pelleting (P₁), *Rhizobium* liquid (P₂), *Rhizobium* powder (P₃) and *Rhizobium* liquid with jaggery (P₄) and 5 levels of nitrogen application i.e. 0% N of RDF (F₁), 40% N of RDF (F₂), 60% N of RDF (F₃), 80% N of RDF (F₄) and 100% N of RDF (F₅), altogether 20 treatment combinations of seed pelleting and nitrogen replicated thrice to find out effect on growth parameters i.e. emergence %, days to 50% flowering, plant height, number of branches per plant, number of pods per plant and number of seeds per pod of cowpea cv. Himachal Lobia-1. Among different pelleting treatments, seed pelleting with *Rhizobium* liquid exhibited better results over other treatments and all growth parameters affect significantly by pelleting treatments. Different level of nitrogen application also had significant effect on growth parameters. 80% N of recommended dose of fertilizer was best treatment over other treatments. Interaction effect of seed pelleting and nitrogen application was non-significant on growth parameters except number of branches per plant. Maximum number of branches was found in P₂F₄ (*Rhizobium* liquid + 80% N of RDF) treatment.

Keywords: Cowpea, *Rhizobium*, pelleting, Himachal Lobia-1, emergence %, vegetative growth

1. Introduction

Cowpea [*Vigna unguiculata* (L.) Walp.] is one of the most important crops which is grown commercially as a grain crop, animal fodder and vegetable crop [1] in the semi-arid regions of Africa, Asia, Southern Europe, Southern United States, and Central and South America [2,3,4]. It belongs to family Leguminosae and is native of South-Africa [5]. The vast majority

of the world's. In this regard, given the relevance of the study, it would be appropriate to consider the environmental problems arising from the excess of nitrogen fertilizers in the soil compared to the use of rhizobia which are more ecologically friendly, although they require more investment. For example, indiscriminate use of nitrogen fertilizers increases emissions of nitrous oxide, a potent cause of the greenhouse effect. Cowpea production (over 95%) takes place in sub-Saharan Africa with about 12.5 million hectares under cultivation worldwide in 2014. Asia is the second largest producing region, representing less than 3% of the global production in average over the 1993-2014 periods. In world, approximately over 5.59 million metric tons during the year 2014 was produced [6]. It is a crop of major importance to the nutrition of poor rural households whose diets tend to heavily rely on starchy foods such as millet, sorghum, maize and cassava. Therefore, it has a tremendous potential to contribute to the alleviation of malnutrition because it contains carbohydrate and proteins in huge quantity along with less quantity of fats, fiber, B-vitamins such as folic acid, thiamine, riboflavin and niacin and micronutrients such as iron and zinc [7].

Cowpea has ability to fix atmospheric nitrogen in soil at the rate of 56 kg/ha in association with symbiotic bacteria under favorable conditions [8,9] and biological nitrogen fixation leads to noteworthy reduction in production cost incurred on nitrogen fertilizers [10]. With a view to make sure the buildup of sufficient bacterial population and to have adequate symbiosis between bacteria and plant roots, *Rhizobium* is applied artificially in many ways like soil and seed applications. The application of *Rhizobium* through seeds in the form of seed pelleting is one of the effective ways to ensure the optimum symbiosis for enhanced crop productivity. Seed pelleting focuses on better establishment and increased productivity by precision sowing and addition of various materials along with filler material that improve the seed quality in many ways [11]. Keeping in view the above prospective, this study had been planned to determine the effect of seed pelleting with *Rhizobium* and nitrogen application on vegetative growth parameters of cowpea crop.

Comment [AFdS1]: As it is relevant information in global terms, it would be advisable to put more current information once, considering that 7 years have passed since 2014.

Comment [AFdS2]: In this regard, given the relevance of the study, it would be appropriate to consider the environmental problems arising from the excess of nitrogen fertilizers in the soil compared to the use of rhizobia which are more ecologically friendly, although they require more investment. For example, indiscriminate use of nitrogen fertilizers increases emissions of nitrous oxide, a potent cause of the greenhouse effect.

2. Materials and Methods

The experiment was carried out at farm of Department of Seed Science and Technology, Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan (HP) is located at an altitude of 1250 meters above mean sea level with latitude of 35.5°N and longitude of 77.8°E in the mid- hill zone of Himachal Pradesh, India. Climate of the area is generally sub-temperate and semi-humid characterized by cold winters. Generally, December and January are the coldest while; May and June are the hottest with moderate rainfall (1000-1300 mm). The soil texture was loam to clay loam having pH ranging from 6.85-7.05.

Table 1. Meteorological data on rainfall, temperature and relative humidity during the course of investigation

MONTH	MEAN TEMPERATURE (°C)	MEAN RAINFALL (mm)	RELATIVE HUMIDITY (%)
June	38.05	91.1	60
July	38.20	294.40	79
August	38.05	102.20	80
September	36.75	41.60	68
October	19.65	34.60	58

Source: Meteorological Observatory, Department of Environmental Sciences, Dr. Y S Parmar University of Horticulture and Forestry, Nauni-173 230, Solan (HP)

Genetically pure seed of cowpea cv. Himachal Lobia-1 was purchased from Directorate of Extension Education, Dr Y S P University of Horticulture and Forestry, Nauni, Solan. Liquid and powder form of *Rhizobium* culture was obtained from Soil microbiology laboratory of Department of Soil Science and water management, UHF Nauni, Solan. Seed pelleting was done at CSIR-IHBT (Institute of Himalayan Bioresource Technology) Palampur, Himachal Pradesh. For seed pelleting clay and adhesive (45% Gum Arabica) was used. There were total 20 treatment combinations ($P \times F$) and each treatment was replicated three times so there were total 60 plots of $1.8\text{m} \times 1.5\text{m}$ (i.e. 2.7 m^2) size. There were total 40 plants in every plot and spacing was $45\text{cm} \times 15\text{cm}$. Urea is applied to these plots and recommended dose for cowpea is 45 kg/hectare. Cultural practices were done according to package of practices for

vegetable crops, Directorate of Extension Education, Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan (HP). Recorded observations were emergence %, days to 50% flowering, plant height, number of branches per plant, number of pods per plant and number of seeds per pod. The statistical analysis for RBD was done as per design of experiment as suggested by [12]

3. Results and discussion

Emergence percentage of cowpea seeds was significantly affected by different seed pelleting with *Rhizobium* treatments and different dose of nitrogen application. When seeds were pelleted with *Rhizobium* (liquid form), emergence percentage was maximum (93%) and non-pelleted seeds shown only 89.67% emergence in field condition. Seed pelleted with dry form of *Rhizobium* gave 91.17% emergence. Among different dose of nitrogen application, 80% N of RDF (F₄) gave maximum (92.50%) emergence and minimum (89.79%) emergence was found in control (0% N of RDF). Interaction effect of seed pelleting and nitrogen application was non-significant on emergence of cowpea seeds. It might be due to continuous supply of nutrients to seeds present in pelleting material which helped in survival of seedlings. Similar increased emergence due to seed pelleting was also found by Raj [13] in cowpea. Such beneficial influence (higher germination percentage) of seed pelleting with *Rhizobium* was also found in Bengal gram, Moth beans, and Green gram and peas by Pawar *et al.* [14]. The hike in germination also could be attributed to higher cytokinin production due to biofertilizer which led to higher cell division. Ahmed *et al.* [15] also reported increased germination in French bean due to seed pelleting with *Rhizobium*.

Table 2: Effect of seed pelleting and nitrogen application on growth parameters of cowpea

Treatment	Total emergence (%)	Days to 50% flowering	Plant height	Number of branches per plant	Number of seeds per pod	Number of pods per plant	Number of seeds per plant
-----------	---------------------	-----------------------	--------------	------------------------------	-------------------------	--------------------------	---------------------------

Seed pelleting							
P₁	89.67 (9.52)	41.08	83.80	3.86	12.25	12.90	158.05
P₂	93.00 (9.70)	40.33	90.18	4.44	12.90	14.05	181.24
P₃	91.17 (9.60)	40.61	89.28	4.25	12.64	13.39	169.31
P₄	90.67 (9.57)	40.85	85.54	4.24	12.39	13.36	165.51
CD _{0.05}	0.06	0.38	0.43	0.08	0.20	0.16	3.08
Nitrogen application							
F₁	89.79 (9.53)	40.98	86.32	4.00	12.27	13.06	160.40
F₂	90.63 (9.57)	41.01	86.74	4.07	12.43	13.31	165.45
F₃	91.25 (9.60)	40.93	87.10	4.18	12.50	13.43	167.98
F₄	92.50 (9.67)	40.33	88.11	4.41	12.87	13.72	176.58
F₅	91.46 (9.62)	40.34	87.72	4.31	12.65	13.61	172.21
CD _{0.05}	0.07	0.43	0.48	0.09	0.23	0.18	3.44
Interaction (P × F)							
CD _{0.05}	NS	NS	NS	0.17	NS	NS	NS
P ₁ -Control, P ₂ - <i>Rhizobium</i> liquid + clay + adhesive, P ₃ - <i>Rhizobium</i> powder + clay + adhesive, P ₄ - <i>Rhizobium</i> liquid formulation + jaggary F ₁ -0 % N of RDF, F ₂ -40 % N of RDF, F ₃ -60 % N of RDF, F ₄ -80 % N of RDF, F ₅ -100 % N of RDF Figures in parenthesis are square root transformed.							

Different types of pelleting and different doses of nitrogen had significant effect on days to 50% flowering of cowpea. Among different pelleting methods, minimum (40.33) days for 50% flowering was taken when seed were pelleted with liquid form of *Rhizobium* and maximum (41.08) days was taken when seeds were not pelleted. Among different dose of nitrogen application, 50% of flowering was first occurred when 80% N of RDF was applied and maximum days for 50% flowering when 40% N of RDF was applied. Interaction effect of seed pelleting and nitrogen application was non-significant on days to 50% flowering. Reduction in days to 50% flowering might be attributed to the fact that pelleted seeds germinated early leading to faster vegetative growth of plant and flowering growth started early in comparison to plants which were germinated from non-pelleted seeds. Nitrogen is

known to stimulate vegetative growth therefore, days to 50% flowering increased with increase in nitrogen dose. Reduced days to 50% flowering due to seed pelleting with *Rhizobium* was also reported by Patel and Jadav [16] in cowpea cv. Pusa Phalguni.

Plant height of cowpea was increased when seeds were pelleted with *Rhizobium* and maximum (90.18 cm) plant height was obtained when seed were pelleted with liquid form of *Rhizobium* and minimum (83.80 cm) was found in control (P_1). Different level of nitrogen application had also significant effect on plant height of cowpea. 80% N of RDF gave maximum (88.11 cm) plant height and minimum (86.32 cm) was found in control (0% N of RDF) treatment. Interaction effect of seed pelleting and nitrogen application on plant height of cowpea was non-significant. This may be attributed to the fact that pelleting material maintained nutritive reserve and dehydrogenase activity in seeds and nitrogen is an essential part of chlorophyll and nucleic acids, which might have played major role in promoting plant growth in terms of plant height, number of branches per plant, number of seeds per pod etc. hike in plant height due to seed pelleting with *Rhizobium* was also reported by Khaitov *et al.* [17] in chickpea. Increased plant height due to nitrogen application was also reported by Upadhyay and Singh [18] in cowpea and by Starling *et al.* [19] in soybean.

Number of branches per plant was affected significantly by seed pelleting and different dose of nitrogen application. Minimum number of branches was obtained when seeds were not pelleted and 0% N of RDF was applied. When seeds were pelleted with liquid form of *Rhizobium*, maximum (4.44) branches per plant was obtained. In non-pelleted seeds there were only 3.86 (minimum) branches per plant. Increment in nitrogen dose up to 80% N of RDF also increase number of branches per plant. Minimum number of branches was found when nitrogen was not applied. Interaction of seed pelleting and nitrogen application also affected number of branches significantly. Maximum number of branches per plant (4.67) was found in P_2F_4 (*Rhizobium* culture pelleting + 80% N of RDF) which was at par with P_2F_3

(4.53) and P₂F₅ (4.58). However, minimum number of branches per plant (3.80) was found in P₁F₁ (control + 0% N of RDF). This might be due to constant and optimal supply of nitrogen and other nutrients under *Rhizobium* treatments which led to more plant height and plants with more height bear a greater number of branches. Similar results of increased number of branches per plant due to seed inoculation with *Rhizobium* and nitrogen application was reported by Yadav and Malik [20] in cowpea.

Table 3. Interaction effect of seed pelleting and nitrogen application on growth parameters of cowpea.

Interaction treatment	Total emergence (%)	Days to 50% flowering	Plant height (cm)	Number of branches per plant	Number of seeds per pod	Number of pods per plant	Number of seeds per plant
P ₁ F ₁	88.33 (9.45)	41.32	82.93	3.80	11.99	12.60	151.10
P ₂ F ₁	90.83 (9.58)	40.34	89.36	4.19	12.71	13.40	170.23
P ₃ F ₁	90.83 (9.58)	41.42	88.25	4.05	12.29	13.07	160.67
P ₄ F ₁	89.17 (9.50)	40.82	84.74	3.96	12.10	13.19	159.61
P ₁ F ₂	89.17 (9.50)	41.12	83.00	3.86	12.17	12.77	155.40
P ₂ F ₂	91.67 (9.63)	40.75	89.46	4.21	12.83	13.82	177.28
P ₃ F ₂	90.83 (9.58)	40.94	89.13	4.11	12.33	13.39	165.10
P ₄ F ₂	90.83 (9.58)	41.22	85.38	4.10	12.39	13.24	164.03
P ₁ F ₃	89.17 (9.50)	41.87	83.33	3.84	12.26	12.90	158.20
P ₂ F ₃	93.33 (9.71)	40.75	90.14	4.53	12.88	14.07	181.20
P ₃ F ₃	90.83 (9.58)	40.37	89.41	4.15	12.44	13.44	167.23
P ₄ F ₃	91.67 (9.63)	40.72	85.52	4.21	12.43	13.30	165.31
P ₁ F ₄	91.67 (9.63)	40.31	85.22	3.95	12.49	13.14	164.05
P ₂ F ₄	95.00 (9.80)	40.03	91.20	4.67	13.09	14.53	190.12
P ₃ F ₄	91.67 (9.63)	39.90	89.87	4.55	13.33	13.57	180.81
P ₄ F ₄	91.67	41.08	86.16	4.47	12.56	13.65	171.35

	(9.63)						
P₁F₅	90.00 (9.54)	40.76	84.50	3.85	12.35	13.07	161.50
P₂F₅	94.17 (9.76)	39.80	90.74	4.58	12.99	14.43	187.37
P₃F₅	91.67 (9.63)	40.39	89.73	4.37	12.80	13.49	172.74
P₄F₅	90.00 (9.54)	40.42	85.92	4.46	12.45	13.43	167.23
CD _{0.05}	NS	NS	NS	0.17	NS	NS	NS
P ₁ -Control, P ₂ - <i>Rhizobium</i> liquid + clay + adhesive, P ₃ - <i>Rhizobium</i> powder + clay + adhesive, P ₄ - <i>Rhizobium</i> liquid formulation + jaggary F ₁ -0 % N of RDF, F ₂ -40 % N of RDF, F ₃ -60 % N of RDF, F ₄ -80 % N of RDF, F ₅ -100 % N of RDF Figures in parenthesis are square root transformed.							

Number of seeds per pod and pods per plant were increased when seeds were pelleted with *Rhizobium*. Maximum number of seed per pod and pods per plant was obtained when liquid form of *Rhizobium* was used for seed pelleting over other treatments and minimum numbers were found in control (P₁). Different dose of nitrogen application also affected these growth parameters significantly. 80% N of RDF (F₄) produced maximum seeds per pods and pods per plant and in control (F₁) treatment. Interaction effect of seed pelleting and nitrogen application was non-significant on number of pods per plant and seeds per pod. The production of more pods per plant and seeds per pod with application of higher doses of nitrogen in comparison to lower doses and/or with no nitrogen also could be attributed to the fact that more biomass in terms of vegetative growth might have been put up with higher levels of nitrogen and nitrogen is known to stimulate vegetative growth and it might be also due to more number of flowers. Similar results of increased number of pods per plant due to seed pelleting with *Rhizobium* and nitrogen application were reported by Swaroop *et al.* [21] in cowpea cv. Arka Garima.

Different type of seed pelleting and different dose of nitrogen application had significant effect on number of seeds per plant. Number of seeds per plant was increased by 14.67% when seeds were pelleted with liquid form of *Rhizobium* over control. Among

different dose of nitrogen, 80% N of RDF gave maximum (176.58) number of seeds per plant which were 10% more of control (F₁). Interaction effect of seed pelleting and nitrogen application was non-significant in number of seeds per plants. The increased number of seeds per plant might be due to seed pelleting with *Rhizobium* and nitrogen doses, more nodulation because of nitrogen and *Rhizobium* pelleting has occurred which led to a greater number of seeds per pod and pods per plant thereby, increase number of seeds per plant. Similar results of increased number of seeds per plant due to nitrogen application were reported by Farahvash and Mirshekari [22] in cowpea.

4. Conclusion

It is concluded that seed of cowpea shows better emergence% and vegetative growth in term of plant height, number of branches, number of pods per plant and number of seeds per pod when they are treated with *Rhizobium*.

References:

1. Nidhi, Singh G, Kumar R, Malik A. Seed Production Technology of Cowpea and Soybean. In: Singh N, Malik A, Punia H. Seed Production Technology. 1st ed. Karnataka: INSC International Publishers; 2021
2. Singh N, Thakur AK, Kaushal R, Mehta DK, Bhardwaj RK. Effects of seed pelleting on seed quality of cowpea (*Vigna unguiculata* L.) during storage. International Journal of Economic Plants. 2018;5(2):76-79.
3. Timko MP, Ehlers JD, Roberts PA. Cowpea. In: Chittaranjan K. Pulses, sugar and tuber crops. 1st ed. Berlin: Springer; 2007.
4. Diouf D. Recent advances in cowpea [*Vigna unguiculata* (L.) Walp.] “omics” research for genetic improvement. African Journal of Biotechnology. 2011;10(15):2803-10.
5. Steenkamp ET, Stępkowski T, Przymusiak A, Botha WJ, Law IJ. Cowpea and peanut in southern Africa are nodulated by diverse *Bradyrhizobium* strains harboring nodulation genes that belong to the large pantropical clade common in Africa. Molecular Phylogenetics and Evolution. 2008;48(3):1131-44.
6. Anonymous. Annual report: Food and Agriculture Organization Corporate Statistical Database. 2014; Accessed 07 May 2022. Available: <https://www.fao.org/faostat/en/>.

Comment [AFdS3]: I would like to draw attention to the conclusion of this study. The conclusion notes: “It is concluded that seed of cowpea shows better emergence% and vegetative growth in term of plant height, number of branches, number of pods per plant and number of seeds per pod when they are treated with *Rhizobium*”. In addition to this information, to complete the study, it would be advisable in the final part of the work to pay attention to the practical meaning of the study, in addition to specific recommendations for farmers.

7. Graham RD, Welch RM, Saunders DA, Ortiz-Monasterio I, Bouis HE, Bonierbale M, et al. Nutritious subsistence food systems. *Advances in agronomy*. 2007;92:1-74.
8. Giridhar K, Raju PS, Pushpalatha G, Patra C. Effect of plant density on yield parameters of cowpea (*Vigna unguiculata* L.). *International Journal of Chemical Studies*. 2020;8(4):344-7.
9. Singh AK, Bhatt BP, Sundaram PK, Kumar S, Bahrati RC, Chandra N, Rai M. Study of site specific nutrients management of cowpea seed production and their effect on soil nutrient status. *Journal of Agricultural Science*. 2012;4(10):191-8.
10. Soumare A, Diedhiou AG, Thuita M, Hafidi M, Ouhdouch Y, Gopalakrishnan S, Kouisni L. Exploiting biological nitrogen fixation: a route towards a sustainable agriculture. *Plants*. 2020;9(8):1011.
11. Afzal I, Javed T, Amirkhani M, Taylor AG. Modern seed technology: Seed coating delivery systems for enhancing seed and crop performance. *Agriculture*. 2020;10(11):526.
12. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. 1st ed. Hoboken, New Jersey, United States: John Wiley & sons; 1984.
13. Raj AB, Raj SK, Prathapan K, Radhakrishnan NV, Swadija OK. Effect of seed invigoration on yield enhancement in grain cowpea [*Vigna unguiculata* (L.) Walp]. *Legume Research: An International Journal*. 2021;44(9):1-6.
14. Pawar VA, Pawar PR, Bhosale AM, Chavan SV. Effect of *Rhizobium* on seed germination and growth of plants. *Journal of Academia and Industrial Research*. 2014;3(2):84-8.
15. Ahmed I, Khan MA, Ahmed N, Khan N, Khan S, Marwat FY. Influence of *Rhizobium* inoculation on nodules, growth and yield of french beans cultivars. *International Journal of Bioscience*. 2016;9(6):226-33.
16. Patel BN, Jadav DK. Effect of *Rhizobium* seed inoculation, nitrogen and phosphorus on growth, nodulation, flowering and seed yield of cowpea cv. PUSA PHALGUNI (*Vigna unguiculata* Walp). *International Journal of Agricultural Sciences*. 2010;6(2):361-4.
17. Khaitov B, Kurbonov A, Abdiev A, Adilov M. Effect of chickpea in association with *Rhizobium* to crop productivity and soil fertility. *Eurasian Journal of Soil Science*. 2016;5(2):105-12.
18. Upadhyay RG, Singh A. Effect of nitrogen and zinc on nodulation, growth and yield of cowpea (*Vigna unguiculata*). *Legume Research-An International Journal*. 2016;39(1):149-51.

19. Starling ME, Wood CW, Weaver DB. Starter nitrogen and growth habit effects on late-planted soybean. *Agron. J.* 1998;90:658–662.
20. Yadav RD, Malik CV. Effect of *Rhizobium* inoculation and various sources of nitrogen on growth and yield of cowpea [*Vigna unguiculata* (L.) Walp.]. *Legume Research-An International Journal.* 2005;28(1):38-41.
21. Swaroop K, Ganeshamurthy AN, Rathore SV. Response of vegetable cowpea to P, K and *Rhizobium* inoculation under Andaman condition. *Indian Journal of Horticulture.* 2001;58(3):254-9.
22. Farahvash F, Mirshekari B. Yield and yield components of cowpea as affected by different sources and application rates of nitrogen fertilizers. *J Food Agric Environ.* 2011;9(3-4):295-8.

UNDER PEER REVIEW