

Original Research Article

Evaluation of performance of parents and their resultant hybrid for yield and quality traits in brinjal (*Solanum melongena* L.) over season under salt affected soil

Abstract

The present investigation was carried out in order to obtain information based on *per se* performance of parents and their cross combinations for genetic improvement in Brinjal. Ten parents were crossed in diallel fashion excluding reciprocal. Half diallel set of crosses and ~~all~~ their ~~all~~-possible 45 F₁'s (excluding reciprocals) in brinjal (*Solanum melongena* L.) were evaluated in Randomized Complete Block Design (RCBD) with three replications for nineteen yields and yield contributing traits during *Kharif* 2020-21 (Y₁) and 2021-22 (Y₂). Investigation was conducted at the Main Experiment Station (MES) of Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar, Kumarganj, Ayodhya (U.P.) India. The study revealed ~~that~~ highly significant differences ~~were observed~~ for most of the traits under study. Based on the *per se* performance, the parent P₉ (2.61 kg) registered highest fruit yield per plant followed by P₇ (2.49 kg) in both the year and pooled. The *per se* performance of crosses *viz.*, P₂ x P₉ (3.11 kg), P₄ x P₆ (2.66 kg), P₁ x P₈ (2.55 kg) and P₆ x P₇ (2.42 kg) ~~were~~ produced significantly highest fruit yield per plant than the general mean. These hybrids may be subjected to multi-locational and multi-seasonal trials for their release for commercial cultivation.

Key word: *Per se* performance, brinjal, yield, hybrids.

Introduction

Brinjal or eggplant (*Solanum melongena* L. 2n=24) is one of the most cultivated solanaceous vegetable, ~~which it~~ is mainly grown for its edible fruit. In India it is known by many regional name *viz.*, Baigan (Hindi), Badanekai (kannada), Vangi (Marathi), Katharikai (Tamil), Vankai (Telugu) while in worldwide it is popularly known as aubergine (France) or guinea squash. It is the most popular and major fruit vegetable crop in India and many other countries of the world. It is a perennial plant but grown as an annual crop ~~in which~~ with self-pollination ~~should be found~~ but ~~some time~~ cross pollination also occurs. According to **Zeven and Zhukovsky (1975) [15]** brinjal is claimed to be originated in India and spread to China, which became a secondary center of origin. It is ~~a flexible crop~~ adapted to different agro-

climatic regions and can be grown throughout the year. It is an important crop in the tropical regions of world and is ~~being~~ grown commonly in India, China, Turkey, Japan, Italy, Indonesia, Iraq, Syria, Spain and Phillipines.

Its immature fruits are generally used as vegetable and other culinary preparations, unripe fruit is essentially consumed as cooked vegetable in various forms and the dried shoots are used as firewood in rural areas. Brinjal is consumed ~~by in~~ many ways ~~like as~~ salad, bhaji, stuffed brinjal, bhatha, pickles etc., ~~this has~~ ~~make~~ ~~made the~~ brinjal ~~is a~~ more popular vegetables in India. Its fruits are ~~widely consumed in various culinary preparations and are~~ rich source of protective nutrients (Hedges and Lister, 2007) [6].

Comment [MM1]: Repetition

Globally, India ~~is~~ ranked second in vegetable production next to China and contributed 10.80 M ha and 196.26 MT to global vegetable area and production, respectively. In India, brinjal occupies an area of 0.758 million hectares with 13.154 million tonnes of annual production which have the 17.5 tonnes per hectare productivity (**Anonymous, 2021**) [2]. In Uttar Pradesh brinjal is ~~existence in cultivation~~ ~~cultivated~~ on an area of 0.080 million hectare with annual production of 2.75 million tonnes with 34.40 tonnes per hectare productivity. In Uttar Pradesh, Agra, Meerut, Lucknow, Kanpur, Aligarh, Chitrakoot and Gorakhpur district share more area and production in the state (**Anonymous 2018**) [1].

The peel of brinjal ~~have has~~ a great amount of anthocyanin. A higher anthocyanin and ~~low glyco-alkaloid content~~ are considered essential for good fruit quality. These antioxidant act as anti-ageing agent. In particular antioxidant found in brinjal helps prevent skin cancer. Brinjal is also a good source of bone building vitamin K and magnesium as well as heart healthy copper which give nourishment to ~~your the~~ scalp and ~~keeps~~ it healthy. White brinjal are highly beneficial for regulation of blood sugar levels in human body and also controls absorption of glucose. This makes them the best option for people suffering from diabetes.

Comment [MM2]: Is the glycol-alkaloid level low in brinjal?

A number of cultivars are under cultivation depending upon the yield, consumer preference about the colour, size and shape of the fruit. But it is not possible to have one common cultivar to suit different localities and local preferences. It is therefore required to improve the locally preferred cultivars for yield and adaptation or development of new hybrid combinations. Earlier, egg plant breeding ~~was~~ relied both on mass selection and pure line selection from land races for the development of improved varieties. Brinjal, being native to India and often cross pollinated crop, possesses considerable diversity for plant type, fruit colour, fruit shape, fruit size, yield and other quality traits (Ravali *et al.*, 2017) [9], which offers much scope for improvement through heterosis breeding.

The estimation of heterosis for yield and its component traits would be useful to identify the best hybrid combination for exploitation of superior hybrids. The exploitation of hybrid vigour has become a potential tool for improvement in eggplant Bavage *et al.*, 2005 [3] and Dharwad *et al.*, 2011 [4]. The selection of parents is the important step in heterosis breeding which could combine well and produce desirable hybrids. Several research workers reported the importance of genetic diversity in crops Reddy *et al.*, 2017 [10], Triveni *et al.*, 2017 [14], and Srivatsava *et al.*, 2019 [14] with this perception, the present investigation was carried out with the objective of studying the *per se* performance of hybrids and parents for yield and quality traits.

Materials and Methods

The present investigation was carried out at the main experiment station, Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar (Kumarganj), Ayodhya (UP), India, during *Kharif*, 2020-21 (Y₁) and 2021–22 (Y₂). The experimental farm falls under humid subtropical climate and is located between 24.47° and 26.56°N latitude and 82.12° and 83.58°E longitude at an altitude of 113 m above mean sea level. The experimental farm had saline alkali soil with pH above 8.

The experimental materials comprised of ten promising and diverse inbreds and varieties of brinjal selected on the basis of genetic variability from the germplasm stock maintained in the department. The selected parental lines *i.e.*; Balfahava (P₁), Punjab Sadabahar (P₂), NDB-3 (P₃), NDB-2 (P₄), NDB White-1 (P₅), Pusa Kranti (P₆), Pant Samrat (P₇), Mukta Keshi (P₈), NDB Sel-1 (P₉) and Co-2 (P₁₀) were crossed in **all possible cross combinations**, excluding reciprocals, during the year, 2019-20 to get 45 F₁'s for the study of heterobeltiosis and economic heterosis.

Comment [MM3]: Briefly indicate how crossing was done

The experiments were conducted in a Randomized Complete Block Design (RCBD) with three replications to assess the performance of 45 F₁'s hybrids and their 10 parental lines of brinjal. The crop was planted in two rows spaced at 75 cm apart with a plant to plant in row spacing of 60 cm. The experiments were transplanted on 12 August, 2020 and 17 August, 2021.

Observations were recorded for nineteen economic traits including biochemical traits, *viz.* days to 50% flowering, days to first fruit harvest, leaf length (cm), leaf width (cm), plant height (cm), number of primary branches per plant, harvest duration (days), fruit length (cm), fruit equatorial circumference (cm), number of fruits per plant, average fruit weight (kg), marketable fruit yield per plant (kg), unmarketable fruit yield per plant (kg), reducing

sugars (%), non-reducing sugar (%), total sugars (%), chlorophyll content (mg/g), total phenol content and total fruit yield per plant (kg). *Per se* performance were evaluated for parents and hybrids following method suggested by Panse and Sukatme, 1987 [8] for analysis of variance of experimental for nineteen yield and yield contributing traits.

Result and discussion

The success of any breeding programme depends on the identification of suitable parents and selection of proper breeding methodology for the improvement of a specific trait. The selection of parents with high *per se* performance would be of merit in producing better hybrids and hence the parents selected for crossing programme were evaluated based on their *per se* performance.

Perusal of Table-1 and 2 revealed that the mean squares due to genotypes, parents and hybrids were found highly significant for all the traits in both the seasons (Y_1 , Y_2) and over seasons (pooled) except plant height and primary branches per plant in the parents during the year 2020-21 (Y_1) while, days to 50% flowering in the parents during the year 2021-22 (Y_2). The mean squares due to parents vs. hybrids also found significant for all the traits studied during both the seasons (Y_1 , Y_2) and over seasons (pooled) except plant height, fruit equatorial circumference, unmarketable fruit yield per plant and chlorophyll content during the year 2020-21 (Y_1) while, days to first fruit harvest, fruit equatorial circumference and number of fruits per plant during the year 2021-22 (Y_2).

The pooled analysis of variance (Table-2) divided the source of variation into environments, replication, environments x replication, genotypes and environments x genotypes were found highly significant for all the traits under study except primary branches per plant, reducing sugars, non-reducing sugar and total sugars. Mean squares due to replication and environments x replication were found significant for only one traits *i.e.* primary branches per plant in replication and number of fruits per plant in environments x replication under the study. The mean squares due to genotypes and environments x genotypes were found highly significant for all the traits under study during over season pooled.

Among the parents highest *per se* performance (Table-3) for most desirable traits total fruit yield per plant were exhibited by parents P_9 and P_7 in both the season (Y_1 , Y_2) and in over season (pooled). The above mentioned genotypes may be used as donor parents in hybridization programme for developing high yielding varieties of respective groups. Some other genotypes exhibiting very high mean performance for the characters other than total

Comment [MM4]: Any reference to support this opinion

fruit yield per plant are also listed in Table-3. In this context, none of the parents found significantly early for days to 50 % flowering and days to first fruit harvest.

For quality traits, most desirable parents were P₃ followed by P₁ and P₇ for reducing sugars in both the season (Y₁, Y₂) as well as in over season pooled recorded higher values over the grand mean. Maximum non-reducing sugar among the parents was observed in parent P₁₀ in Y₁, Y₂ and in over season pooled. Maximum total sugars was exhibited by parents P₃ followed by P₁ and P₉ in both the season (Y₁, Y₂) as well as in over season pooled. Parents P₄ and P₈ in Y₁ and in over season pooled while, none of the parents in Y₂ showed significant chlorophyll content among the parents over the grand mean. Parental line P₉ in Y₁ while, the parents P₄ and P₇ in Y₂ and P₄ in over season pooled were exhibited maximum total phenol content over the grand mean.

Among the hybrids highest mean performance (Table-3) for most desirable traits total fruit yield per plant were exhibited by cross combination P₂ x P₉ followed by P₆ x P₇, P₇ x P₈, P₇ x P₉ and P₁ x P₇ in Y₁, P₂ x P₉ followed by P₆ x P₇, P₇ x P₈, P₇ x P₁₀ and P₈ x P₉ in Y₂ while, crosses P₂ x P₉ followed by P₄ x P₆, P₁ x P₈, P₆ x P₇ and P₇ x P₁₀ in over season pooled. Some other crosses exhibiting very high mean performance for the characters other than total fruit yield per plant are also listed in Table-3. In this context, the most desirable cross combination which produced significantly early 50 % flower than the grand mean were P₆ x P₇ followed by P₃ x P₇, P₈ x P₉ in Y₁, P₆ x P₇ in Y₂ and P₅ x P₆ in over season pooled. Cross combination P₃ x P₈ followed by P₆ x P₇ and P₈ x P₉ in Y₁, P₆ x P₇ and P₁ x P₂ in Y₂ and P₈ x P₉ followed by P₅ x P₆ and P₇ x P₉ in over season pooled found significantly early for days to first fruit harvest.

Among the crosses, quality traits *viz.* reducing sugars, non-reducing sugar, total sugars, chlorophyll content and total phenol content were more anticipated traits would be useful to identify the best hybrid combination for exploitation of superior hybrids. Out of forty five crosses, top three cross combination P₂ x P₄ followed by P₂ x P₉, P₃ x P₆ in Y₁; P₂ x P₄ followed by P₂ x P₉, P₃ x P₆ in Y₂ and P₂ x P₄ followed by P₂ x P₉ and P₈ x P₁₀ in over season pooled produced significantly high reducing sugars than the general mean. The cross combination P₄ x P₆ followed by P₃ x P₁₀ and P₃ x P₉ in Y₁; P₃ x P₉ followed by P₃ x P₁₀ and P₄ x P₆ in Y₂ and P₃ x P₈ followed by P₃ x P₁₀ and P₄ x P₆ in over season pooled exhibited significantly high non-reducing sugars than the general mean. Cross combination for total sugars were P₃ x P₆ followed by P₁ x P₅ and P₂ x P₆ in Y₁; P₃ x P₆ followed by P₁ x P₄ and P₃

x P₄ in Y₂ and P₃ x P₆ followed by P₄ x P₄ and P₂ x P₉ in over season pooled showed significantly highest total sugars than the general mean. Crosses which produced significantly highest chlorophyll content than the general mean were P₅ x P₆ followed by P₃ x P₇ and P₃ x P₉ in Y₁; P₅ x P₈ followed by P₃ x P₉ and P₃ x P₇ in Y₂; P₃ x P₉ followed by P₅ x P₆ and P₁ x P₅ in over season pooled. The crosses that produced significantly maximum total phenol content than the general mean were P₃ x P₈ followed by P₁ x P₈ and P₂ x P₇ in Y₁; P₁ x P₇ followed by P₂ x P₃ and P₄ x P₈ in Y₂ and P₅ x P₆ followed by P₅ x P₇ and P₂ x P₁₀ in over season pooled. The superiority of *per se* performance of parents and F₁ for various traits in seasons and over the season, pooled have also been reported by earlier ~~workers~~ researchers Kanchana, *et al.* 2021 [7]; Tripathy, *et al.* 2021 [13]; Timmareddygar, *et al.* 2021 [12]; Gadhiya, *et al.* 2016 [5].

Conclusion

It may be concluded from the present study that, on the basis of *per se* performance, four F₁ crosses *viz.* P₂ x P₉ followed by P₄ x P₆, P₁ x P₈ and P₆ x P₇ were identified as superior crosses for total fruit yield plant. These F₁ crosses may be recommended for multi-locational trials for commercialization of the crop.

References

1. Anonymous. Indian Horticulture Database. National Horticulture Board, Ministry of agriculture, Government of India, Gurgaon. 2018.
2. Anonymous. Horticulture Statistics Division, Department of Agriculture, Co-operation and Farmer's Welfare, Ministry of Agriculture, India. 2020-2021.
3. Bavage MS, Madalageri MB, and Mulge R. Hybrid performance in round fruited brinjal (*Solanum melongena* L.). Karnataka Journal of Horticulture. 2005; 1(3):95-97.
4. Dharwad NA, Patil SA and Salimath PM. Study on genetic diversity and its relation to heterosis in brinjal (*Solanum melongena* L.). Karnataka Journal of Agricultural Science. 2011; 24(2):110-113.
5. Gadhiya AD, Chaudhari KN, Rai VP and Patel AP. *Per se* Performance of brinjal (*Solanum melongena* L.) for yield and its contributing traits. Advances in Life Sciences. 2016; 5(2):449-458.
6. Hedges LJ, and Lister CE. Nutritional attributes of spinach, silver beet and eggplant. Crop and Food Research Confidential, Report No. 1928. 2007.

Comment [MM5]: Section is more of results than discussion

7. Kanchana R, Vijayalatha KR, Anitha T, Sandeep G and Paramaguru P. Per se Performance of Hybrids and Parents for Yield and Quality in Brinjal (*Solanum melongena* L.). The Pharma Innovation Journal. 2021; 10(10): 14-21.
8. Panse VG, Sukhatme PV. Genetics of quantitative characters in relation to plant breeding. Indian Journal of Genetics and Plant Breeding. 1987; 17:318-328.
9. Ravali B, Saidaiah P, Reddy RK, Shivraj N and Geetha A. Study on character association and path analysis in brinjal (*Solanum melongena* L.). Journal of Pharmacognocny and Phytochemistry. 2017; 6(6):393-397.
10. Reddy RD, Saidaiah P, Reddy KR and Pandravada SR. Mean performance of cluster bean genotypes for yield, yield parameters and quality traits. International Journal of Current Microbiology and Applied Science. 2017; 6(9): 3685-3693
11. Srivastava S, Saidaiah P, Shivraj N and Reddy KR. Yield and quality based phenotypic evaluation of germplasm of brinjal (*Solanum melongena* L.) under semi-arid conditions. International Journal of Current Microbiology and Applied Science. 2019; 8(7): 415-422.
12. Timmareddygari S, Saidaiah P, Natarajan S, Geetha A and Komatireddy RR. *Per se* performance of hybrids for yield, yield attributes and quality parameters in brinjal (*Solanum melongena* L.). International Journal of Current Microbiology and Applied Sciences. 2021; 10(2): 32-45.
13. Tripathy B, Tripathy P, Sahu GS, Dash SK, Pradhan BD, Lenka D and Das S. *Per Se* performances of brinjal (*Solanum melongena* L.) F1 crosses for fruit yield and bacterial wilt tolerance. The Pharma Innovation Journal. 2021; 10(10): 490-494.
14. Triveni D, Saidaiah P, Reddy RK and Pandravada SR. Mean performance of the parents and hybrids for yield and yield contributing traits in tomato. International Journal of Current Microbiology and Applied Science. 2017; 6(11): 613-619.
15. Zeven AC and Zhukovsky PM. Dictionary of cultivated plants and their centres of diversity. PUDOC, Wageningen. 1975; 219 p.

Table-1: ANOVA (mean squares) for a set of 10 x 10 diallel crosses for different traits in brinjal during year, 2020-21 (Y₁)

Source of variation	d.f	days to 50% flowering	Days to first fruit harvest	Leaf length	Leaf width	Plant height	Primary branches per plant	Crop duration	Fruit length	Fruit equatorial circumference	Number of fruits per plant
Replications	2	0.90	2.14	3.03	0.21	14.74	1.28	4.12	0.09	1.52	0.58
Genotypes	54	13.06**	14.38**	7.13**	5.63**	164.05**	2.16**	21.90**	22.38**	16.91**	95.07**
Parents	9	7.37*	15.22**	2.69**	2.01**	82.73	0.41	16.54**	37.60**	47.22**	127.15**
Hybrids	44	13.37**	14.14**	7.88**	6.08**	184.11**	2.41**	21.90**	18.86**	11.10**	90.30**
Parents vs. Hybrids	1	50.43**	17.29**	13.88**	18.55**	13.91	7.04**	69.98**	40.49**	0.06	16.46**
Error	108	4.19	5.84	0.81	0.15	45.50	0.21	9.33	0.34	0.50	1.55

Source of variation	d.f	Average fruit weight	Marketable fruit yield per plant	Unmarketable fruit yield per plant	Reducing sugar	Non reducing sugar	Total sugar	Chlorophyll content	Total Phenol content	Total fruit yield per plant
Replications	2	13.59	0.027	0.014	0.004	0.001	0.001	0.084	0.003	0.006
Genotypes	54	1759.66**	0.563**	0.085**	0.180**	0.025**	0.196**	0.078**	0.107**	0.733**
Parents	9	4004.24**	0.546**	0.031**	0.235**	0.032**	0.369**	0.088**	0.060**	0.633**
Hybrids	44	1303.25**	0.577**	0.098**	0.160**	0.021**	0.162**	0.078**	0.113**	0.768**
Parents vs. Hybrids	1	1640.80**	0.119**	0.000	0.557**	0.153**	0.127**	0.010	0.282**	0.108**
Error	108	27.72	0.011	0.003	0.002	0.001	0.003	0.005	0.002	0.014

*, ** Significant at 5 per cent and 1 per cent probability levels, respectively.

Table-1: ANOVA (mean squares) for a set of 10 x 10 diallel crosses for different traits in brinjal during year, 2021-22 (Y₂)(Contd.)

Source of variation	d.f	Days to 50% flowering	Days to first fruit harvest	Leaf length	Leaf width	Plant height	Primary branches per plant	Crop duration	Fruit length	Fruit equatorial circumference	Number of fruits per plant
Replications	2	0.26	0.51	1.13	0.25	3.40	0.15	4.82	0.89	0.56	0.80
Genotypes	54	9.27**	10.75**	7.76**	3.91**	183.58**	3.25**	25.32**	18.47**	14.71**	76.95**
Parents	9	5.10	9.94**	3.86**	1.43**	18.63**	1.87**	11.69**	31.78**	46.74**	114.46**
Hybrids	44	9.74**	11.05**	8.51**	4.14**	221.10**	3.38**	26.49**	15.20**	8.49**	71.00**
Parents vs. Hybrids	1	26.40**	4.61	9.72**	16.18**	17.48**	9.93**	96.72**	42.40**	0.25	0.98
Error	108	4.35	3.99	1.16	0.21	34.42	0.12	11.36	0.57	0.44	1.34

Source of variation	d.f	Average fruit weight	Marketable fruit yield per plant	Unmarketable fruit yield per plant	Reducing sugar	Non reducing sugar	Total sugar	Chlorophyll content	Total Phenol content	Total fruit yield per plant
Replications	2	26.39	0.002	0.011	0.006	0.003	0.001	0.024	0.005	0.009
Genotypes	54	1868.54**	0.563**	0.069**	0.168**	0.025**	0.188**	0.068**	0.107**	0.694**
Parents	9	5426.36**	0.477**	0.017**	0.264**	0.034**	0.407**	0.053**	0.167**	0.528**
Hybrids	44	1173.29**	0.586**	0.081**	0.139**	0.020**	0.145**	0.071**	0.096**	0.733**
Parents vs. Hybrids	1	438.50**	0.302**	0.013**	0.579**	0.148**	0.142**	0.092**	0.030**	0.494**
Error	108	26.93	0.006	0.002	0.003	0.001	0.004	0.004	0.002	0.009

*, ** Significant at 5 per cent and 1 per cent probability levels, respectively.

Table-2: ANOVA (mean squares) for a set of 10 x 10 diallel crosses for different traits in brinjal during over season pooled.

Source of variation	d.f	Days to 50% flowering	Days to first fruit harvest	Leaf length	Leaf width	Plant height	Primary branches per plant	Crop duration	Fruit length	Fruit equatorial circumference	Number of fruits per plant
Environment	1	163.10**	598.05**	5.56**	15.14**	108.10**	0.50	2581.60**	1.30**	1.65**	20.68**
Replication	2	1.09	0.55	1.96	0.37	47.51	1.11**	11.28	0.62	0.75	0.80
Env. x Rep.	2	0.20	0.07	1.03	0.06	19.36	0.35	0.22	0.31	0.08	3.53**
Genotypes	54	11.41**	14.42**	6.87**	4.77**	157.02**	2.19**	26.47**	28.82**	29.24**	107.19**
Env. x geno.	54	10.52**	8.66*	7.35**	3.52**	175.21**	2.35**	19.36**	12.39**	8.01**	78.01**
Error	216	4.38	4.83	0.91	0.16	36.52	0.16	10.46	0.47	0.39	1.51

Source of variation	d.f	Average fruit weight	Marketable fruit yield per plant	Unmarketable fruit yield per plant	Reducing sugar	Non reducing sugar	Total sugar	Chlorophyll content	Total Phenol content	Total fruit yield per plant
Environment	1	57.43**	0.662**	0.597**	0.001	0.000	0.002	0.157**	0.048**	2.419**
Replication	2	31.28	0.017	0.020	0.014	0.003	0.006	0.076	0.006	0.001
Env. x Rep.	2	12.30	0.009	0.000	0.000	0.001	0.001	0.016	0.001	0.012
Genotypes	54	2923.14**	0.591**	0.077**	0.271**	0.034**	0.315**	0.088**	0.102**	0.793**
Env. X geno.	54	1052.30**	0.418**	0.067**	0.107**	0.021**	0.116**	0.061**	0.112**	0.485**
Error	216	25.52	0.008	0.002	0.003	0.001	0.003	0.005	0.002	0.010

*, ** Significant at 5 per cent and 1 per cent probability levels, respectively.

Table-3 Mean performance, general mean, range, coefficient of variation and critical difference for nineteen characters of diallel set of 45 F₁'s and their 10 parents in Y₁ (2020-21), Y₂ (2021-22) and pooled

Sr. No.	Genotypes	Days to 50% flowering			Days to first fruit harvest			Leaf length (cm)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	46.67	45.00	48.33	67.67	64.67	68.50	13.01	13.43	12.69
2	P ₁ × P ₃	47.33	45.67	47.68	70.67	67.67	69.00	13.51	13.50	13.10
3	P ₁ × P ₄	48.00	46.00	49.50	72.67	67.67	68.17	12.84	11.61	12.29
4	P ₁ × P ₅	48.67	45.67	47.33	70.82	69.67	70.67	11.49	14.45	13.89
5	P ₁ × P ₆	51.00	49.00	48.83	71.91	70.33	70.33	13.59	13.34	13.32
6	P ₁ × P ₇	48.67	46.67	49.68	72.19	71.00	70.33	13.39	13.33	11.99
7	P ₁ × P ₈	50.67	48.67	50.00	70.67	69.33	71.33	12.99	13.07	13.02
8	P ₁ × P ₉	47.67	48.33	48.83	69.00	67.67	67.67	13.05	9.50	13.04
9	P ₁ × P ₁₀	49.67	50.33	49.67	70.33	70.33	70.67	13.28	12.91	13.37
10	P ₂ × P ₃	50.33	49.67	51.00	70.73	69.67	72.50	12.20	12.08	11.83
11	P ₂ × P ₄	49.00	47.00	46.83	72.00	69.00	68.33	9.94	9.76	11.38
12	P ₂ × P ₅	46.33	50.33	48.83	69.00	69.33	70.00	14.29	14.61	14.06
13	P ₂ × P ₆	52.33	51.33	49.67	72.01	70.33	71.50	14.36	14.09	13.47
14	P ₂ × P ₇	49.67	48.33	48.50	70.67	69.00	69.91	14.65	13.96	12.72
15	P ₂ × P ₈	49.00	49.67	52.33	72.34	68.67	70.29	11.92	12.19	12.89
16	P ₂ × P ₉	47.00	45.67	47.17	68.33	67.00	69.60	12.83	12.92	13.15
17	P ₂ × P ₁₀	47.00	48.67	49.67	71.67	69.67	70.17	11.82	11.62	12.30
18	P ₃ × P ₄	47.33	46.33	47.00	70.00	67.33	68.17	14.78	15.23	14.14
19	P ₃ × P ₅	48.67	46.33	48.00	71.63	68.33	69.33	16.20	16.63	14.95
20	P ₃ × P ₆	46.67	47.33	48.83	68.63	66.33	68.53	11.87	11.86	12.03
21	P ₃ × P ₇	45.33	48.00	48.50	67.67	66.67	69.33	12.82	13.24	11.69
22	P ₃ × P ₈	45.67	47.00	46.67	65.67	66.00	67.50	10.45	10.77	12.53
23	P ₃ × P ₉	46.67	45.00	48.67	72.33	69.00	70.50	14.65	14.57	14.45
24	P ₃ × P ₁₀	47.00	48.67	49.17	71.51	68.00	69.33	9.74	10.08	12.37
25	P ₄ × P ₅	52.33	49.67	49.33	71.00	69.67	71.00	13.78	13.41	12.66
26	P ₄ × P ₆	50.00	49.33	48.17	71.67	70.67	69.50	12.55	12.24	12.53
27	P ₄ × P ₇	50.67	48.33	47.67	73.00	69.00	70.33	12.08	12.16	11.99
28	P ₄ × P ₈	50.33	48.33	47.83	72.67	69.00	69.50	12.81	13.11	13.95
29	P ₄ × P ₉	52.67	50.00	49.33	71.93	70.33	70.98	12.49	12.33	14.26
30	P ₄ × P ₁₀	49.67	47.67	47.17	70.74	68.00	68.32	11.03	11.55	11.71
31	P ₅ × P ₆	47.67	45.67	45.50	68.05	65.33	66.50	14.56	16.46	14.64
32	P ₅ × P ₇	53.00	51.00	48.33	72.00	69.33	67.50	12.95	12.91	11.68
33	P ₅ × P ₈	47.67	45.67	46.17	67.33	68.33	70.33	16.34	14.20	14.43
34	P ₅ × P ₉	52.00	49.67	48.33	74.82	69.00	70.25	13.79	17.03	13.38
35	P ₅ × P ₁₀	52.33	49.67	51.00	71.78	71.33	71.17	17.51	13.37	13.57
36	P ₆ × P ₇	45.33	44.67	47.33	66.00	63.67	67.67	15.23	15.73	14.14
37	P ₆ × P ₈	50.00	50.00	50.33	69.00	66.33	69.67	14.07	15.53	13.81
38	P ₆ × P ₉	49.00	49.00	49.67	67.19	66.00	69.33	14.33	13.81	13.31
39	P ₆ × P ₁₀	50.33	47.67	50.17	68.31	66.67	69.30	14.58	13.50	13.00
40	P ₇ × P ₈	49.00	46.00	47.83	69.00	65.00	67.87	15.50	15.24	13.13
41	P ₇ × P ₉	49.67	46.00	46.83	68.98	65.00	66.53	14.12	13.35	13.96
42	P ₇ × P ₁₀	47.67	47.00	50.00	67.42	66.33	69.17	13.24	13.60	13.28
43	P ₈ × P ₉	45.33	46.00	46.83	66.03	65.00	66.17	14.51	14.87	15.60
44	P ₈ × P ₁₀	51.00	48.67	50.33	72.67	69.67	72.24	11.27	12.39	13.09
45	P ₉ × P ₁₀	50.00	47.67	50.00	68.65	66.67	69.39	13.66	13.25	15.38
F₁ Hybrid mean		48.93	47.90	48.61	70.19	68.05	69.43	13.34	13.31	13.20
Parents										
1	P ₁	51.67	48.67	50.17	72.33	70.00	71.17	11.95	11.61	11.78
2	P ₂	49.67	49.00	49.33	70.33	69.33	69.83	12.69	12.63	12.66
3	P ₃	53.00	50.00	51.50	68.67	67.33	68.00	12.96	13.57	13.26
4	P ₄	49.00	50.67	49.83	71.67	68.67	70.17	13.33	13.69	13.51
5	P ₅	48.67	48.00	48.33	70.33	67.00	68.67	13.30	13.97	13.64
6	P ₆	49.67	47.33	48.50	69.67	65.67	67.67	10.65	10.71	10.68
7	P ₇	51.33	49.67	50.50	73.33	69.67	71.50	12.97	13.32	13.14
8	P ₈	49.33	48.33	48.83	67.67	67.33	67.50	12.58	12.23	12.41
9	P ₉	49.00	47.00	48.00	71.00	68.00	69.50	13.82	13.61	13.72
10	P ₁₀	52.33	50.67	51.50	75.33	72.00	73.67	11.57	11.42	11.49
Parental mean		50.37	48.93	49.64	71.03	68.50	69.76	12.58	12.67	12.62
Grand mean		49.19	48.08	48.82	70.34	68.13	69.49	13.20	13.19	13.09
CV		4.16	4.33	3.60	3.43	2.93	2.67	6.83	8.17	6.27
CD 5%		3.27	3.33	2.84	3.86	3.19	2.99	1.44	1.72	1.34
Range	Lowest	45.33	44.67	45.5	65.67	63.67	66.17	9.74	9.50	10.68
	Highest	53.00	51.33	52.33	75.33	72.00	73.67	17.51	17.03	15.06

Table-3. Contd...

Sr. No.	Genotypes	Leaf width (cm)			Plant height (cm)			Primary branches /plant		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	8.46	8.42	7.87	72.63	76.27	78.72	5.13	5.30	5.61
2	P ₁ × P ₃	7.90	7.80	8.06	74.15	73.78	75.00	5.33	5.53	5.70
3	P ₁ × P ₄	9.11	7.25	7.69	72.31	76.26	68.49	6.46	6.58	6.11
4	P ₁ × P ₅	7.50	9.09	8.20	71.15	73.78	71.98	6.07	6.25	5.84
5	P ₁ × P ₆	7.75	7.71	8.14	68.31	70.26	72.06	5.92	6.01	6.17
6	P ₁ × P ₇	7.84	7.64	7.01	92.94	71.36	71.52	5.95	5.86	5.95
7	P ₁ × P ₈	7.54	8.38	7.68	69.50	68.56	86.21	6.16	7.74	6.53
8	P ₁ × P ₉	7.56	9.46	8.81	72.85	69.21	74.98	5.96	6.49	6.41
9	P ₁ × P ₁₀	7.66	7.53	8.29	70.13	90.06	71.50	6.18	5.87	6.04
10	P ₂ × P ₃	7.56	8.43	7.89	74.42	68.29	83.00	5.96	6.47	6.38
11	P ₂ × P ₄	6.03	6.65	7.56	83.77	71.62	76.56	5.92	6.29	5.71
12	P ₂ × P ₅	8.07	8.77	8.34	74.34	95.62	73.55	5.88	7.48	6.40
13	P ₂ × P ₆	9.19	9.17	9.14	71.89	80.48	83.10	6.54	6.69	6.57
14	P ₂ × P ₇	10.78	9.46	8.48	69.47	72.94	74.61	5.88	6.08	6.07
15	P ₂ × P ₈	7.54	8.20	7.98	94.17	93.88	74.92	6.18	7.87	6.90
16	P ₂ × P ₉	9.55	7.92	7.88	76.95	78.06	83.02	4.68	4.68	5.32
17	P ₂ × P ₁₀	7.62	6.84	7.19	72.11	81.53	79.00	5.00	5.18	5.68
18	P ₃ × P ₄	7.77	6.91	7.24	74.26	73.10	75.03	5.86	5.98	5.97
19	P ₃ × P ₅	9.77	9.96	8.81	68.50	68.50	69.15	6.15	6.15	6.17
20	P ₃ × P ₆	9.26	8.24	7.90	78.48	77.21	75.99	3.64	3.71	4.84
21	P ₃ × P ₇	8.28	8.35	7.19	72.16	68.17	88.77	5.86	5.41	5.67
22	P ₃ × P ₈	7.16	7.26	7.67	95.99	77.56	76.96	6.08	6.15	6.01
23	P ₃ × P ₉	11.17	10.27	9.73	84.82	93.76	81.69	4.42	4.57	5.55
24	P ₃ × P ₁₀	6.55	6.72	8.75	69.25	79.57	71.09	4.50	5.23	5.55
25	P ₄ × P ₅	7.38	8.62	8.08	70.44	91.49	84.07	6.92	6.70	6.44
26	P ₄ × P ₆	6.98	8.09	8.82	68.40	72.71	74.38	6.08	6.50	5.59
27	P ₄ × P ₇	7.91	8.81	8.22	72.84	73.97	74.30	5.38	5.57	5.28
28	P ₄ × P ₈	8.36	10.50	9.13	68.90	71.82	73.30	4.96	4.66	5.26
29	P ₄ × P ₉	10.16	11.39	10.58	72.60	76.48	68.73	6.03	6.27	6.21
30	P ₄ × P ₁₀	8.24	8.80	9.09	95.31	72.34	75.64	5.79	6.49	5.07
31	P ₅ × P ₆	7.87	9.20	8.74	66.92	68.97	80.14	6.46	4.82	5.34
32	P ₅ × P ₇	8.63	8.63	7.90	76.15	72.79	84.17	5.69	6.16	6.12
33	P ₅ × P ₈	9.03	9.13	10.15	77.05	88.12	74.20	4.88	7.46	5.94
34	P ₅ × P ₉	7.51	10.78	8.66	75.62	72.35	74.47	3.68	6.55	5.52
35	P ₅ × P ₁₀	12.76	7.66	7.52	78.26	63.57	71.14	5.63	4.59	5.75
36	P ₆ × P ₇	11.52	10.97	8.97	85.19	79.68	73.67	5.30	5.78	5.93
37	P ₆ × P ₈	11.34	8.06	7.99	72.36	71.84	82.22	3.64	4.15	4.77
38	P ₆ × P ₉	9.75	9.75	9.05	76.06	78.94	74.38	4.79	4.81	4.88
39	P ₆ × P ₁₀	8.85	9.84	10.00	79.89	91.60	81.29	4.21	3.64	4.83
40	P ₇ × P ₈	11.06	10.88	9.56	71.43	79.87	85.15	6.37	6.31	6.05
41	P ₇ × P ₉	8.90	8.81	8.34	93.08	89.98	66.34	4.88	4.85	5.66
42	P ₇ × P ₁₀	7.91	9.33	8.98	69.22	75.00	83.12	4.88	5.06	5.38
43	P ₈ × P ₉	9.31	9.38	9.20	79.28	65.76	74.02	3.71	3.83	4.35
44	P ₈ × P ₁₀	8.18	8.22	7.87	72.89	90.09	72.43	4.59	5.22	4.45
45	P ₉ × P ₁₀	8.93	8.80	10.78	70.79	70.99	72.75	3.46	3.80	4.71
F₁ Hybrid mean		8.63	8.17	8.46	75.94	76.77	76.37	5.40	5.71	5.70
Parents										
1	P ₁	7.32	7.80	7.56	81.17	76.95	79.06	5.91	6.43	6.17
2	P ₂	8.32	8.21	8.27	76.22	78.27	77.25	5.88	5.69	5.79
3	P ₃	8.13	8.36	8.24	66.71	70.05	68.38	5.65	5.71	5.68
4	P ₄	7.31	7.13	7.22	72.60	76.90	74.75	5.44	5.62	5.53
5	P ₅	8.56	8.84	8.70	75.56	75.68	75.62	6.32	6.87	6.59
6	P ₆	6.37	6.90	6.64	73.84	77.53	75.68	6.04	7.32	6.68
7	P ₇	6.99	7.28	7.13	82.36	76.81	79.58	5.31	5.06	5.19
8	P ₈	8.17	8.17	8.17	81.66	78.25	79.95	6.33	6.69	6.51
9	P ₉	9.06	8.84	8.95	71.37	74.94	73.15	6.20	6.61	6.41
10	P ₁₀	7.35	7.50	7.43	70.39	73.91	72.15	6.28	7.43	6.86
Parental mean		7.76	7.90	7.83	75.19	75.92	75.55	5.94	6.34	6.14
Grand mean		8.47	8.57	8.35	75.80	76.62	76.22	5.50	5.81	5.78
CV		4.61	5.33	4.09	8.89	7.65	7.03	8.31	5.91	5.78
CD 5%		0.62	0.73	0.57	10.79	9.38	8.68	0.73	0.55	0.53
Range	Lowest	6.03	6.65	6.64	66.71	63.57	66.34	3.46	3.64	4.35
	Highest	12.76	11.39	1.78	95.99	95.62	88.77	6.92	7.87	6.90

Table-3. Contd...

Sr. No.	Genotypes	Crop duration (Days)			Fruit length (cm)			Fruit equatorial circumference (cm)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	194.00	198.00	196.17	15.82	16.32	13.69	12.45	11.83	12.82
2	P ₁ × P ₃	193.00	198.00	192.50	13.56	13.56	15.02	10.51	10.90	9.26
3	P ₁ × P ₄	188.67	200.33	196.83	16.58	16.77	16.92	14.71	11.47	9.26
4	P ₁ × P ₅	195.67	193.67	191.83	16.64	16.48	17.84	12.59	13.14	10.82
5	P ₁ × P ₆	189.67	193.67	193.67	15.45	15.71	13.37	13.66	13.36	12.74
6	P ₁ × P ₇	185.33	196.00	193.00	12.29	15.30	14.16	13.96	12.64	14.26
7	P ₁ × P ₈	191.33	190.00	190.33	13.35	12.93	11.90	15.20	13.55	14.67
8	P ₁ × P ₉	191.33	196.00	195.00	15.11	13.90	13.64	12.30	14.36	15.76
9	P ₁ × P ₁₀	194.67	199.67	196.33	13.09	12.99	13.72	16.12	15.69	16.72
10	P ₂ × P ₃	196.00	200.67	194.67	16.97	16.57	11.92	7.75	8.29	10.44
11	P ₂ × P ₄	192.00	197.33	195.83	18.47	18.14	16.98	12.59	12.62	12.53
12	P ₂ × P ₅	193.67	196.00	194.50	13.89	15.53	14.55	12.59	10.75	10.63
13	P ₂ × P ₆	191.67	196.67	194.83	13.26	13.95	15.26	16.44	15.93	15.32
14	P ₂ × P ₇	188.00	198.33	196.50	14.20	13.89	15.27	10.01	12.66	12.62
15	P ₂ × P ₈	190.67	195.67	194.33	17.04	16.81	16.13	14.56	14.56	14.11
16	P ₂ × P ₉	190.00	195.00	191.33	16.72	16.21	14.25	11.63	12.29	13.13
17	P ₂ × P ₁₀	192.00	197.33	194.83	12.74	12.95	13.15	12.94	13.27	14.24
18	P ₃ × P ₄	188.67	194.00	192.67	18.72	17.92	16.52	10.56	11.29	11.79
19	P ₃ × P ₅	191.33	196.33	195.50	18.29	17.64	15.37	9.39	10.05	13.09
20	P ₃ × P ₆	188.33	193.33	194.00	17.62	17.13	17.05	9.94	10.98	9.36
21	P ₃ × P ₇	189.00	194.00	193.00	14.35	13.83	16.15	11.67	11.56	12.07
22	P ₃ × P ₈	198.00	168.67	199.00	13.13	13.47	13.68	11.32	11.61	12.10
23	P ₃ × P ₉	190.67	198.67	195.17	16.70	16.10	14.68	10.96	12.29	14.37
24	P ₃ × P ₁₀	193.67	198.67	194.50	15.34	14.85	14.52	11.95	12.19	11.10
25	P ₄ × P ₅	195.00	200.00	195.33	17.29	16.55	16.80	11.35	11.20	12.88
26	P ₄ × P ₆	191.67	197.00	193.50	18.19	17.46	17.09	11.76	12.06	11.84
27	P ₄ × P ₇	196.67	200.67	197.50	17.07	16.74	14.74	11.60	11.76	12.35
28	P ₄ × P ₈	194.67	199.33	195.67	13.17	13.03	15.88	13.14	12.50	11.53
29	P ₄ × P ₉	192.33	197.00	194.17	16.04	15.29	16.79	12.55	12.23	10.81
30	P ₄ × P ₁₀	194.00	198.00	193.17	13.99	13.99	15.80	12.14	12.13	11.03
31	P ₅ × P ₆	190.67	200.33	194.67	16.37	11.17	12.76	13.98	12.91	12.29
32	P ₅ × P ₇	191.33	196.33	197.17	14.74	14.66	13.90	15.10	15.10	13.31
33	P ₅ × P ₈	196.33	195.67	193.17	11.65	16.37	16.54	13.49	13.46	12.31
34	P ₅ × P ₉	194.00	198.00	195.83	13.82	12.09	13.71	12.78	13.23	12.59
35	P ₅ × P ₁₀	193.33	199.33	197.17	12.09	13.53	15.41	13.14	13.36	12.36
36	P ₆ × P ₇	191.33	196.67	194.17	17.19	19.27	18.73	13.21	12.88	12.32
37	P ₆ × P ₈	195.33	199.33	199.83	12.63	12.93	15.00	12.46	15.96	13.78
38	P ₆ × P ₉	192.33	196.67	195.67	13.02	12.54	12.85	13.99	13.96	13.55
39	P ₆ × P ₁₀	195.67	200.33	196.33	12.89	13.30	14.67	15.99	12.34	12.44
40	P ₇ × P ₈	197.00	201.67	197.83	17.26	16.48	15.23	10.88	10.51	11.33
41	P ₇ × P ₉	192.67	194.67	192.67	12.85	12.85	14.61	12.17	16.61	15.30
42	P ₇ × P ₁₀	189.67	196.67	195.67	12.85	12.67	13.17	16.85	12.11	13.61
43	P ₈ × P ₉	197.00	201.00	198.67	10.89	11.23	11.44	13.89	14.29	13.89
44	P ₈ × P ₁₀	190.33	195.33	194.67	7.72	11.16	12.49	14.74	15.28	14.03
45	P ₉ × P ₁₀	197.33	201.33	197.33	8.92	8.44	10.26	13.88	13.36	13.25
F₁ Hybrid mean		192.58	196.70	195.03	14.67	14.68	14.74	12.78	12.77	12.66
Parents										
1	P ₁	194.33	199.33	196.83	11.05	11.29	11.17	13.82	13.88	13.85
2	P ₂	187.00	194.33	190.67	16.47	16.79	16.63	7.63	7.32	7.47
3	P ₃	193.33	198.00	195.67	17.07	17.20	17.13	7.04	7.44	7.24
4	P ₄	190.00	194.00	192.00	19.20	17.45	18.32	8.49	8.16	8.33
5	P ₅	192.33	198.00	195.17	11.03	10.90	10.96	12.13	11.63	11.88
6	P ₆	190.00	194.33	192.17	13.03	13.63	13.33	15.88	15.16	15.52
7	P ₇	190.33	195.00	192.67	10.88	11.98	11.43	15.78	15.65	15.71
8	P ₈	192.00	197.00	195.50	13.39	12.99	13.19	17.15	16.91	17.03
9	P ₉	191.00	195.33	193.17	14.44	14.32	14.38	17.74	17.99	17.87
10	P ₁₀	189.00	194.00	191.50	7.27	7.14	7.30	12.58	12.52	12.55
Parental mean		190.93	195.93	193.53	13.38	13.36	13.38	12.82	12.66	12.74
Grand mean		192.28	196.56	194.76	13.43	14.44	14.49	12.78	12.75	12.68
CV		1.58	1.70	1.33	4.06	5.24	3.79	5.52	5.22	4.36
CD 5%		4.88	5.93	4.19	0.93	1.21	0.89	1.13	1.06	0.90
Range	Lowest	185.33	168.27	190.33	7.27	7.14	7.3	7.04	7.32	7.24
	Highest	198.00	201.67	199.83	19.20	19.27	18.73	17.74	17.99	17.87

Table-3. Contd...

Sr. No.	Genotypes	Number of fruits /plant			Average fruit weight (g)			Marketable fruit yield per plant (kg)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	16.90	16.57	16.24	108.47	108.69	115.56	1.04	1.00	1.29
2	P ₁ × P ₃	20.24	22.58	26.28	80.07	80.44	65.04	1.24	1.21	1.03
3	P ₁ × P ₄	31.20	22.92	27.88	78.12	84.71	69.86	1.48	1.33	1.18
4	P ₁ × P ₅	22.55	28.42	27.06	83.77	77.93	67.06	1.34	1.44	1.12
5	P ₁ × P ₆	23.18	23.18	23.02	78.54	78.42	77.85	1.09	1.09	1.20
6	P ₁ × P ₇	31.69	26.55	22.47	102.87	76.95	92.21	1.81	1.39	1.43
7	P ₁ × P ₈	23.88	28.60	24.27	112.46	95.78	109.72	1.92	1.82	1.87
8	P ₁ × P ₉	23.83	22.88	19.35	76.26	107.39	117.03	1.37	1.93	1.73
9	P ₁ × P ₁₀	31.36	30.93	24.17	58.46	63.07	107.32	1.13	1.19	1.59
10	P ₂ × P ₃	34.63	34.75	32.90	45.75	43.58	56.90	0.94	0.97	0.97
11	P ₂ × P ₄	26.62	25.58	21.24	66.33	65.16	86.82	1.17	1.23	1.13
12	P ₂ × P ₅	21.81	31.62	25.93	111.35	63.77	71.92	1.92	1.45	1.34
13	P ₂ × P ₆	21.83	22.77	26.99	73.69	78.61	78.36	1.20	1.22	1.35
14	P ₂ × P ₇	33.22	22.58	22.57	63.90	111.03	97.40	1.43	1.93	1.64
15	P ₂ × P ₈	21.32	21.52	22.35	79.73	80.39	79.47	1.42	1.45	1.27
16	P ₂ × P ₉	35.88	35.62	33.65	101.30	102.35	102.61	2.40	2.46	2.13
17	P ₂ × P ₁₀	27.84	29.11	26.50	60.59	60.95	86.70	1.16	1.21	1.56
18	P ₃ × P ₄	36.45	36.91	30.37	51.75	51.93	64.10	1.28	1.30	1.34
19	P ₃ × P ₅	18.29	18.26	24.81	81.49	79.53	69.00	1.15	1.18	1.16
20	P ₃ × P ₆	18.27	18.68	26.66	82.38	83.17	64.46	1.21	1.25	1.09
21	P ₃ × P ₇	31.73	26.61	26.61	55.99	63.86	65.9	1.34	1.39	1.28
22	P ₃ × P ₈	19.34	19.76	20.78	85.13	85.65	98.50	1.13	1.19	1.55
23	P ₃ × P ₉	18.26	20.15	20.99	109.15	110.52	92.11	1.44	1.48	1.34
24	P ₃ × P ₁₀	25.26	24.96	29.09	58.95	59.91	61.90	0.83	0.86	1.14
25	P ₄ × P ₅	22.21	22.48	21.90	67.47	68.44	74.08	0.92	0.93	1.18
26	P ₄ × P ₆	25.48	24.87	30.38	76.56	76.10	88.70	1.34	1.43	1.91
27	P ₄ × P ₇	19.65	20.19	24.02	75.70	76.94	68.77	1.18	1.23	1.19
28	P ₄ × P ₈	17.31	17.43	26.94	101.53	103.35	77.55	1.44	1.46	1.37
29	P ₄ × P ₉	16.46	17.67	17.98	73.41	72.20	76.85	0.81	0.84	1.00
30	P ₄ × P ₁₀	22.11	23.47	20.87	67.36	65.60	73.99	0.91	0.92	1.06
31	P ₅ × P ₆	21.92	17.21	24.47	105.34	91.13	73.56	1.66	1.24	1.29
32	P ₅ × P ₇	19.53	20.35	19.85	104.25	103.44	94.28	1.39	1.41	1.27
33	P ₅ × P ₈	16.78	26.79	22.52	91.38	105.32	107.23	1.20	1.69	1.57
34	P ₅ × P ₉	25.18	20.65	22.95	101.99	68.53	63.74	2.28	0.93	0.88
35	P ₅ × P ₁₀	19.86	25.55	23.88	70.08	102.33	84.90	0.90	2.30	1.61
36	P ₆ × P ₇	30.34	27.76	26.62	115.21	116.07	96.32	2.18	2.22	1.78
37	P ₆ × P ₈	26.23	23.78	21.71	87.10	84.97	80.34	1.72	1.21	1.20
38	P ₆ × P ₉	20.88	26.42	21.86	103.43	104.47	103.00	1.84	1.91	1.68
39	P ₆ × P ₁₀	23.80	26.68	21.57	82.49	89.03	81.22	1.17	1.74	1.28
40	P ₇ × P ₈	21.88	23.31	22.71	127.14	127.81	97.58	2.28	2.34	1.62
41	P ₇ × P ₉	21.41	19.35	20.63	135.60	84.33	94.83	2.26	0.92	1.29
42	P ₇ × P ₁₀	16.63	21.80	20.67	84.50	122.87	113.56	0.89	2.28	1.84
43	P ₈ × P ₉	24.27	24.02	20.40	116.35	114.84	103.11	2.28	2.33	1.77
44	P ₈ × P ₁₀	18.35	18.56	21.87	95.95	96.57	99.28	1.35	1.38	1.83
45	P ₉ × P ₁₀	25.52	26.36	23.11	73.54	77.12	73.60	1.28	1.34	1.12
F₁ Hybrid mean		23.81	24.14	23.97	85.84	85.89	82.67	1.42	1.44	1.38
Parents										
1	P ₁	15.92	17.59	16.75	122.43	128.97	125.70	1.58	1.64	1.61
2	P ₂	29.98	32.73	31.36	49.65	43.19	46.42	0.84	0.88	0.86
3	P ₃	32.85	33.85	33.55	55.02	47.93	51.48	1.03	1.02	1.03
4	P ₄	25.69	27.51	26.60	56.19	52.33	54.26	0.80	0.83	0.81
5	P ₅	22.86	23.44	23.15	77.27	75.42	76.35	1.32	1.30	1.31
6	P ₆	18.38	20.41	19.39	107.47	97.33	102.40	1.47	1.45	1.46
7	P ₇	19.94	21.92	20.93	123.66	123.99	123.83	1.92	1.85	1.89
8	P ₈	15.82	17.58	16.70	126.68	123.86	125.27	1.53	1.53	1.53
9	P ₉	17.41	18.74	18.08	151.57	160.72	156.15	1.99	1.89	1.94
10	P ₁₀	31.05	29.60	30.33	70.23	47.45	58.84	0.97	0.94	0.95
Parental mean		22.99	24.33	23.68	94.02	90.12	92.07	1.35	1.33	1.33
Grand mean		23.66	24.17	23.92	87.33	86.66	84.38	1.40	1.42	1.37
CV		5.25	4.79	4.30	6.02	5.98	5.28	7.40	5.32	5.79
CD 5%		1.99	1.85	1.67	8.42	8.30	7.45	0.16	0.12	0.13
Range	Lowest	15.82	16.57	16.24	45.75	43.19	46.42	0.80	0.83	0.81
	Highest	36.45	36.91	33.65	151.57	160.72	156.15	2.40	2.46	2.13

Table-3. Contd...

Sr. No.	Genotypes	Unmarketable fruit yield per plant (kg)			Reducing sugar			Non reducing sugar		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	0.79	0.81	0.60	1.22	1.24	1.28	0.53	0.53	0.51
2	P ₁ × P ₃	0.43	0.50	0.55	1.14	1.15	1.15	0.45	0.40	0.38
3	P ₁ × P ₄	0.53	0.58	0.53	0.69	1.25	1.31	0.41	0.59	.55
4	P ₁ × P ₅	0.55	0.57	0.55	1.24	0.78	0.91	0.63	0.39	0.44
5	P ₁ × P ₆	0.48	0.53	0.43	0.95	0.99	0.68	0.53	0.55	0.39
6	P ₁ × P ₇	0.87	0.60	0.51	0.94	1.09	1.09	0.58	0.62	0.57
7	P ₁ × P ₈	0.38	0.79	0.69	1.17	0.95	1.10	0.55	0.58	0.55
8	P ₁ × P ₉	0.56	0.46	0.48	1.07	1.18	1.21	0.65	0.53	0.46
9	P ₁ × P ₁₀	0.47	0.57	0.62	0.97	1.01	1.10	0.60	0.60	0.55
10	P ₂ × P ₃	0.51	0.58	0.53	1.23	1.21	1.13	0.51	0.49	0.55
11	P ₂ × P ₄	0.44	0.52	0.65	1.33	1.38	1.30	0.41	0.41	0.47
12	P ₂ × P ₅	0.26	0.56	0.49	0.98	0.69	0.91	0.63	0.52	0.49
13	P ₂ × P ₆	0.36	0.48	0.51	1.30	1.30	0.99	0.52	0.48	0.45
14	P ₂ × P ₇	0.46	0.35	0.45	0.63	1.06	1.15	0.53	0.63	0.63
15	P ₂ × P ₈	0.25	0.35	0.42	1.08	1.14	1.05	0.52	0.52	0.52
16	P ₂ × P ₉	0.98	1.09	0.98	1.32	1.35	1.14	0.46	0.46	0.52
17	P ₂ × P ₁₀	0.62	0.73	0.56	1.00	1.04	1.10	0.63	0.63	0.59
18	P ₃ × P ₄	0.66	0.74	0.65	1.21	1.23	1.15	0.60	0.60	0.60
19	P ₃ × P ₅	0.28	0.48	0.47	0.88	0.90	0.93	0.41	0.41	0.52
20	P ₃ × P ₆	0.24	0.39	0.45	1.32	1.30	1.26	0.56	0.57	0.54
21	P ₃ × P ₇	0.35	0.46	0.45	0.74	0.74	1.03	0.51	0.55	0.48
22	P ₃ × P ₈	0.52	0.61	0.44	0.99	0.97	0.97	0.49	0.49	0.56
23	P ₃ × P ₉	0.61	0.72	0.54	0.73	0.80	1.05	0.67	0.67	0.60
24	P ₃ × P ₁₀	0.72	0.80	0.63	0.69	0.75	0.69	0.67	0.64	0.58
25	P ₄ × P ₅	0.51	0.65	0.45	0.54	0.62	0.85	0.49	0.49	0.51
26	P ₄ × P ₆	0.36	0.52	0.75	0.67	0.86	1.09	0.69	0.63	0.54
27	P ₄ × P ₇	0.24	0.34	0.48	0.99	1.04	1.02	0.48	0.45	0.54
28	P ₄ × P ₈	0.32	0.40	0.53	0.78	0.80	1.00	0.50	0.50	0.55
29	P ₄ × P ₉	0.34	0.42	0.35	0.97	0.98	0.93	0.58	0.58	0.49
30	P ₄ × P ₁₀	0.52	0.62	0.43	0.91	0.90	1.11	0.44	0.41	0.49
31	P ₅ × P ₆	0.74	0.36	0.36	0.60	0.82	0.78	0.59	0.30	0.41
32	P ₅ × P ₇	0.37	0.41	0.46	0.66	0.71	0.85	0.58	0.53	0.51
33	P ₅ × P ₈	0.30	0.81	0.71	0.80	0.62	0.68	0.30	0.60	0.63
34	P ₅ × P ₉	0.28	0.48	0.60	0.60	0.64	0.66	0.57	0.59	0.63
35	P ₅ × P ₁₀	0.42	0.35	0.43	0.65	0.63	0.59	0.60	0.57	0.53
36	P ₆ × P ₇	0.90	0.92	0.64	0.85	0.83	0.75	0.47	0.47	0.58
37	P ₆ × P ₈	0.60	0.77	0.50	1.13	1.01	1.00	0.56	0.51	0.49
38	P ₆ × P ₉	0.48	0.54	0.43	0.79	0.88	0.83	0.64	0.63	0.56
39	P ₆ × P ₁₀	0.71	0.66	0.50	1.04	1.10	1.04	0.54	0.54	0.56
40	P ₇ × P ₈	0.59	0.64	0.58	1.18	1.11	1.01	0.44	0.44	0.44
41	P ₇ × P ₉	0.49	0.59	0.67	0.98	0.75	0.68	0.40	0.49	0.54
42	P ₇ × P ₁₀	0.53	0.55	0.46	0.77	0.94	0.80	0.49	0.40	0.49
43	P ₈ × P ₉	0.39	0.46	0.38	1.14	1.05	0.92	0.56	0.56	0.43
44	P ₈ × P ₁₀	0.37	0.45	0.37	1.31	1.29	0.94	0.45	0.44	0.51
45	P ₉ × P ₁₀	0.60	0.61	0.51	0.87	0.92	0.78	0.53	0.52	0.56
F₁ Hybrid mean		0.50	0.57	0.52	0.96	0.98	0.97	0.53	0.52	0.52
Parents										
1	P ₁	0.39	0.54	0.47	1.33	1.37	1.35	0.50	0.50	0.50
2	P ₂	0.59	0.63	0.61	1.16	1.21	1.18	0.36	0.36	0.36
3	P ₃	0.48	0.52	0.50	1.36	1.36	1.36	0.50	0.52	0.51
4	P ₄	0.53	0.57	0.55	1.05	1.07	1.06	0.49	0.49	0.49
5	P ₅	0.34	0.43	0.38	0.37	0.36	0.37	0.24	0.23	0.24
6	P ₆	0.42	0.46	0.44	1.08	1.04	1.06	0.52	0.51	0.51
7	P ₇	0.58	0.62	0.60	1.24	1.31	1.28	0.43	0.43	0.43
8	P ₈	0.50	0.54	0.82	1.23	1.24	1.24	0.38	0.33	0.36
9	P ₉	0.68	0.67	0.68	1.20	1.27	1.24	0.50	0.49	0.49
10	P ₁₀	0.48	0.52	0.50	1.05	1.09	1.07	0.61	0.58	0.60
Parental mean		0.50	0.55	0.55	1.10	1.13	1.12	0.45	0.51	0.44
Grand mean		0.50	0.57	0.53	0.98	1.01	1.00	0.52	0.51	0.50
CV		10.48	8.53	8.11	4.76	5.75	4.39	6.26	6.51	4.45
CD 5%		0.08	0.07	0.07	0.07	0.09	0.07	0.05	0.05	0.05
Range	Lowest	0.24	0.34	0.35	0.37	0.36	0.37	2.24	0.23	0.24
	Highest	0.98	1.09	0.98	1.36	1.38	1.36	0.69	0.67	0.63

Table-3. Contd...

Sr. No.	Genotypes	Total sugar			Chlorophyll content (mg/g)			Total Phenol content		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	1.75	1.77	1.80	1.38	1.47	1.35	1.38	1.19	1.06
2	P ₁ × P ₃	1.59	1.55	1.53	1.09	1.19	0.70	0.94	0.82	0.97
3	P ₁ × P ₄	1.10	1.84	1.86	1.39	1.32	1.30	1.19	1.13	1.13
4	P ₁ × P ₅	1.87	1.18	1.35	1.29	1.40	1.45	0.83	1.15	1.03
5	P ₁ × P ₆	1.48	1.54	1.08	1.13	1.17	1.17	0.99	1.08	0.95
6	P ₁ × P ₇	1.52	1.71	1.65	1.40	0.98	1.14	1.12	1.44	1.14
7	P ₁ × P ₈	1.72	1.53	1.60	1.32	1.42	1.28	1.42	1.12	1.12
8	P ₁ × P ₉	1.72	1.71	1.66	0.93	1.38	1.41	0.83	0.76	0.83
9	P ₁ × P ₁₀	1.57	1.61	1.65	1.22	1.27	1.12	0.79	1.16	1.15
10	P ₂ × P ₃	1.74	1.70	1.68	1.03	1.07	1.14	1.21	1.41	1.09
11	P ₂ × P ₄	1.74	1.79	1.77	1.03	1.09	1.24	1.12	0.94	1.16
12	P ₂ × P ₅	1.60	1.21	1.40	1.42	1.34	1.21	0.86	0.99	0.96
13	P ₂ × P ₆	1.82	1.78	1.44	1.23	1.25	1.31	0.86	0.91	1.05
14	P ₂ × P ₇	1.16	1.69	1.78	1.29	1.47	1.38	1.41	0.90	0.86
15	P ₂ × P ₈	1.60	1.66	1.57	1.22	1.26	1.20	1.32	0.84	0.92
16	P ₂ × P ₉	1.78	1.80	1.66	1.31	1.39	1.39	0.88	1.21	1.16
17	P ₂ × P ₁₀	1.63	1.67	1.70	1.05	1.11	1.21	1.08	1.12	1.27
18	P ₃ × P ₄	1.80	1.83	1.77	1.11	1.16	1.04	1.09	0.86	0.84
19	P ₃ × P ₅	1.29	1.31	1.44	1.01	1.07	1.15	0.94	0.86	0.82
20	P ₃ × P ₆	1.88	1.87	1.80	1.22	1.28	1.16	1.11	0.94	1.07
21	P ₃ × P ₇	1.25	1.29	1.52	1.48	1.48	1.25	1.35	1.09	1.11
22	P ₃ × P ₈	1.48	1.46	1.53	1.34	1.35	1.39	1.44	0.88	0.87
23	P ₃ × P ₉	1.40	1.47	1.65	1.45	1.51	1.37	0.86	1.16	1.01
24	P ₃ × P ₁₀	1.36	1.38	1.27	1.17	1.21	1.25	0.76	0.88	1.15
25	P ₄ × P ₅	1.03	1.11	1.36	1.06	1.09	1.15	1.25	0.82	1.07
26	P ₄ × P ₆	1.35	1.48	1.63	1.00	1.08	1.19	1.16	1.32	1.10
27	P ₄ × P ₇	1.47	1.48	1.55	1.22	1.28	1.17	0.94	0.87	0.97
28	P ₄ × P ₈	1.28	1.29	1.55	1.31	1.29	1.20	0.87	1.35	1.22
29	P ₄ × P ₉	1.55	1.56	1.42	1.38	1.41	1.21	1.13	1.21	1.07
30	P ₄ × P ₁₀	1.35	1.31	1.60	1.33	1.31	1.27	0.98	0.94	1.02
31	P ₅ × P ₆	1.19	1.12	1.19	1.49	1.44	1.46	0.88	1.26	1.30
32	P ₅ × P ₇	1.24	1.24	1.36	1.36	1.40	1.37	1.19	1.13	1.28
33	P ₅ × P ₈	1.10	1.22	1.31	1.41	1.54	1.49	1.21	1.01	0.94
34	P ₅ × P ₉	1.17	1.23	1.29	1.03	0.95	1.06	1.32	1.19	0.98
35	P ₅ × P ₁₀	1.25	1.20	1.12	0.91	1.09	1.07	0.82	0.94	1.10
36	P ₆ × P ₇	1.32	1.30	1.33	1.43	1.44	1.22	0.94	0.86	1.01
37	P ₆ × P ₈	1.69	1.52	1.50	1.35	1.25	1.24	1.17	1.32	1.13
38	P ₆ × P ₉	1.43	1.51	1.40	1.37	1.41	1.36	1.16	1.17	1.02
39	P ₆ × P ₁₀	1.58	1.64	1.59	1.22	1.39	1.38	0.99	0.99	1.06
40	P ₇ × P ₈	1.62	1.55	1.45	1.09	1.12	1.22	0.88	1.33	1.15
41	P ₇ × P ₉	1.37	1.24	1.22	1.03	1.21	1.35	1.17	1.19	1.04
42	P ₇ × P ₁₀	1.26	1.33	1.29	1.20	1.08	1.22	1.23	0.99	1.09
43	P ₈ × P ₉	1.71	1.61	1.36	1.42	1.42	1.41	1.34	1.17	1.19
44	P ₈ × P ₁₀	1.76	1.73	1.45	1.32	1.35	1.19	1.19	1.22	1.27
45	P ₉ × P ₁₀	1.40	1.44	1.34	1.39	1.42	1.17	0.99	0.88	0.85
F₁ Hybrid mean		1.49	1.50	1.49	1.24	1.28	1.24	1.08	1.07	1.05
Parents										
1	P ₁	1.82	1.87	1.85	1.23	1.26	1.25	0.94	0.78	0.86
2	P ₂	1.51	1.57	1.54	0.96	0.97	0.97	1.12	1.12	1.12
3	P ₃	1.88	1.88	1.88	1.28	1.31	1.30	1.13	0.83	0.98
4	P ₄	1.53	1.56	1.55	1.49	1.34	1.42	0.91	1.42	1.16
5	P ₅	0.62	0.59	0.60	1.17	1.18	1.18	0.83	0.94	0.88
6	P ₆	1.60	1.55	1.57	1.30	1.32	1.31	0.84	1.12	0.98
7	P ₇	1.67	1.74	1.71	1.15	1.18	1.17	1.12	1.38	1.25
8	P ₈	1.61	1.57	1.59	1.43	1.35	1.39	0.90	0.83	0.86
9	P ₉	1.69	1.76	1.73	0.97	1.02	1.00	1.15	1.12	1.13
10	P ₁₀	1.66	1.67	1.67	1.21	1.24	1.23	0.78	0.79	0.79
Parental mean		1.56	1.57	1.56	1.22	1.21	1.22	0.97	1.03	1.00
Grand mean		1.50	1.51	1.51	1.24	1.27	1.24	1.06	1.06	1.04
CV		3.83	4.11	3.23	5.60	5.10	4.58	4.28	4.46	3.06
CD 5%		0.09	0.09	0.08	0.11	0.10	0.09	0.07	0.07	0.05
Range	Lowest	0.62	0.59	0.60	0.91	0.95	0.7	0.76	0.76	0.79
	Highest	1.88	1.88	1.88	1.49	1.54	1.49	1.44	1.44	1.30

Table-3. Contd...

Sr. No.	Genotypes	Total fruit yield per plant (kg)		
		Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	1.83	1.81	1.90
2	P ₁ × P ₃	1.67	1.71	1.57
3	P ₁ × P ₄	2.06	1.91	1.71
4	P ₁ × P ₅	1.89	2.04	1.69
5	P ₁ × P ₆	1.57	1.62	1.64
6	P ₁ × P ₇	2.68	1.99	1.95
7	P ₁ × P ₈	2.30	2.60	2.55
8	P ₁ × P ₉	1.94	2.39	2.21
9	P ₁ × P ₁₀	1.60	1.76	2.21
10	P ₂ × P ₃	1.44	1.55	1.50
11	P ₂ × P ₄	1.60	1.75	1.79
12	P ₂ × P ₅	2.18	2.00	1.84
13	P ₂ × P ₆	1.57	1.70	1.88
14	P ₂ × P ₇	1.89	2.28	2.09
15	P ₂ × P ₈	1.67	1.80	1.68
16	P ₂ × P ₉	3.38	3.54	3.11
17	P ₂ × P ₁₀	1.78	1.94	2.12
18	P ₃ × P ₄	1.94	2.04	1.99
19	P ₃ × P ₅	1.43	1.66	1.63
20	P ₃ × P ₆	1.45	1.64	1.54
21	P ₃ × P ₇	1.66	1.78	1.69
22	P ₃ × P ₈	1.64	1.80	1.99
23	P ₃ × P ₉	2.06	2.21	1.89
24	P ₃ × P ₁₀	1.55	1.66	1.77
25	P ₄ × P ₅	1.42	1.58	1.62
26	P ₄ × P ₆	1.70	1.95	2.66
27	P ₄ × P ₇	1.42	1.57	1.68
28	P ₄ × P ₈	1.76	1.86	1.90
29	P ₄ × P ₉	1.15	1.26	1.35
30	P ₄ × P ₁₀	1.42	1.54	1.50
31	P ₅ × P ₆	2.40	1.60	1.63
32	P ₅ × P ₇	1.76	1.82	1.73
33	P ₅ × P ₈	1.50	2.50	2.28
34	P ₅ × P ₉	2.56	1.41	1.48
35	P ₅ × P ₁₀	1.32	2.65	2.03
36	P ₆ × P ₇	3.08	3.14	2.42
37	P ₆ × P ₈	2.32	1.97	1.70
38	P ₆ × P ₉	2.32	2.45	2.10
39	P ₆ × P ₁₀	1.88	2.41	1.78
40	P ₇ × P ₈	2.87	2.98	2.28
41	P ₇ × P ₉	2.75	1.51	1.96
42	P ₇ × P ₁₀	1.41	2.83	2.30
43	P ₈ × P ₉	2.67	2.79	2.15
44	P ₈ × P ₁₀	1.72	1.83	2.20
45	P ₉ × P ₁₀	1.87	1.94	1.63
F₁ Hybrid mean		1.91	2.02	1.91
Parents				
1	P ₁	1.99	2.10	2.05
2	P ₂	1.43	1.51	1.47
3	P ₃	1.51	1.55	1.53
4	P ₄	1.33	1.40	1.37
5	P ₅	1.65	1.73	1.69
6	P ₆	1.90	1.91	1.91
7	P ₇	2.50	2.47	2.49
8	P ₈	2.03	2.07	2.05
9	P ₉	2.67	2.56	2.61
10	P ₁₀	1.45	1.46	1.46
Parental mean		1.84	1.87	1.86
Grand mean		1.90	1.99	1.90
CV		6.24	4.71	4.93
CD 5%		0.19	0.15	0.16
Range	Lowest	1.15	1.26	1.35
	Highest	3.38	3.54	3.11