

Original Research Article

Effect of Gibberellic acid and Naphthalene acetic acid on growth and yield of Cucumber (*Cucumis sativus* L.) in Prayagraj agro climatic conditions

ABSTRACT

An experiment was conducted on effect of gibberellic acid and Naphthalene acetic acid to determine the best suitable concentrations on growth and yield of cucumber at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during mid-March 2021. Total Nine treatments were tried and replicated three times in a Randomized Block Design. Treatments include two growth regulators having concentrations of GA₃ (15, 25, 35, 45 ppm) and NAA (75,100,125,150 ppm) and Control (Water spray) were sprayed at two leaf stage and four-leaf stages. The application of plant growth regulators significantly affects the vegetative as well as reproductive attributes of the crop. Application of 25 ppm GA₃ at 2, 4 leaf stage was found most effective in terms of Vine length at 20, 40, 60 days (39.12, 91.02, 180.32 cm) Days to first appearance of male flowers in 50% plants (32.86 days) , No. of fruits per plant (12.20) Average fruit yield per plant (1870.37 g) fruit weight (153.33g) Days to first appearance of female flowers in 50% plants (38.93 in 35 ppm GA₃) , Days to first harvest (50.6 days in 35 ppm GA₃) , length of the fruit(14.66 cm in 15 ppm GA₃) ,fruit diameter (4.28 cm in 15 ppm GA₃), Average yield (18.77 t/ha in 25 ppm GA₃)

Keywords: Cucumber, GA₃, NAA, Growth, yield.

1. INTRODUCTION

Cucumber (*Cucumis sativus* L.) belongs to the family Cucurbitaceae with monoecious nature having 2n=14 as chromosome number is an important warm season vegetable crop (Bailey et al., 1969) It is cross pollinated and propagated by seed. *Cucumis hardwickii* is progenitor of present cultivating cucumber. The fruit of cucumber is said to have cooling effect, prevent constipation and check jaundice and indigestion. During 2020-2021 productivity of cucumber was about 1,18,000 ha with annual production of 1,665 mt was recorded (DAC & FW). In cucumber staminate flowers are much more than pistillate flowers so application of optimum concentration of PGR at correct stage of plant growth modifies the morphology and physiology of plant by inhibiting or by stimulating the enzymes which regulate the metabolism of plants in altering the sequence of flowering. Plant growth regulators can be used flexibly for the fine tuning of crops that grow uncontrollable and unpredictable environmental conditions (Rademacher, 2016). GA₃ and NAA are the two important plant growth regulators that alter the growth and yield contributing characters of cucumber. Exogenous application of GA₃ at three leaf stage and tendril initiation stage helps in enhancing vine length, branches, promoting and better fruit setting while NAA which is a synthetic form of auxin helps in enhancement in vegetative growth, fruit and seed yield (Nagamani 2015). Increasing length of main axis and number of branches (Hadvani,2010). Application of auxins, gibberellins, growth retardants shifts towards femaleness in sex expression (Chowdary et al 1959) Inadequate data was observed PGR's on growth, yield of cucumber. Therefore an experiment was conducted to determine the best concentration of gibberellic acid and naphthalene acetic acid on growth and yield of cucumber.

Comment [j1]: The experiment was conducted to determine the

Comment [j2]: Whats it means?

Comment [j3]: Do you only mentioned the GA₃ and whats hapen with NAA?
The comparisons was against GA₃ vs NAA, GA₃ vs Control or NAA vs Control?

Comment [j4]: Bibliography

Comment [j5]: What PGR means?

Comment [j6]: Bibliography?
What are the enzymes?

Comment [j7]: Who do it?

Comment [j8]: how do you related this with text?

Comment [j9]: Inadequate? Why?

2. MATERIAL AND METHODS

The field experiment was conducted at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during mid-March 2021. The University is about 7 km away from Prayagraj city and it is geographically situated at 25.5° N latitude, 81.08° E longitudes. The altitude of this location is about 98 meter above the mean sea level. The soil of the experimental field was sandy loam with pH of 6.8. Recommended package of cultural practices were adopted at last ploughing FYM- 2.0 kg/m² and half dose of nitrogen, full dose of P₂O₅ and K₂O were applied as basal and the remaining half nitrogen was applied at 30 days after sowing 100:60:60 kg/ha was applied in the form of Urea, Single super phosphate and Muriate of potash. The field experiment was carried out as a randomized block design with three replications. Two seeds per pit were sown with a spacing of 1.5 x 0.6 m in march 2021. GA₃ (15,25,35,45ppm) and NAA(75,100,125,150ppm) solutions, control (water spray) were prepared from the respective stock solutions with distilled water. The solutions were sprayed at 2 and 4 leaf stage with hand sprayer early in the morning. Irrigation ,weeding, hoeing and plant protection measures were employed as per the need of the crop. For each treatment of a replication five plants were selected and observations were recorded on various parameters of growth, and yield of the crop as per the planned schedule .The recorded data was analyzed statistically by ANOVA.

Comment [j10]: you should write it by sections. This way of writing is typical of results

3. RESULTS AND DISCUSSION

Germination parameters

The seeds were not treated with the growth regulators and the experimental results revealed that there is no significant difference with respect to days taken to germination it can be affected by various environmental factors such as the difference between day and night temperature, soil moisture, light, pH of the soil in cucumber. The statistically analysed data was presented in Table no.1.

Comment [j11]: These aren't results, you only mentioned it but didn't describe anything. The table 1 didn't describe if the significance was against the control and what treatment. The text didnt describe the results in table 1.

Growth parameters

Application of growth regulators has significant effect on vegetative growth as well as reproductive attributes (Mahala *et al.*, 2010). In the experiment the results revealed that the vine length increased significantly with the application of GA₃ 25 ppm (39.12,91.02,180.32 cm) which was at par with GA₃ 15 ppm (34.9,89.76,178.84 cm) followed by NAA 75 ppm (35.03,86.16,176.72 cm) and minimum was recorded (29.8,73.21,162.5 cm) under control at 20 ,40 ,60 days after sowing(Chovatia *et al.*,2010 ; Sure *et al.*,2012) GA₃ increases the rate of photosynthesis, efficiency of utilization of photosynthates which results in rapid cell elongation and cell division, stimulates RNA and protein synthesis which ensures in increasing the vine length. These finding are in the confirmity with earlier findings (Ajay *et al.*,2018). The data indicated in Table no 1: shows that days to first appearance of male flowers in 50 % plants of cucumber were significantly influenced by the various doses of NAA ,GA₃ as compared to control(Acharya *et al.*,2020). Minimum days taken to first appearance of male flowers in 50% plants were recorded in GA₃ 25ppm (32.86) which is at par with GA₃ 15ppm (33.4) where as maximum was recorded in control-water spray (39). This might be due to reducing the sugars there by inducing metabolic activity of cell which induces early flowering (Kadi *et al* 2018). Minimum days to first appearance of female flowers in 50% plants were recorded in GA₃ 35 ppm (38.93 days) which is par at GA₃ 25 ppm (39.9) where as maximum days under control-water spray(46.2). This might be due to effect of GA₃ which alters the sexual differentiation in potentially male buds to female buds that enhances female flower formation earlier. The similar results of the study are in agreement with were Rahman *et al.*,(2018) ;Dixit *et al.*,(2001) The statistically analysed data was presented in Table no. 1.

Comment [j12]: You are comparing the results of the GA3 through time, with the control, but did you do any statistics to probe it?

Comment [j13]: Did you do any experiment to comprove it or you readed in any paper?

Comment [j14]: What were these doses?

Comment [j15]: That's your result or the result of some one else?

Comment [j16]: Any statistics for it?

Table no. 1 Effect of Gibberellic acid and Naphthalene acetic acid on days to germination, vine length, Days to first appearance of male flowers in 50% plants and first appearance of female flowers in 50% plants.

Notation	Treatment	Days to germination	Vine length(cm)			Days to 1 st appearance of male Flowers in 50% plants	Days to 1 st appearance of female flowers in 50% plants
			20 DAS	40 DAS	60 DAS		
T ₁	Control	4	29.89	73.21	162.75	39.93	46.26
T ₂	GA ₃ 15 ppm	5	34.96	89.76	178.84	33.40	41.06
T ₃	GA ₃ 25 ppm	4	39.12	91.02	180.32	32.86	39.93
T ₄	GA ₃ 35 ppm	5	33.02	82.31	172.79	34.13	38.93
T ₅	GA ₃ 45 ppm	5	32.48	79.88	170.16	36.05	44.86
T ₆	NAA 75 ppm	4	35.03	86.16	176.72	37.06	45.80
T ₇	NAA100 ppm	5	34.14	85.13	175.94	33.86	42.13
T ₈	NAA125 ppm	5	33.28	84.15	173.79	35.26	43.73
T ₉	NAA150 ppm	4	30.79	77.80	166.75	44.33	44.33
F-Test		NS	S	S	S	S	S
SE(d) ±		0.150	0.812	1.231	1.072	0.285	0.579
CD _{0.05}		-	1.735	2.631	2.292	0.608	1.238

S-significant ; NS-Non significant; DAS-days after sowing




Fig 1: Effect of Gibberellic acid and Naphthalene acetic acid on vine length of cucumber




Fig 2: Effect of Gibberellic acid and Naphthalene acetic acid on Days to first appearance of male and female flowers in 50% of the plants and Harvesting of cucumber

Yield Parameters

In the study the application of gibberellic acid and naphthalene acetic acid shows significant difference on yield parameters as compared to control. The results obtained in the present study shows that minimum days taken to first harvest of cucumber was recorded in GA₃ 35 ppm (49.8) which was at par with GA₃ 25 ppm(50.60), where as the maximum days taken to first harvest in control-water spray (55.2).The treated plants are more physiologically active translocates food to develop fruits which results in early maturity of fruits.The maximum fruit weight (153.33g) was recorded in GA₃ 25 ppm followed by GA₃ 15 ppm (151.11g) where as minimum was recorded in control-water spray (142.90 g). GA₃ and NAA application increases the fruit mesocarp by increasing the cell size (Jyoti *et al.*,2016 ;Hossain *et al.*, 2006)).The length of the fruit(14.66 cm) and the diameter (4.42cm) was recorded highest with application of GA₃ 15ppm.This may be due to increasing the accumulation of carbohydrates and traslocation of food towards fruit. The results are similar with (kadi *et al.*,2018).The maximum number of Fruits per plant (12.2), Average yield per plant (1870.378 g) and Average yield tonnes per hectare (18.70 t/ha) was recorded in GA₃ 25 ppm where as minimum (8.13,1162.3 g,11.61 t/ha) was recorded in control-water spray respectively (Das *et al.*,2001;Anayat *et al.*,2020) and similar in Dalai *et al.*,2015) The Yield which depends on multiple factors like number of fruits per plant, fruit weight, length and diameter of the fruits. GA₃ has an effect on both vegetative as well as reproductive characters, mainly female flower formation and fruit setting which ultimately increases the number of fruits, fruit weight both combinedly increased fruit yield per plant resulted in maximum Average yield per hectare.This may be due to better enzymatic activity and the synthesis of growth regulators endogenously which was similar findings of (sure *et al.*,2012, Anayat *et al.*,2020) The statistically analysed data was presented in Table no. 2.

Table no. 2 Effect of Gibberellic acid and Naphthalene acetic acid on Days to first harvest, fruit weight, length of the fruit, fruit diameter, no. of fruits per plant, average yield per plant and average yield per hectare.

Notation	Treatment	Days to first harvest	Fruit weight (g)	Length of the fruit (cm)	Fruit diameter (cm)	No of fruits per plant	Average yield per plant (g)	Average yield (t/ha)
T ₁	Control	55.20	142.90	12.36	3.96	8.13	1,162.36	11.61
T ₂	GA ₃ 15 ppm	51.20	151.11	14.66	4.42	11.13	1,682.39	16.81
T ₃	GA ₃ 25 ppm	50.60	153.33	14.12	4.25	12.20	1,870.37	18.70
T ₄	GA ₃ 35 ppm	49.80	146.86	14.05	4.15	8.53	1,253.14	12.53
T ₅	GA ₃ 45 ppm	53.33	145.67	13.88	4.11	8.20	1,194.58	11.94
T ₆	NAA 75 ppm	53.86	149.72	14.23	4.28	10.73	1,606.73	16.06
T ₇	NAA100 ppm	51.80	148.01	13.74	4.07	9.66	1,430.71	14.30
T ₈	NAA125 ppm	52.33	147.79	14.19	4.26	9.26	1,369.56	13.69
T ₉	NAA150 ppm	52.60	143.90	13.54	4.02	8.20	1,180.03	11.79
F-TEST		S	S	S	S	S	S	S
SE(D) ±		0.513	1.217	0.184	0.037	0.190	29.083	0.291
CD _{0.05}		1.131	2.602	0.394	0.080	0.407	62.184	2.515




Fig 3: Effect of gibberellic acid and naphthalene acetic acid on fruit weight of cucumber




Fig 4: Effect of gibberellic acid and naphthalene acetic acid on length and diameter of cucumber




Fig 5: Effect of gibberellic acid and naphthalene acetic acid on no of fruits per plant of cucumber




Fig 6: Effect of gibberellic acid and naphthalene acetic acid on average yield of cucumber




Fig 7: Effect of gibberellic acid and naphthalene acetic acid on average yield per plant of cucumber

4. CONCLUSION

It is concluded that the effect of gibberellic acid and naphthalene acetic acid on growth and yield of cucumber at 2, 4 leaf stage was found superior in T₃ - GA₃ 25 ppm in terms of Vine length , Days to first appearance of female flowers in 50% plants, Days to first appearance of male flowers in 50% plants, Days to first Harvest, Fruit weight in grams (g) , Length of the fruit (cm), Fruit diameter (cm), Number of Fruits per plant, Average Yield per plant(g),Average Yield (t/ha) as compared to control.

REFERENCES

- Acharya, S. K., Thakar, C., Brahmabhatt, J. H., and Joshi, N. (2020).** Effect of plant growth regulators on cucurbits. *Asian Journal of Pharmacognosy and Phytochemistry*, 9(4), 540-544.
- Anayat, R., Mufti, S., Rashid, Z., Wani, S., and Khan, I. M. (2020).** Effect of Gibberellic Acid and Cycocel on Yield and Quality of Bitter Gourd. *International Journal of Pure and Applied Biosciences* 8, 4, 402-06.
- Bailey, L.H. (1969).** Manual of cultivated plants. Macmillan Company, New York. 1. 116 pp.
- Choudhury B, Phatak SC.** Sex egression and sex ratio in cucumber as affected by plant regulator sprays. *Indian Journal of Horticulture*. 1959; 16(3):162-169.
- Chovatia, R.S.; Ahlawat, T.R.; Kavathia, Y.A.; Jivani, L.L.; and Kaila, D.C. (2010).** Effect of plant growth regulators on vegetative, flowering and yield of bitter gourd cv. Priya. *Indian J. Hort.*, 67: 254-256.
- Dalai, S., Singh, M. K., Singh, K. V., Kumar, M., Malik, S., and Kumar, V. (2015).** Effect of foliar application of GA₃ and NAA on growth, flowering yield and yield attributes of cucumber [*Cucumis Sativus* L.]. *Annals of Horticulture*, 8(2), 181-194.
- Das, R.; Rabha, B.K.; and Dey, S.C.; (2001).** Effect of growth regulators on sex expression and fruit yield in cucumber (*Cucumis sativa* L.) under plastic greenhouse. *Annals of Agricultural Research.*, 22(3): 312-317.
- Dixit, A.; Rai, N.; and Kumar, V. (2001).** Effect of plant growth regulators on growth, earliness and sex ratio in Watermelon under Chhatisgarh region. *Indian Journal of Agricultural Research*, 35: 173-176.
- Farhana, U. (2015).** Effects of plant growth regulators on flowering behavior and yield of cucumber (Doctoral dissertation, Department of agricultural botany Sher-E-Bangla Agricultural University Dhaka-1207). *Journal of science and Technology*, 33-42
- Garg, P., Dev, R., Raj, S., Patel, V. J. K., and Singh, V. K. (2020).** Influence of plant growth regulators (PGRs) on growth parameters and sex ratio in cucumber (*Cucumis sativus* L.). *Journal of Pharmacognosy and Phytochemistry*, 9(3), 1658-1661.
- Hadvani VG.** Effect of plant growth regulators on growth, flowering, yield and quality of muskmelon (*Cucumis melo* Linn.) cv. Rasmadhuri. Thesis M.Sc (Hort.) Junagadh Agricultural University, 2010.
- Hossain, D.; Karin, M.A.; Pramani, M.H.R.; and Rehman, A.A.S. (2006).** Effect of foliar application of IAA and GA₃ on sex expression, yield attributes and yield of bitter gourd (*Momordica charantia* L.). The Chittagong University. *Journal of Biological Sciences.*, 5: 55-62.
- Jyoti, S., Patel, N. B., and Patel, J. B. (2016).** Effect of growth regulators and stages of spray on seed yield and seed quality parameters of ridge gourd [*Luffa acutangula* (Roxb) L.]. *Journal of Applied and Natural Science*, 8(3), 1551-1555.
- Kadi, A. S., Asati, K. P., Barche, S., and Tulasigeri, R. G. (2018).** Effect of different plant growth regulators on growth, yield and quality parameters in cucumber (*Cucumis sativus* L.) under

polyhouse condition. *International Journal of Current Microbiology and Applied Sciences*, 7(4), 3339-3352.

Mahala P, Choudhary MR, Yadav TV, Garhwaland OP, Singh P. Effect of plant growth regulators on yield, quality and economics of bottle gourd (*Lagenaria siceraria* (Mol.) Standl.). *Annals of Agri-Bio Research*. 2014; 19(1):137-139.

M.A. Rahman, S. Sikder, M.M. Bahadur and S.K. Pramanik (2018). Influence of gibberellic acid on growth, flowering, and fruit yield of cucumber *Journal of Science and Technology*: 33-42

MR, S., Helmy, Y. I., Ahmed, A. A., and Ghoname, A. A. (2016). Effect of foliar application of growth regulators (GA3 and Ethereal) on growth, sex expression and yield of summer squash plants (*Cucurbita pepo* L.) under plastic house condition. *International Journal of ChemTech Research*, 9(6), pp 70-7.

Nagamani S, Sudipta B, Lal SK, Behera TK, Chakrabarty SK, Talukdar A. Effect of plant growth regulators on sex expression, fruit setting, seed yield and quality in the parental lines for hybrid seed production on bitter gourd. (*Momordica charantia* L.). *Indian Journal of Agricultural sciences*. 2015; 85(9):1185-1191

Pandey, P., Shukla, I. N., and Upadhyay, A. (2021). Effect of different plant growth regulators on growth, yield and quality parameters of cucumber (*Cucumis sativus* L.) cv. Kalyanpur green. *Journal of Pharmacognosy and Phytochemistry*, 10(1), 2681-2684.

Pradeep Garg, Rishabh Dev, Shani Raj, Vijay Jitendra Kumar Patel and VK Singh (2018). Influence of plant growth regulators (PGRs) on growth parameters and sex ratio in cucumber (*Cucumis sativus* L.) *Journal of Pharmacognosy and Phytochemistry*, 9(3): 1658-1661

Rademacher W. Chemical regulators of gibberellin status and their application in plant production. *Annual Plant Reviews*. 2016; 49:359-403.

Sandra, N., Sudipta, B., Sukhbir, S., Lal, S. K., Behera, T. K., Chakrabarty, S. K., and Talukdar, A. (2015). Effect of plant growth regulators on sex expression, fruit setting, seed yield and quality in the parental lines for hybrid seed production in bitter gourd (*Momordica charantia*). *Indian Journal of Agriculture Sciences*, 85, 1185-191.

Sure, S.; Arooie, H.; and Azizi, M. (2012). Influence of plant growth regulators (PGRs) and planting method on growth and yield in Oil pumpkin (*Cucurbitapepo* var. *styriaca*). *Not. Sci. Biol.* 4(2): 101-107.

Tantasawat, P. A., Sorntip, A., and Pornbungkerd, P. (2015). Effects of exogenous application of plant growth regulators on growth, yield, and in vitro gynogenesis in cucumber. *HortScience*, 50(3), 374-382.

Thappa, M., Kumar, S., and Rafiq, R. (2011). Influence of plant growth regulators on morphological, floral and yield traits of cucumber (*Cucumis sativus* L.). *Agriculture and Natural Resources*, 45(2), 177-188.

Vyas, M. N., Leua, H. N., Jadav, R. G., Patel, H. C., Patel, A. D., and Patel, A. S. (2015). Effect of plant growth regulators on growth, flowering and yield of ridge gourd (*Luffa acutangula* roxb l.) Cv. Pusa nasdar. *Ecology environment and conservation journal*, 902