

Integrated Zinc and Iron management practices for enhancing productivity and flower quality of Marigold

ABSTRACT:

The present research was conducted to examine the influence of iron and zinc both as soil application and foliar spray for better production and flower quality of marigold. The research was performed at the Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore during 2021-2022. The experiment was laid out in completely randomized design (CRD) with eight treatments and three replications. The eight treatments included an untreated control, 100% RDF, soil application of enriched FYM with zinc sulphate and iron sulphate at different proportions, foliar spray of zinc sulphate and iron sulphate at different ratios and TNAU prepared liquid multi micronutrient. The results of the study indicated that foliar spray of zinc and iron resulted in superior performance of the crop compared to soil application. Among all the treatments foliar application of 0.5% ZnSO₄+ 0.5% FeSO₄ at two stages (20DAT and 35DAT) along with RDF produced significantly enhanced vegetative growth, quality of flower. Foliar application of 1.0 % liquid multi micronutrient along with RDF also produced significant effect on xanthophyll content compared to control. However both soil application and foliar spray of zinc and iron showing significant result compared to control for better productivity and flower quality of marigold.

Keywords: Zinc, Iron, Multi micronutrient, Enriched FYM, Marigold, Xanthophyll

1. INTRODUCTION

Marigold (*Tagetes erecta*) belonging to the family Asteraceae is one of the important annual commercial flower crops cultivated around the world. It is widely cultivated owing to its ease of growing. It is grown for loose flower and cut flower, garland making and garden display. It stands in second place next to Chrysanthemum among the annual flowers. The crop is also known for its potential to control root knot nematodes. The flowers are an important source of carotenoids [2]. It covers an area of 66.13 thousand hectares with 603.18 thousand metric tonnes of production in 2015-16 [3]. In India Andhra Pradesh and Tamil Nadu are major producers of marigold flower. India's maximum producer and the main trading centre for marigold is Erode, a city of Tamil Nadu. It covers an area of 2761 ha with 72.389 thousand metric tonnes of production and 26.22 metric tonnes productivity per hectare. (TANHODA, Chennai, 2019)

In Indian agriculture, the relevance of micronutrients is well understood as their application has greatly enhanced the production of a variety of crops. Micronutrients are as important as macronutrients in terms of plant development, yield, and quality. In the past, there was no need to supplement micronutrients in soil because these were naturally present. However, in most soils, due to intensive farming, increased salinity, and soil pH, these nutrients are not available to plants [1]. Micronutrient application has not only contributed in enhancing the food grain production but also helped in sustaining soil health. In India, micronutrient status is normally low. It has been estimated that the deficiency levels of available nutrients viz., Fe, Zn, Mn, Cu and B in Tamil Nadu are 12.62%, 63.30%, 7.37%, 12.01% and 20.65% respectively. Among all micronutrients, deficiency of zinc is commonly prevalent in many of the districts in Tamil Nadu [4].

Iron plays a significant role in various physiological and biochemical pathways in plants. It serves as an element of many vital enzymes such as cytochromes of the electron transport chain, and it is thus required for a wide range of biological functions. In plants, iron is involved in the synthesis of chlorophyll, and it is essential for the maintenance of chloroplast structure and function [8]. Fe is needed for the synthesis of enzymes that

produce chlorophyll it is used for catalytic activity, other metabolic activities like biological oxidation reduction in plants, oxidative photophosphorylation during cell respiration. Iron plays major role in the metabolism of carbohydrates and for the protein synthesis [12].

Zn is an important element for growth of plant as well as for other metabolic functions namely enzymatic activities, protein synthesis (regulatory cofactor), photosynthesis, synthesis of auxin, cell division, construction of membrane and sexual fertilization. Next to iron (Fe), zinc (Zn) is the second most abundant microelement in organisms, and it plays a vital role in a variety of biological activities [8]. Plants require zinc as a critical micronutrient because it is a component of many enzymes and proteins in organisms; it is an influential metal for appropriate growth and development in plants [24].

This study aimed to investigate the effect of varying quantities of iron and zinc in combination, when applied as soil with dissolved irrigation water and applied as a foliar spray, affecting the growth, blooming, and chemical composition of marigold under shade net house condition.

2. MATERIALS AND METHODS

The experiment was conducted during 2021-2022 at the Radioisotope (Tracer) Laboratory, Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore. Coimbatore located in the North Western agro climatic zone of Tamil Nadu at (11 °N latitude, 77 °E longitudes and at an altitude of 426 meters above MSL). The mean annual maximum and minimum temperature are 31.5°C and 21.2 °C, respectively. The mean annual rainfall is 657mm.

Shade net house was constructed with 50% green shade net. It provided a partially controlled climate and environment for the marigold crop by decreasing light intensity and effective heat during the day.

The pot experiment was conducted with marigold cv. Local Orange. The pot soil was clay loam in texture, slightly alkaline in reaction (pH of 7.52), medium in organic carbon (0.54 %) with low in available nitrogen (191.0 kg ha⁻¹), medium in available phosphorous (19.0 kg ha⁻¹) and high in available potassium (297.0 kg ha⁻¹), sufficient in both DTPA Mn (2.08ppm) and DTPA Cu (2.20ppm) but deficient in DTPA Zn (0.74ppm) and DTPA Fe (1.32ppm). The investigation included eight treatment combinations (listed out in Table) replicated three times, was laid out in completely randomized design.

2.1 Chart 1: Treatment details:

Notation	Treatment details
T ₁	Absolute Control
T ₂	100% RDF (90:90:75 NPK kg ha ⁻¹)
T ₃	RDF + soil application of FYM enriched with ZnSO ₄ @10 kg ha ⁻¹ and FeSO ₄ @15 kg ha ⁻¹
T ₄	RDF +soil application of FYM enriched with ZnSO ₄ @15 kg ha ⁻¹ and FeSO ₄ @20 kg ha ⁻¹
T ₅	RDF +soil application of FYM enriched with ZnSO ₄ @20 kg ha ⁻¹ and FeSO ₄ @25 kg ha ⁻¹
T ₆	RDF + 0.5% ZnSO ₄ (foliar spray) + 0.5% FeSO ₄ (foliar spray)
T ₇	RDF + 0.5%ZnSO ₄ (foliar spray) + 1.0%FeSO ₄ (foliar spray)

T ₈	RDF + 1.0% liquid multi micronutrient (foliar spray)
----------------	------------------------------------------------------

2.2 Zinc and Fe enriched FYM

Farmyard manure (FYM) was collected from the farm at the Tamil Nadu Agricultural University, Coimbatore and used for application. ZnSO₄ and FeSO₄ were mixed with FYM in 1:10 ratio as per the treatments. The mixture was kept in polythene bags for incubation under anaerobic condition for 21 days. The mixture was turned over periodically and the moisture content was checked twice in a week.

2.3 Crop husbandry

Earthen pots of height 25cm with inner diameter of 25cm were used for the experiment. Each pot was filled with 10kg of processed soil and the soil was watered and made to a fine tilth. Before transplanting, the soil was fertilized as per the blanket recommendation viz., 90:90:75 kg N: P₂O₅: K₂O ha⁻¹. Full dose of the recommended phosphorus and potassium were applied as basal using single super phosphate and muriate of potash respectively as source of fertilizers. Half of the recommended dose of nitrogen was applied basally and the remaining half was applied as top dressing after transplanting. Enriched FYM was mixed with RDF and applied before transplanting. At the time of transplanting, a thin film of water was maintained in the pot. Twenty one days old seedlings were transplanted in the pot with three seedlings in each pot.

2.4 Preparation of micronutrients solution

Fresh stock solution of micronutrients with a known concentration was prepared. The micronutrients were first dissolved in a small amount of water, and then combined with water to create a stock solution. The needed concentrations of solution were then generated by dilution from the stock solution.

Liquid multi micronutrient developed by the Department of Soil Science and Agricultural Chemistry, TNAU, Coimbatore was used. It contains Zn (0.3567%), Fe (0.9651%), Mn (0.1456%) and Cu (1.1345%). ZnSO₄, FeSO₄ and multi micronutrient mixture were applied as foliar spray at two stages namely (20 DAT and 35 DAT).

The observations recorded included vegetative growth parameters namely plant height, number of branches, fresh weight of plant, dry weight of plant, flower yield and quality parameters namely diameter of flower, individual flower weight, shelf life of flower and xanthophyll content in the flowers.

2.5 Biochemical traits

Total Xanthophyll content was estimated by AOAC method (Lawrence, 1990) [19].

2.6 Statistical analysis

The data obtained from the experiment were subjected to statistical analysis using AGRSS software version 7.01. The level of significance used was $p < 0.05$. Critical difference (CD) values were calculated for the $p < 0.05$ whenever "F" test was found significant [11].

3. RESULTS AND DISCUSSION

3.1 Growth parameters

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum plant height (81.6 cm at 60 DAT) which was followed by T₈ (81.32cm), where RDF was applied along with FS of 1.0% liquid multi micronutrient. The greater plant height in T₆ can be attributed to the foliar application of Zn and Fe which might have increased the cell division, cell elongation and other growth promoting hormones which was ultimately reflected as taller plants in the said treatment. Similar results were obtained earlier [9], [10], [27].

The highest number of branches (18.33 at 60 DAT) was recorded in T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) followed by T₈ (17.00), whereas control showed minimum number of branches (9.67). Such a significant increase in number of branches is due to foliar application of zinc and iron which might have increased vegetative growth parameters of plant by involving in tryptophan and chlorophyll synthesis, which in turn helps to maintain polarity and apical dominance of plant. Similar results were obtained by earlier workers [8], [26].

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum fresh weight of plant (359.86g) which was followed by T₈ (350.05g) whereas T₁ (control) recorded minimum (250.11g). The treatment T₆ recorded the maximum dry weight of plant (78.08g) which was followed by T₈ (77.20g) while T₁ (control) recorded minimum (50.11g). This is due to application of zinc and iron which play major role in production of vegetative growth and ultimately encourage the biomass of plant, resulting in increased fresh and dry weight of plants. [6], [25], [26].

Table 1: Effect of Zn and Fe on growth parameters of marigold cv. Local Orange

Treatment	Plant height (cm)			No. of primary branches per plant			Fresh weight of plants (g)			Dry weight of plants (g)		
	30 DAT	45 DAT	60 DAT	30 DAT	45 DAT	60 DAT	30 DAT	45 DAT	60 DAT	30 DAT	45 DAT	60 DAT
T ₁	25.80	32.94	42.92	5.00	8.00	9.67	170.05	210.06	250.11	26.10	38.11	50.11
T ₂	26.32	33.44	43.42	7.00	10.00	11.67	189.79	240.87	272.71	41.06	53.07	65.07
T ₃	28.55	35.64	47.64	8.00	11.00	13.67	205.82	256.89	315.81	46.04	58.05	70.05
T ₄	29.22	36.32	48.32	10.00	13.00	15.33	210.68	261.75	327.67	48.06	60.07	72.07
T ₅	29.98	37.08	49.07	11.00	14.00	16.67	216.18	267.25	340.11	50.53	62.54	74.54
T ₆	34.39	41.50	53.49	13.00	16.00	18.33	247.95	299.00	359.86	54.07	66.08	78.08
T ₇	31.37	38.56	50.64	11.00	14.00	15.67	226.15	277.22	343.15	51.65	63.66	75.66
T ₈	34.10	41.24	53.25	12.00	15.00	17.00	245.89	296.94	350.05	53.19	65.20	77.20
SEd	0.12	0.09	0.07	0.82	0.04	0.53	0.83	0.78	0.08	1.16	0.57	0.57
CD (0.05)	0.25	0.19	0.14	0.09	1.12	1.12	1.76	1.65	0.16	2.46	1.21	1.21

3.2 Quality parameters:

Flowering, quality attributes namely individual flower weight, flower diameter, shelf life of flower were significantly influenced by Zn and Fe (Table 2).

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum individual diameter of flower (6.76cm) which was followed by T₈ (6.62cm), where RDF was applied along with FS of 1.0% liquid multi micronutrient. The treatment T₁ (control) recorded the minimum (5.62cm) diameter of flower. This could be due to the fact that iron and zinc work together to regulate semi-permeability of cell walls that allowed enormous water to enter into flowers and also help for increasing iron synthesis, which promotes cell expansion in turn resulting in increased flower diameter [14], [15].

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum fresh flower weight (4.69g) which was followed by T₈ (4.65g), where RDF was applied along with FS of 1.0% liquid multi micronutrient. The treatment T₁ (control) recorded the minimum fresh flower weight (3.67g). Both zinc and iron enhance biomass of the plant which ultimately resulted in maximum fresh and dry weight of flower [16], [17], [20].

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum dry weight of flower (1.05g) which was followed by T₈ (1.03g), where RDF was applied along with FS of 1.0% liquid multi

micronutrient. The treatment T₁ (control) recorded the minimum dry weight of flower (0.75g). The results revealed application of zinc and iron might have accumulate dry matter alongside with increased availability and uptake of micronutrients which ultimately increased dry flower weight.[18], [5].

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) showed the maximum shelf life of flower (5.20 days) which was followed by T₈ (5.05 days), where RDF was applied along with FS of 1.0% liquid multi micronutrient. The treatment T₁ (control) recorded the minimum (2.50 days). The longest shelf life of flower in days observed in T₆ due to the foliar application of Zn and Fe because they play an important role in physiology *viz.*, carbohydrate metabolism, plant growth regulation and enzyme synthesis which results in good flower quality. Further this creates favourable condition inside the stem and gives strength against adverse condition after harvest. Similar results were reported by earlier [15], [21], [22].

3.3 Yield attributes:

The flower yield attributes namely number of flowers per plant, number of flowers per pot and flower yield were also significantly influenced by application of zinc and iron (Table 2).

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum number of flowers plant⁻¹ (15.67) which was followed by T₈ (14.67), where RDF was applied along with FS of 1.0% liquid multi micronutrient. The treatment T₁ (control) recorded the minimum number of flowers plant⁻¹ (6.33). Application of iron and zinc either through foliar spray or soil application might have increased vegetative growth and this in turn might have led to production of more food material resulting in better development enabled by enhanced RNA metabolism, formation of DNA, synthesis of proteins, formation of pollen [23], [24], [25] ultimately leading and increase in number of flowers per plant.

The treatment T₆ (RDF + 0.5% ZnSO₄ FS + 0.5% FeSO₄ FS) recorded the maximum flower yield plant⁻¹ (73.48g) which was followed by T₈ (68.20g), where RDF was applied along with FS of 1.0% liquid multi micronutrient whereas the treatment T₁ (control) recorded the minimum flower yield plant⁻¹ (23.24g). This results showed that application of zinc and iron increased the synthesis of chlorophyll, growth promoting substances and mobility of minerals, water, photosynthesis and amino acids from source to sink which might have increased the flower production and ultimately flower yield. Similar results were also obtained by earlier workers [5], [23], [24].

Table 2: Effect of Zn and Fe on quality, yield and biochemical parameter of marigold cv. Local Orange

Treatment	Diameter of flower (cm)	Fresh weight of flower (g)	Dry weight of flower (g)	No. of flowers per plant	Number of flowers per pot	Yield of flowers per plant (g/plant)	Yield of flowers per pot (g/pot)	Shelf life of flowers (days)	Xanthophyll content (mg/g)
T ₁	5.62	3.67	0.78	6.33	19.33	23.24	70.95	3.50	1.30
T ₂	5.82	3.80	0.83	8.67	25.67	32.93	97.53	3.90	1.36
T ₃	5.90	4.34	0.90	11.00	32.67	47.74	141.77	4.30	1.45
T ₄	5.95	4.40	0.95	11.33	34.00	49.87	149.60	4.51	1.49
T ₅	6.01	4.50	0.97	12.67	38.33	57.00	172.50	4.49	1.55

T ₆	6.76	4.69	1.05	15.67	46.33	73.48	217.30	5.20	1.67
T ₇	6.21	4.62	0.98	13.67	41.00	63.14	189.42	4.53	1.65
T ₈	6.62	4.65	1.03	14.67	44.00	68.20	204.60	5.06	1.73
SEd	0.51	0.10	0.01	0.53	1.28	2.46	6.79	0.09	0.03
CD (0.05)	1.08	0.22	0.02	1.12	2.71	14.40	14.40	0.20	0.06

3.4. Biochemical parameter:

The treatment T₈ (RDF + 1.0% liquid multi micronutrient) recorded the maximum xanthophyll content in flower (1.73mg/g) which was followed by T₆ (1.67mg), where (RDF+0.5% ZnSO₄ FS + 0.5% FeSO₄ FS). The treatment T₁ (control) recorded the minimum xanthophyll content (1.30mg/g). This could be due to fact that iron plays a vital role in chlorophyll biosynthesis and zinc is favourable for influencing photosynthetic pigments, application of Zn and Fe might have helped in improving the photosynthetic electron transfer and photosynthesis rates leading to increased xanthophyll content in flowers. Similar results were reported earlier [5], [6], [27],[28].




Fig. 1. Effect of zinc and iron on xanthophyll content of marigold cv. Local Orange

4. CONCLUSION

The importance of Zn and Fe in the cultivation of marigold could be well demonstrated in the present experiment. The results indicated that marigold cv. Local Orange responded better to foliar application of ZnSO₄ @ 0.5% + FeSO₄ @ 0.5% at two stages namely 20 and 35 days after transplanting. These micronutrients improved the vegetative growth attributes, flower yield and quality. Application of 1.0% liquid multi micronutrient which contains zinc and iron along with RDF significantly enhanced the biochemical trait namely xanthophyll content (mg/g) of flowers.

REFERENCES

1. Ahmad, Iftikhar, M. Aslam Khan, M. Qasim, Rashid Ahmad, and Mahmood A. Randhawa. "Growth, yield and quality of *Rosa hybrida* L. as influenced by various micronutrients." *Pak. J. Agric. Sci* 47, no. 1 (2010): 5-12.
2. Pawar, Ajit, D. Saraladevi, M. Kannan, and A. SabirAhamed. "Effect of micronutrients and plant growth regulators on growth, flowering and yield attributes of marigold (*Tagetes erecta* L.)." *Madras Agricultural Journal* 106, no. march (1-3) (2019): 1.
3. Anonymous (2017). Horticultural statistics at a glance 2017. Horticulture statistics Division, Department of Agriculture, Cooperation and Farmers Welfare, Government of India, 481p.
4. Shukla, Arvind K., Sanjib K. Behera, Abhijit Pakhre, and S. K. Chaudhari. "Micronutrients in soils, plants, animals and humans." *Indian Journal of Fertilisers* 14, no. 3 (2018): 30-54.
5. Balakrishnan, V., M. Jawaharlal, T. Senthil Kumar, and M. Ganga. "Response of micro-nutrients on flowering, yield and xanthophyll content in African marigold (*Tagetes erecta* Linn.)." *Journal of Ornamental Horticulture* 10, no. 3 (2007): 153-156.
6. Subhash, Bishnoi, N. D. Polara, and A. L. Regar. "Response of micro-nutrients on flowering, yield, quality and xanthophyll yield in African marigold (*Tagetes erecta* Linn.)." *Trends in Biosciences* 10, no. 2 (2017): 626-628..
- 7. Broadley, Martin R., Philip J. White, John P. Hammond, Ivan Zelko, and Alexander Lux. "Zinc in plants." *New phytologist* 173, no. 4 (2007): 677-702.**
8. Neha, Chopde, M. J. Patokar, A. C. Bhaskarwar, and Patil Siddhi. "Response of marigold to foliar application of zinc and iron." *Trends in Biosciences* 10, no. 2 (2017): 892-894.
9. Choudhary, A., A. Mishra, P. K. Bola, S. K. Moond, and M. Dhayal. "Effect of foliar application of zinc and salicylic acid on growth, flowering and chemical constitute of African marigold cv. pusa narangi gainda (*Tagetes erecta* L.)." *Journal of Applied and Natural Science* 8, no. 3 (2016): 1467-1470.
10. Choudhary, A., A. Mishra, K. K. Nagar, and R. MEENA. "Foliar application of zinc and salicylic acid on African marigold." *Journal of Crop and Weed* 12, no. 3 (2016): 107-111.
11. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John wiley & sons;1984
12. Rout, Gyana R., and Sunita Sahoo. "Role of iron in plant growth and metabolism." *Reviews in Agricultural Science* 3 (2015): 1-24.
- 13. Jagtap, H. D., V. J. Golliwari, and S. A. Thakre. "Effect of foliar application of micronutrients on growth and flowering of rose under polyhouse conditions." *Asian Journal of Horticulture* 7, no. 1 (2012): 25-27.**
14. Jat, R. N., S. K. Khandelwal, and K. N. Gupta. "Effect of foliar application of urea and zinc sulphate on growth and flowering parameters in African marigold (*Tagetes erecta* Linn.)." *Journal of Ornamental Horticulture* 10, no. 4 (2007): 271-273.
15. Kakade, D. K., S. G. Rajput, and K. I. Joshi. "Effect of foliar application of Fe and Zn on growth, flowering and yield of China aster (*Callistephus chinensis* L. Nees)." *Asian Journal of Horticulture* 4, no. 1 (2009): 138-140.

16. Karuppaiah, P. "Effect of zinc and iron on growth, yield and quality of chrysanthemum (*Dendratherium grandiflorum* Tzeuleu)." *Asian Journal of Horticulture* 9, no. 1 (2014): 232-236.
17. Kumar, S., and K. Haripriya. "Effect of foliar application of iron and zinc on growth flowering and yield of Nerium (*Nerium odorum* L.)." *Plant Archives* 10, no. 2 (2010): 637-640.
18. Kumar, Prabhat, and J. S. Arora. "Effect of micronutrients on gladiolus." *Journal of Ornamental Horticulture* 3, no. 2 (2000): 91-93.
19. Lawrence, J. "Determination of total xanthophyll and marigold oleoresin." *Journal of Association of Official Analytical Chemists* 2 (1990): 970-975.
20. Jain, P. A. L. A. K. "Effect of micronutrients on growth and flowering of tuberose (*Polianthes tuberosa* L.)." PhD diss., M. Sc. Thesis, College of Horticulture, Mandasaur, India, 2014.
21. Surya, Narayan. "Effect of foliar application of NAA and micro-elements on vigour and flowering of marigold (*Tagetes erecta* L.) cv. Pusa Basanti." *HortFlora Research Spectrum* 4, no. 3 (2015): 264-267.
22. Pal, Sunder, A. V. Barad, A. K. Singh, B. S. Khadda, and Daleep Kumar. "Effect of foliar application of Fe and Zn on growth, flowering and yield of gerbera (*Gerbera jamesonii*) under protected condition." *Indian Journal of Agricultural Sciences* 86, no. 3 (2016): 394-8.
23. Bhute, P., D. Panchbhai, V. Raut, Chopde Neha, and Khobragade Hemlata. "Studies on flower production in annual chrysanthemum in response to iron and zinc." *Plant Arch* 17 (2017): 1017-1019.
24. Shah, Syed Tanveer, Sami Ullah, Nadeem Khan, Muhammad Sajid, Abdur Rab, Noor UI Amin, Asif Iqbal et al. "Effect of zinc as a foliar spray on growth and flower production of Marigold (*Tagetes erecta* L.)." *Pure and Applied Biology* 5, no. 4 (2016): 1.
25. Hembrom, Raimani, and Anil K. Singh. "Effect of iron and zinc on growth, flowering and bulb yield in liliium." *International Journal of Agriculture, Environment and Biotechnology* 8, no. 1 (2015): 61.
26. Saeed, Tariq, Imran Hassan, Ghulam Jilani, and Nadeem Akhtar Abbasi. "Zinc augments the growth and floral attributes of gladiolus, and alleviates oxidative stress in cut flowers." *Scientia Horticulturae* 164 (2013): 124-129.
27. Thirumalmurugan, V., K. Manivannan, and S. Nanthakumar. "Influence of micronutrients on growth, flowering and yield of African marigold (*Tagetes erecta* L.)." *Journal of Pharmacognosy and Phytochemistry* 10, no. 3 (2021): 461-463.
28. Yadegari, M. "Foliar application of Fe, Cu, Mn and B on growth, yield, and essential oil yield of marigold (*Calendula officinalis*)." *Journal of Applied Science and Agriculture* 8, no. 5 (2013): 559-567.