

Original Research Article

This manuscript needs much :[1MM]Comment efforts and data to be acceptable for publication in any journal

Effects of plant growth regulators GA₃ and NAA on growth and yield and quality of brinjal (*Solanum melongena* L.) cv. Kashi Sandesh

ABSTRACT

A field experiment was conducted to find out the Effects of ~~plant growth regulator~~ two concentrations of GA₃ and NAA and their ~~on combinations on~~ growth and yield and quality of brinjal (*Solanum melongena* L.) cv. Kashi Sandesh. The experiment was carried out at the Horticultural Research Farm, Department of Horticulture, Udai Pratap (Autonomous) College, Varanasi during the year 2020-21. The experiment was laid out in Randomized Block Design with three replications and nine treatments. ~~There were two plant growth regulators viz. GA₃ and NAA was used with different concentrations and applied by foliar application to assess the effect of growth, yield and quality of brinjal.~~ Excellent promising results was found in Treatment T₈ - NAA + GA₃ (40 ppm + 50 ppm) for all the characters viz. maximum plant height (71.14 cm), Number of leaves per plant (71.65), Number of branches (12.77), Days to 50% flowering (43.12), Number of fruit per plant (16.17), Fruit length (11.66 cm), Fruit weight (180.48 g), Fruit diameter (9.48 cm), Fruit yield (2.91 kg/plant), Fruit yield (29.22 kg/plot), Fruit yield (383.95 q/ha), Net return (421750.86 Rs./ha) and B:C ratio (2.73) which was followed by treatment T₇ and T₆ and in other hand T₉ - Control was gives poor performance among all the treatments. Hence considering the positive effect of NAA 40 ppm + GA₃ 50 ppm was found to be beneficial for brinjal growing farmers under Varanasi (U.P.) condition.

Very poor???? :[2MM]Comment

Key Words- Brinjal, PGRs, Growth, ~~Yield and Economics.~~

Add new strong key words :[3MM]Comment

INTRODUCTION

Brinjal (*Solanum melongena* L.) is one of the most common, popular and principal vegetable crops grown in India and other parts of the world. The brinjal is grown extensively in India and other Asian countries like Bangladesh, Pakistan, and Philippines. Major brinjal producing countries are China, Turkey, Japan, Egypt, Indonesia, Iraq, Italy, Syria and Spain. The cultivated brinjal of Indian origin has been in cultivation for long time (Thompson and Kelly, 1957). In India, it is one of the most common and popular vegetable crop grown throughout the country, except the higher altitude. The production of brinjal is governed not only by the inherent genetic yield potential of the cultivars but it is greatly influenced by several environmental factors and cultivation practices.

Brinjal fruits are fairly good source of Ca, P, Fe, and vitamins particularly 'B' group (Kiran *et al.* 2010). Brinjal is also valued for its medicinal properties and role of brinjal in treatment of liver disease, cough due to allergy, rheumatism, leucorrhoea and intestinal worms has been mentioned (Das and Barua; 2013). Like other vegetables, it provides dietary fiber, minerals, vitamins, carbohydrate and protein. Plant growth regulators are known to influence on higher yields as well as quality in horticultural crops. Recently, there has been global realization of the important role of PGR's in increasing crop yield. GA₃ is an important growth regulator that may have many uses to modify the growth, yield and yield contributing characters of plant (Rafeekheret *al.* 2002). Plant growth regulators are used widely to improve plant performance. GA₃ is one of those growth regulators that have positive effect on plant growth through the effect on cell division and elongation (Batlanget *al.*, 2006). It recorded dipping of brinjal seedling roots in NAA at 0.1 or 0.2 ppm for 24 hours influenced growth and development (Vajaet *al.* 2017). The advantage of plant growth regulators like Gibberellic (GA₃), Indol Acetic Acid (IAA), Nephthaline Acetic Acid (NAA), 2-4- Diclorophenoxy Acetic Acid (2,4-D) can be taken to increase the yield of local variety of brinjal.

Very poor introduction :[4MM]Comment

MATERIAL AND METHODS

The present experiment was carried out in research farm of Department of Horticulture at udaipratap (autonomous) college, Varanasi under agro- climatic conditions of Eastern region of Uttar Pradesh during the *rabi* season 2020-21 on brinjal variety Kashi Sandesh (Round). Seedling were uprooted from the nursery beds and transplanted in the **main plots** at a spacing of 70 cm row to row and 60 cm plant to plant. Transplanting was done in the **evening** hours immediately followed by irrigation for proper establishment of the seedlings. All the **recommended package of practices** was followed to raise a healthy crop. The experiment consisted of two levels of gibberellic acid (GA₃), two levels of naphthalene acetic acid (NAA) and their combinations was arranged in randomized block design with three replications and nine treatments *viz.* (T₁) - NAA (20 ppm), (T₂) - NAA (40 ppm), (T₃) - GA₃ (25 ppm), (T₄) - GA₃ (50 ppm), (T₅) - NAA + GA₃ (20 ppm + 25 ppm), (T₆) - NAA + GA₃ (20 ppm + 50 ppm), (T₇) - NAA + GA₃ (40 ppm + 25 ppm), (T₈) - NAA + GA₃(40 ppm + 50 ppm) and (T₉) - Control.

The required weight of the PGRs was taken using electronic sensitive balance and solution was prepared by dissolving in 1 mg L⁻¹. The solution was poured into hand-held sprayer and was directly sprayed on the plants three times at 30, 45 and 60 days after transplanting. Spraying was performed early in the morning to avoid rapid drying of the spray solution, due to transpiration. Data were collected from randomly selected five plants in each row. The collected data includes Plant heights (cm), No. of **leafleaves**/plant, No. of branches **per** /plant, **Days to** 50% flowering, No. of fruit **per** /plant, **Fruit length**(cm), Weight offruits (g), Fruit diameter(cm), **Fruit Yield**(Kg/plant), **Yield**(kg/plot) and **Yield**(q/ha). The data **was** **were** analyzed using analysis of variance (ANOVA) by **software** and **mean separation** was carried out at 5% probability level.

You are testing a round fruit :[5MM]Comment
shape variety ????

All of these give the same :[6MM]Comment
results

Which software you used??? :[7MM]Comment

????? :[8MM]Comment

RESULTS AND DISCUSSIONS

The data was present in various attributes showed that table 1 at 30 DAT maximum plant height was found in T₈- NAA + GA₃ (40 ppm + 50 ppm) (41.92) followed by T₇ - NAA + GA₃ (40 ppm + 25 ppm) (36.18) and minimum was found in T₉ - Control (27.18), at 45 DAT maximum plant height in T₈- NAA + GA₃ (40 ppm + 50 ppm) (61.20) followed by (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (57.84) and minimum was found in T₉-Control (37.75). At 60 DAT maximum plant height was found in T₈- NAA + GA₃ (40 ppm + 50 ppm) (71.14) followed by T₇- NAA + GA₃ (40 ppm + 25 ppm) (69.40) and minimum was found in T₉- Control (45.47). Gibberellin can promotes the activity of xyloglucan endotransglycosylase (XET) which cause loosening of cell wall and increase cell permeability (Saptari and Dawi 2013). Similar result was found in brinjal by Meena and Dhaka (2003). The no. of leaves per plant increased continuously from 30 to 60 DAT in all the treatments of Kashi Sandesh. At 30 DAT, the maximum no. of leaves (44.35 leaves) was recorded in treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was found significantly superior over other treatments followed by (43.12 and 42.31 leaves) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum no. of leaves (32.26 leaves) was recorded in treatment (T₉) - Control. At 45 DAT, the maximum no. of leaves (47.12 leaves) was recorded in treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was found significantly superior over other treatments followed by (43.52 and 43.16 leaves) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum no. of leaves (35.15 leaves) was recorded in treatment T₉- Control. At 60 DAT, the maximum no. of leaves (71.65 leaves) was recorded in treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was found significantly superior over other treatments followed by (68.59 and 67.38 leaves) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum no. of leaves (41.63 leaves) was recorded in T₉-Control. The similar result was also found by (Hemlata& Raza; 2016), Kropi&Phonglosa; (2020) and Khaleghi *et al.*,(2021).

Number of branches increased continuously from 30 DAT to 60 DAT in all the treatments. At 30 DAT, no. of branches was found significantly higher in treatment T₈ - NAA + GA₃ (40 ppm + 50 ppm (9.49 plant-1) followed by T₇- GA₃ @ 50 ppm (8.40) and T₆- NAA @ 40 ppm (7.39), whereas, lowest no. of branches (3.80) was recorded in T₁- NAA (20 ppm). At 45 DAT, significantly higher no. of branches (10.07) was recorded in (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (8.41) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm) (7.60), whereas, lowest no. of branches (4.19) was recorded in T₉- Control. At 60 DAT, significantly higher no. of branches (12.77 plant-1) was recorded in (T₈) - NAA + GA₃ (40 ppm + 50 ppm) followed by (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (11.80) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm) (10.53), whereas, lowest no. of branches (4.48) was recorded in T₉- Control. All the treatments were significantly influenced days taken to 50 per cent flowering. The minimum days to 50 per cent flowering (29.66) was obtained in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was found significantly superior over other treatments followed by (30.54) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm), whereas, maximum days to 50 per cent flowering (43.12) were noted in T₉- Control. Among the treatments, (T₈) - NAA + GA₃ (40 ppm

+ 50 ppm) recorded significantly higher number of fruits plant⁻¹ (16.17 fruits) and it was observed significantly superior over other treatments. It is followed (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (15.20 fruits). While, significantly lower number of fruits plant⁻¹ was recorded with T₁-NAA (20 ppm) (8.67). The maximum fruit length (11.66 cm) was recorded in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was observed significantly superior over other treatment followed by (11.29 and 10.25 cm) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas, minimum fruit length (5.45 cm) was recorded in T₉-Control. The maximum fresh weight (180.48 g) was recorded in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was observed significantly superior over other treatments followed by (177.99 and 176.50 g) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas minimum fresh weight (163.59 g) was recorded in T₉- Control. The maximum fruit diameter (9.48 cm) was recorded in the treatment (T₈) - NAA + GA₃ (40 ppm + 50 ppm) and it was observed significantly superior over other treatments followed by (8.49 and 7.43 cm) in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm), whereas minimum fruit diameter (3.77 cm) was recorded in T₉- Control. The result was conformed with the previous findings by Khaleghi *et al.*(2021), Bagaleet *al.* (2022) and Ruidaset *al.*(2022).

Among all the treatments, (T₈) - NAA + GA₃ (40 ppm + 50 ppm) recorded higher fruit yield plant⁻¹ (2.91 kg) over other treatments. It is closely followed by 2.71 and 2.39 kg in (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm) fruit yield plant⁻¹. While, T₁ NAA (20 ppm) had the minimum (1.44 kg) fruit yield per plant. Yield kg per plot among the treatments, (T₈) - NAA + GA₃ (40 ppm + 50 ppm) recorded significantly higher yield (29.22 kg) and it was observed significantly superior over other treatments. It is followed (T₇) - NAA + GA₃ (40 ppm + 25 ppm) (28.13 kg). While, significantly lowest yield was recorded with T₉-Control (8.67 kg). Yield of brinjal in term of quintal per hectare was recorded maximum in treatment T₈-NAA + GA₃ (40 ppm + 50 ppm) i.e. (383.95) it is followed by (T₇) - NAA + GA₃ (40 ppm + 25 ppm) and (T₆) - NAA + GA₃ (20 ppm + 50 ppm) that was (379.84 and 376.85) respectively. In other hand least yield quintal per hectare (206.69) was observed in T₉- Control for this character. It was revealed that the maximum net return of Rs. 422710.86 q/ha with Benefit:Cost ratio of 2.75 was recorded in T₈ (@ 40 ppm NAA + @ 50 ppm GA₃). That was followed by T₇ (@ 40 ppm NAA + 25 ppm GA₃) with net income of Rs. 416581.86 q/ha along with Benifit:Cost ratio of 2.71 and minimum net return of Rs.185598.25 q/ha with Benefit:Cost ratio of 1.49 was observed in control. The result was supported with the findings of Athaneriaet *al.* (2011), Patel *et al.* (2012), Veishnavet *al.* (2012), Kiranmayiet *al.* (2014) and Vandnaet *al.* (2014).

As for fruit sample observed and given below in table 3 we can say that there are no any differences was found in fruit colour, Fruit shape and fruit size with the application of various doses of growth hormone, its combination and control. All the treatment including control was showed Purple colour, Round in shape and Medium size fruits. The similar result was recorded by Khaleghi *et al.*(2021) and Ruidaset *al.*(2022).

Generally: **[9MM]Comment**
You are presenting your results as a MSc thesis
You don't have a discussion
I am not happy to read this section

CONCLUSION

On the basis of present investigation it can be concluded that the treatment T₈ (NAA - 40 ppm + GA₃-25 ppm) was produces maximum effect on most of all the growth characters and yield. Similarly, NAA - 40 ppm + GA₃-25 ppm was supported not only in growth and yield parameters while also observed economically feasible in term of higher return (benefit cost ratio) as compare to other treatments.

You don't have data to support your conclusion, and you did not do any economical analysis

UNDER PEER REVIEW

Table 1 Effect of PGR on plant height, Number of leaf per plant and number of branches per plant at different successive stages in brinjal

Treatment	Plant height (cm)			No. of leaf/plant			No. of branches/plant		
	30DAT	45DAT	60DAT	30DAT	45DAT	60DAT	30DAT	45DAT	60DAT
T ₁	27.96	44.72	61.45	37.41	37.67	54.00	3.38	4.41	5.82
T ₂	31.06	47.66	62.18	38.55	38.84	55.32	4.33	5.11	6.45
T ₃	32.55	51.03	64.42	39.36	40.37	62.53	5.35	5.74	7.34
T ₄	34.52	51.92	65.37	40.51	42.24	64.91	5.84	6.33	8.62
T ₅	34.90	53.22	66.44	41.46	43.16	66.04	6.51	6.74	9.74
T ₆	36.16	54.00	67.61	42.31	43.52	67.38	7.39	7.60	10.53
T ₇	36.18	57.84	69.40	43.12	45.29	68.59	8.40	8.41	11.80
T ₈	41.92	61.20	71.14	44.35	47.12	71.65	9.49	10.07	12.77
T ₉	27.18	37.75	45.47	32.26	35.15	41.63	3.73	4.19	4.86
SE ±	0.55	0.54	0.28	0.27	0.20	0.46	0.20	0.15	0.20
CD(5%)	1.67	1.63	0.84	0.81	0.60	1.38	0.60	0.47	0.61

Table .2 Effect of PGR on Days to 50%flowering, No. of fruit/plant, Fruit length(cm), Weight of fruits (g), Fruit diameter(cm), Fruit Yield(Kg/plant), Yield(kg/plot), Yield(q/ha), Net return and B:C ratio at different successive stages in brinjal

Treatment	Days to 50%flowering	No. of fruit per plant	Fruit length(cm)	Weight of fruits (g)	Fruit diameter(cm)	Fruit Yield(Kg/plant)	Yield(kg /plot)	Yield(q /ha)	Net return(Rs./ha)	B:CR ratio
T ₁	40.57	8.67	5.56	166.51	3.94	1.44	21.90	356.61	381785.34	2.49
T ₂	37.75	9.82	6.28	168.23	3.94	1.65	23.32	362.48	390577.82	2.55
T ₃	36.08	11.13	7.40	170.65	4.72	1.89	24.14	366.97	397301.82	2.59
T ₄	34.43	12.00	7.94	171.79	5.49	2.06	25.42	370.44	402470.82	2.62
T ₅	32.79	12.93	8.88	174.34	6.50	2.25	25.84	373.57	406829.34	2.64
T ₆	31.44	13.56	10.25	176.50	7.43	2.39	27.33	376.85	412073.34	2.68
T ₇	30.54	15.20	11.29	177.99	8.49	2.71	28.13	379.84	416581.82	2.71
T ₈	29.66	16.17	11.66	180.48	9.48	2.91	29.22	383.95	422710.86	2.75
T ₉	43.12	11.63	5.45	163.59	3.77	1.90	19.91	206.69	185598.25	1.49
SE ±	0.42	0.18	0.28	1.18	0.12	0.02	0.18	0.85	-	-

CD(5%)	1.28	0.56	0.84	3.54	0.35	0.07	0.55	2.56	-	-
---------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	---	---

UNDER PEER REVIEW




Table 3 Fruit colour, fruit shape and fruit size is influenced by different treatments

Symbol	Treatment	Fruit colour	Fruit shape	Fruit size
T ₁	NAA(20ppm)	Purple	Round	Medium
T ₂	NAA(40 ppm)	Purple	Round	Medium
T ₃	GA ₃ (25ppm)	Purple	Round	Medium
T ₄	GA ₃ (50ppm)	Purple	Round	Medium
T ₅	NAA+ GA ₃ (20 ppm +25 ppm)	Purple	Round	Large
T ₆	NAA+ GA ₃ (20 ppm + 50 ppm)	Purple	Round	Large
T ₇	NAA+ GA ₃ (40 ppm +25 ppm)	Purple	Round	Large
T ₈	NAA+ GA ₃ (40 ppm + 50 ppm)	Purple	Round	Large
T ₉	Control	Purple	Round	Small

* NAA-Nephthaline Acetic Acid ,GA₃- Gibberellic Acid and ppm- Part Per Million

References

- Athaneriya, M. K., Sengar, N. and Pandey, B. R. (2011). Influence of biofertilizer on growth and yield of chilli. *Vegetable Science*, **38**(1):101-103.
- Bagale, P., Pandey, S., Regmi, P., & Bhusal, S. (2022). Role of Plant Growth Regulator "Gibberellins" in Vegetable Production: An Overview. *International Journal of Horticultural Science and Technology*, **9**(3): 291-299.
- Batlang, V., Emongor, V.E. and Pule-Meulenburg, F. (2006). Effect of benzyladenine and gibberellic acid on yield and yield components of cucumber (*Cucumis sativus* L. cv. 'tempo'). *J. Agron.*, **5**(3): 418-423.
- Das, M., & Barua, N. (2013). Pharmacological activities of *Solanum melongena* Linn. (Brinjal plant). *International Journal of Green Pharmacy (IJGP)*, **7**(4).
- Hemlata, P., & Raza, A. K. (2016). Effect of GA₃ and NAA on growth and yield of brinjal (*Solanum melongena* Lin.) cv. Pusa purple long. *Journal of Natural Resource and Development*, **11**(1): 32-35.

- Khaleghi, S., Baninasab, B., Mobli, M., & Ehtemam, M. H. (2021). Effect of plant growth regulators on two different types of eggplant flowers regarding style length and fruit setting. *Spanish Journal of Agricultural Research*, **19**(4): e0906-e0906.
- Kiran, J., Vyakaranahal, B. S., Raikar, S. D., Ravikumar, G. H., & Deshpande, V. K. (2010). Seed yield and quality of brinjal as influenced by crop nutrition. *Indian Journal of Agricultural Research*, **44**(1): 1-7.
- Kiranmayi, P., Jyothi, K. U., Kumari, K. U., Vani, V. S. and Sneetha, D. R. S. (2014). Effect of NAA, 4-CPA and boron on growth and yield of green chilli (*Capsicum annum* L.) var. Lam-353 in summer. *Agrotechnol.* **2**(4):216-222.
- Kropi, J., & Phonglosa, A. (2020). Response of different plant growth regulators on fruit yield of Brinjal. *International Journal of Agriculture, Environment and Biotechnology*, **13**(2): 129-131.
- Meena, S.S., and Dhaka, R.S. (2003). Economics of plant growth regulators in brinjal (*Solanum melongena* L.) under semiarid condition of Rajasthan. *Annals Agric. Res.*, **24** (2): 273-275.
- Patel, J. S., Sitapara, H. H., and Patel, K. A. (2012). Influence of plant growth regulators on growth, yield and quality of tomato and brinjal. *International Journal of Forestry and Crop Improvement*. **3**(2):116-118.
- Rafeekher, M., Nair, S. A., Sorte, P. N., Hatwal, G. P. and Chandan, P. M. (2000). Effect of growth regulators on growth and yield of summer cucumber. *Journal of Soils and Crops*. **12**(1):108-110.
- Ruidas, S., Karmakar, S., Purkait, A., Gangopadhyay, A., Saha, R., Mukherjee, K., & Hazra, D. K. (2022). Preparation, optimization, and testing of biostimulant formulations as stress management tools and foliar applications on brinjal and onion for growth and yield.
- Saptari, R.T., and Dewi, K., (2013). Effect of borax and gibberellic acid on the growth and development of red chilli (*Capsicum annum* L. "gelora"). *The Third Basic Science International Conference*. (B41):1-3.
- Thompson, C. H. and Kelly, C. W. (1957). *Vegetable Crops*, Mc Graw – Hall Book Co. Inc, New York, p. 502.
- Vaja, A. D., Patel, J. B., Daki, R. N., & Chauhan, S. A. (2017). Effect of nitrogen and plant growth regulators on seed yield per plant and seed quality parameters in brinjal (*Solanum melongena* L.). *Journal of Applied and Natural Science*, **9**(4):2338-2343.
- Vandana, P. and Verma, L. R. (2014). Effect of spray treatment of growth substances at different stages on growth and yield of sweet pepper (*Capsicum annum* L.) cv. Indra under green house. *International Journal of Life Sciences Research*. **2**(4):235-240.

Veishnav, N., Singh, B. K., and Singh, A. K. (2012). Effect of NAA on growth and yield of chilli (*Capsicum annum* L.). *Environment & Ecology*. **30**(4):126.

UNDER PEER REVIEW