

Original Research Article

FINDING THE GAPS: PATIENTS SATISFACTION WITH STRUCTURE, PROCESS AND OUTCOME DIMENSIONS OF TUBERCULOSIS CARE IN PORT HARCOURT, NIGERIA.

ABSTRACT

Background:

Tuberculosis (TB) is the tenth leading cause of death and is the leading cause of death from a single infectious agent. Quality of tuberculosis care and patients satisfaction of care received varies and remains a concern to all stakeholders. The study aimed to assess patients satisfaction with the structure, process and outcome dimensions of tuberculosis care in south-south ~~nigeria~~Nigeria.

Methods: The descriptive cross-sectional study design was adopted for the study. Data collection was done using the Patient Satisfaction Questionnaire for measuring satisfaction with Quality of Care. A total of 102 patients participate. Convenience sampling of the total population attending the TB clinic during the period of study was used. Data were analyzed using SPSS version- 21.

Results: Among 102 participants 96.08% of participants were found to be satisfied ~~in~~with the structure dimension, 86.63% had good satisfaction with the process and 91.99% had good satisfaction with the outcome.

Conclusion: Majority ~~The majority~~ of the respondents had good satisfaction with the structure, process and outcome of care at the DOTS Clinic of the University of Port Harcourt Teaching Hospital.

Keywords: ~~Ppatients'~~Patient satisfaction, quality of care, tuberculosis,

INTRODUCTION

The World Health Organization (WHO) recommends the Directly Observed Treatment Short-course (DOTS) strategy for [the](#) control of TB, targeting to reduce TB deaths by 95% and new infections by 90% between 2015 and 2035 [1]. The strategy has been adopted globally and is considered the most appropriate and cost-effective approach for TB control especially in [resource-resource](#)-scarce settings [2]. Achieving these set goals is predicated upon an all-inclusive quality health care delivery system. The consensus has always been that quality care is at the core of service delivery whether the assessor is the health care provider, the user, or the facility management staff. [3]. Good quality health care, therefore, has to be effective, efficient, accessible, acceptable, patient-centered, equitable, and safe. [4] Evidence from previous studies [have-has](#) revealed an immense variation in the quality of TB care available across healthcare facilities and this continues to affect the global TB control efforts, especially in resource-limited settings like [in](#)-Nigeria [5,6]

The Nigerian National Health Policy [7] highlighted TB as a priority public health challenge and targeted [a](#) reduction in the tuberculosis prevalence rate and the tuberculosis mortality rate in Nigeria by committing to provide universal access to high-quality, client-centered TB diagnosis and treatment services.

The Institute of Medicine defines health care quality as "the degree to which health care services for individuals and populations increase the likelihood of desired health outcomes and must be consistent with current professional knowledge" [8]

Avedis Donabedian's model of quality of care consists of three key elements: structure or the resources available at a health facility, processes involving the interaction between providers and patients and outcomes or the consequences of care. [9] In other words, quality of care describes

the extent to which health care services provided to individuals and patient populations improve desired health outcomes. [7]

Patients' satisfaction is widely accepted as an independent and crucial assessment of the quality of care offered to patients. It is an invaluable tool that can give insights into daily hospital care. It provides genuine information on real experiences as well as pinpoints what is generally important to the patients and their families. [10] The other benefits of knowing patient's satisfaction is that it can identify which aspects of a service need to be changed or improved on, assist organizations in identifying patients likely to drop out or have better treatment success and higher cure rates. [11,12,13] Also, it has been shown to influence healthcare utilization and can be a predictor of subsequent health-related behavior such as whether patients are willing to recommend their healthcare provider to others. [14]

Regarding tuberculosis (TB) program services, outcome measures have often focused on mortality and microbiologic cure and most times neglect patients' perspectives such as satisfaction with care, which are crucial in influencing adherence and clinical outcomes [15,16]

Despite the increased focus on satisfaction as an important outcome measure, satisfaction has remained difficult to classify. While numerous satisfaction surveys have been developed, most with acceptable psychometric properties, the factors individual patients use to consider themselves satisfied remain largely varied or undefined. Consequently, studies measuring predictors of satisfaction have explained only a small portion of satisfaction variance, nearly always less than 20%, because patient's satisfaction is a subjective multi-dimensional health-care construct affected by many variables. [13] Nonetheless, there is an increased global call to correlate health care quality and patient satisfaction.

This study aimed to assess patients' satisfaction with [the](#) quality of care at an out-patient tuberculosis clinic in a tertiary hospital in the South-South region of Nigeria.

2. Materials and Methods

2.1 Study Area

This study was carried out in Port Harcourt, the capital of Rivers State, in the oil-rich South-South Nigeria. The city has a population of 3,171,076 according to the 2021 estimates. [17] The city has two tertiary health facilities and several other government [primary-primaries](#) and secondary health facilities as well as many private health facilities. The DOTS TB Clinic (Directly Observed Treatment, Short Course Clinic) is part of the Outpatient department of the University of Port Harcourt Teaching Hospital (UPTH). UPTH is a tertiary health care facility with about 950 beds and serves as a major referral [centre-center](#) for Rivers and nearby South-South and South-East states [18] The DOTS Clinic is a daily Monday to Friday outpatient clinic.

2.2 Study Population

The study recruited confirmed drug-susceptible TB cases (pulmonary and extra-pulmonary) who were on treatment at the adult DOTS Clinic of University of the Port Harcourt Teaching Hospital.

2.3 Sample size

The minimum sample size of 102 was calculated using the sample size formula for a single proportion [19]

The computation is based on the prevalence of TB reported in Rivers state by Otukunfor *et al.*, 2018 which revealed a 4.9% prevalence of TB in Port Harcourt, Nigeria.

2.4 Sampling technique

Convenience sampling that recruited the total population- of new and previously treated patients presenting to the clinic at the period of study was used until the required sample size was reached. The period of study was between July 2019 and December 2019.

Comment [WU1]: Keep same kine spacing

2.5 Study Instrument/ Data Collection

~~Patient-~~The patient Satisfaction Questionnaire (PSQ-17) was used to assess the quality of care from the perspective of the patient. (adapted from Ameh *et al.*, (2017) [20]-. The PSQ-17 assesses multiple dimensions of patient satisfaction and includes general satisfaction, technical quality, interpersonal relations, communication, time spent with health providers; accessibility and convenience. The PSQ-17 instrument is reflective of Donabedian's Structure-Process-Outcome(SPO) theories and concisely measures patient satisfaction with dimensions of care for which SPO constructs are proposed. It has three main areas of assessment namely: Structure Measures, Process Measures, and Outcome measures, of which a 17-item scale was developed. Three (3) main questions were used to assess Structure, eight (8) main questions were used to assess Process and six (6) questions were used to assess Outcome on a Likert scale rating of: Strongly disagree =1, Disagree =2, Not Sure =3, Agree =4, and Strongly agree =5. All items were scored from one to five (1-5) so that high scores reflect satisfaction. For the overall quality of care, the rating was on a score of 1(lowest quality of care) to 5 (highest quality). Furthermore, dichotomous data were created from the Likert scale scores with a mean score of ≤ 3 representing 'dissatisfaction' while a score of 4 and above (≥ 4) represented 'satisfaction'.

2.6 Validity and Reliability of the Instrument

The instrument was pre-tested with 10 respondents at DOTS Clinic, Ozuoba Health Centre to ensure face validity of the data. The results of the pre-test correlated positively with expected results and so the instrument was considered reliable. The reliability of the scaled items on the

study instrument was determined using Cronbach's α (alpha). It was used to measure the internal consistency of the 17-item *Patient Satisfaction Questionnaire for measuring Satisfaction with Quality of Care*. The Item responses were considered to be consistent as we obtained a reliability score of 0.875 (95%CI: 0.820 - 0.920) according to Snoek, Skovlund&Pouwer [21].

2.7 Data Analysis

Data were extracted from the questionnaires into Microsoft Excel ® version 2010 where it was coded and cleaned, and then imported into the “*IBM Statistical Package for Social Sciences (SPSS) v21 (United States) software* for Data Analysis.” Chi-square (χ^2) test analysis was performed to test for association between two or more categorical variables and their proportions (%) and to determine the level of statistical significance between the variables associated. Differences in continuous data and means were tested using the student t-test as appropriate. Bivariate analysis was performed using an odds ratio. All ORs were reported with their 95% CI and corresponding p values. An observation is said to be statistically significant if the “*p-value* is less than or equal to 0.05 (≤ 0.05)”

2.8 Ethical Consideration

Ethical approval for the study was obtained from the Ethics Committee of the University of Port Harcourt (UPH/CEREMAD/REC/MM61/047). Signed informed consent was obtained from each respondent before they were included in the study. All personal information remained confidential and was used strictly for the study.

2.9 Study Strengths and Limitations

One strength of this study was the use of the patient satisfaction survey to evaluate the quality of care in our DOTS Clinic using Donabedian's theory. Causal relationships between structure-

process-outcome constructs could not be deduced because the study design was cross-sectional.

Convenience sampling is of less statistical certainty so generalization of inference is limited.

3.0 RESULTS

Table 1: Socio-demographic Characteristics (n=102)

Characteristics	Frequency (n=102)	Percentage (%)
Age		
≤19	22	21.57
20-29	26	25.49
30-39	29	28.43
40-49	15	14.71
50-59	10	9.80
Mean	30.31 ± 10.93 years	
Sex		
Male	55	53.92
Female	47	46.08
Marital Status		
Single	67	65.69
Married	33	32.35
Widowed/Divorced	2	1.96
Educational Level		
Primary	18	17.65
Secondary	40	39.22
Post secondary	44	43.14
Occupation		
Student	29	28.43
Artisans	25	24.51
Civil Servant	14	13.73
Trading/Business	12	11.76
Contractor/Company worker	8	7.84
Health worker	5	4.90
House wife	3	2.94
Farming/Fishing	6	5.88

Table 1 shows that 28.43% of respondents were in the age range of 30-39 years with the mean age of 30.31 ± 10.93 years. More than half (53.92%) of respondents were males. Singles were in

the majority (65.69%), while 43.14% of respondents were or have attained tertiary level of education and 28.43% were students.

UNDER PEER REVIEW

Table 2: Clinical Status

Characteristics	Frequency (n=102)	Percentage (%)
How long have you been diagnosed with TB (months)		
0-5	71	69.61
6-10	27	26.47
>10	4	3.92
<i>Mean</i>	<i>4.13 ± 3.99months</i>	
HIV Status		
Positive	18	17.65
Negative	84	82.35

Table 3: Participants' level of Satisfaction with Structure, Process and Outcome

Characteristics	Satisfaction with Structure (n=102)	
	Dissatisfied Scores 1-3	Satisfied Scores 4-5
Staffing at the clinic	0 (0.0)	102 (100)
Drug Stocking at the clinic	2 (1.96)	100 (98.04)
Availability of equipments for tests and other examinations	4 (3.92)	98 (96.08)
Mean score	2(1.96)	100 (98.04)
Characteristics	Satisfaction with Process (n=102)	
	Dissatisfied (Scores 1-3)	Satisfied (Scores 4-5)
Ease of retrieving results of follow-up tests	13 (12.74)	89 (87.26)
Time spent with doctors and health workers during consultations	10 (9.90)	92 (90.10)
Follow-up care	15(14.71)	87 (85.29)

Professionalism at the clinic	2 (1.96)	100 (98.04)
Referral to specialists when the need arises	17 (16.66)	85 (83.34)
Friendliness of the staff	27 (26.47)	75 (73.53)
Defaulter tracing	14 (13.73)	88 (86.27)
Communication on medical care	11 (10.78)	91 (89.22)
Mean Score	13.6 (13.37)	88.38 (86.63)

Characteristics	Satisfaction with Outcome (n=102)	
	Dissatisfied	Satisfied
How do you feel about the tuberculosis program in general	9 (8.82)	93 (91.18)
How do you feel about the clinic to <u>being</u> able to recommend it to others?	2 (1.96)	100 (98.03)
Influence of care on your health	0 (0.00)	102 (100.00)
Impression on patients' waiting time	9 (8.82)	93 (91.18)
The medical terms used by the medical staff	21 (20.59)	81 (79.41)
How is your recovery process?	8 (7.84)	94 (92.15)
Mean Score	8.2 (8.01)	93.8 (91.99)

Table 3 shows that all the respondents reported being satisfied (scores 3 & 4) with all the structure-related dimensions of quality of care.

They were satisfied with staffing ~~at~~ the clinic, drug Stocking and availability at the clinic, and availability- of equipments for tests and other examinations with the composite score for structure averaging 96.04%.

The participants assessed process-related dimensions of care viz; ease of retrieving results of follow-up tests, Time spent with doctors and health workers during consultations, Follow-up

care, Professionalism at the clinic, Referral to specialists when the need arises, Defaulter tracing, and Communication on medical care. They were most satisfied with the professionalism of the doctors and nurses and least satisfied with the ability to refer them to another specialist when the need arises.

In assessing outcome related dimension of care, we looked at the influence of care on patient's patients' health, their impression on patients' waiting time, and the recovery process. All the participants were satisfied with the outcome of care they have received with-satisfaction with the influence of care on patient's health scoring 100%

Table 4: Relationship between Socio-demographic characteristics and Overall Satisfaction (Chi-Square & Bivariate Logistic Regression, using proportions)

Socio-demographic characteristics	Composite Satisfaction Score		Total	D f	χ^2 (p-value)	OR (95% CI)
	Poor	Good				
	Freq (%) n=8	Freq (%) n=94				
Age						
≤29	4 (50.0)	44 (46.81)	48 (47.06)	1	0.00 (1.00)	1.14 (0.27- 4.82)
30 and more	4 (50.0)	50 (53.19)	54 (52.94)			
Sex						
Male	4 (50.0)	51 (54.26)	55 (53.92)	1	0.00 (1.00)	0.84 (0.19- 3.57)
Female	4 (50.0)	43 (45.74)	47 (46.08)			
Marital Status						
Married	0 (0.0)	33 (35.11)	33 (32.350)	1	2.70 (0.100)	Undefined
Single (not in union)	8 (100.0)	61 (64.89)	69 (67.65)			
Educational Status						
≤Secondary	8 (100.0)	50 (53.19)	58 (56.86)	1	4.82 (0.03)*	Undefined
Post secondary	0 (0.0)	44 (46.81)	44 (43.14)			

*Statistically significant ($p < 0.05$); OR=Bivariate Logistic Regression; Undefined= OR cannot be determined due to an "0" value in the equation

A statistically significant association was observed between educational status and satisfaction, as persons with a secondary level of education and below showed a high proportion of dissatisfaction (100% vs. 0%; $p=0.03$).

Discussion

We documented that among 102 participants that took part in this study, 28.43% were in the age range of 30-39 years, with a mean age of 30.31 ± 10.93 years. Mean age in our study, tallies with the mean ages in other Nigerian studies (34.5 ± 13.8) and (35.4 ± 14.7). [22,23]. It is an age-long knowledge that TB is commoner among active age groups who are equally sexually and socially active. What this means is that this age group have a tendency to indulge in alcoholism and smoking, unsafe sex and a high prevalence of HIV/AIDS. More than half (53.92%) of respondents were males, 65.69% were single, and 43.14% had post-secondary education. Students and artisans constituted more than half of the respondents. Studies have shown that poor income, lower socio-economic status and little or no educational attainment are associated with an increased risk of tuberculosis infection, recurrence and relapse.[24] We had a large percentage of singles in our study due to the fact that because students especially post-secondary level students were in the majority. The proximity of the TB centres to tertiary educational centres in the state explains the large proportion of students. About a quarter of our participants (17.65%) were co-infected with HIV. [25]. World-wide, TB is the most common HIV- associated opportunistic infection and AIDS-related death. Also, progression

from TB infection to active disease is faster in ~~HIV~~-~~HIV~~-infected individuals and often exhibiting a higher tendency to poorer outcomes.

The results of this study show that with regards to the structural component of the Donabedian triad, 96.04% of our study participants were very satisfied with the quality of care received after we assessed for staff strength, availability of drugs and equipment for tests and examination. This indicates that the facility is structurally equipped and functional. This could be ~~due to the fact that~~because the tertiary health facility is a federal ~~government~~-~~government~~-owned and therefore well funded and supervised by the National tuberculosis program and partnering NGOs. Most studies reported good satisfaction with TB services but two studies from West and East Africa reported poor satisfaction. [26,27]

In assessing the processes involved in patients care, 87.3% of patients were satisfied with the waiting time and 90% were satisfied with the time spent consulting with the doctor. This result is in tandem with what was obtained in a similar study in south-east Nigeria [22] and Sudan [28] Psychologically, long waiting time before consultation is a demotivator to keeping hospital appointments and a major contributor to poor ~~care~~-~~care~~-seeking behavior of the people. Also reluctance to keep hospital appointments is a correlate of treatment failure and possibly a slide towards ~~drug~~-~~drug~~-resistant tuberculosis.

~~Friendliness~~-The friendliness of hospital staff had the lowest satisfaction rating (73%) in the service process- component of our study. The lower rating of the friendliness of staff agrees with reports from- south-east Nigeria, Northern Nigeria, and Uganda [23,29,26,30]. Anecdotaly, in Nigeria, healthcare service providers particularly in public facilities tend to be less friendly towards their clients. They perceive their service as a favour to the clients. and not a duty. In trying to regulate the attitude of all service providers towards their clients, the federal government launched the SERVICON (service compact with all Nigerians) strategy. Healthcare provider-patient relationship is often identified as a predictor of patients' satisfaction with healthcare services. therefore, staff attitude is important in service utilization. Professionalism was scored very high (98.04%) in our study. as a tertiary hospital, the highest level of expertise and candour is expected from the TB ~~centre~~-~~center~~ much more because the TB programme is vertically organised in partnership with donor funds. There is good evidence that the federal ministry of

health and donor partners are consistent in their monitoring and supervisory roles. A study from East Africa, however, reported a low scoring for professionalism but showed that participants were satisfied in the aspect of clinical consultation, time spent with the clinician, technical skills and communication with patients during consultation.[31,32]

A large proportion of the study participants rated medical care they received high. They were also satisfied with the way their medical conditions were discussed with them by the healthcare providers. On the contrary, in studying trends in patients satisfaction over a 14-year period, a Finnish team noted a 9% decrease in ~~patients'~~ patient's satisfaction [33]

This study also found that 91.99% of participants were satisfied with the outcome parameters.

The findings in this study are in agreement with the reports of Gebrekidan et al [34] and Tobin-

West et al [35]. ~~Majority~~ The majority (91.2%) of our participants claimed they can recommend

the clinic to friends and relations. In the same vein, a similar study in Pakistan- reported that 85%

of their study participants reported they can recommend ~~the~~ TB services to others.[36] All our

participants reported they were satisfied, as the ~~services~~ they received in our ~~centre~~ center

affected their health and quality of life positively. Our result is in line with the treatment

outcomes documented ~~from~~ in Ethiopia [34]. They noted that 99.9% of participants in their

study were either cured or completed their treatment.

From the findings in this study, it can be inferred that most of our participants were satisfied with

the outcome of treatment they received in our TB facility which ultimately translates into a

perception of good quality of care. Our overall result of over 90% of our participants ~~were~~ was

very satisfied with all the component TB treatment services according to Donabedian (SPO)

construct. Cazabon and colleagues[37], in order to address gaps in the care cascade, reviewed

scholarly studies on the quality of tuberculosis care in high burden countries. Their finding lent

credence to our result that a high percentage of patients were very satisfied with the quality of care received at TB [centres/centers](#).

5.2 Conclusion

This study aimed to assess patients' satisfaction with [the](#) quality of care at a tuberculosis clinic. Based on the findings in this study we conclude that majority of the respondents were satisfied with the structure, process and outcome of care at the DOTS clinic of Univerity Port Harcourt Teaching, Port Harcourt.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. World Health Organisation. (2015). WHO End TB Strategy. Global strategy and targets for tuberculosis prevention, care and control after 2015.
2. Corbett EL, Bandason T, Cheung YB, Munyati S, Godfrey-Faussett, P,& Hayes R. Epidemiology of tuberculosis in a high HIV prevalence population provided with enhanced diagnosis of symptomatic disease. 2017, *PLoS Med*, 4(1):e22.
3. Ezegwui I R, Okoye O I, Aghaji A E, Okoye O, Oguego N. Patients' satisfaction with eye care services in a Nigerian teaching hospital. 2014, *Niger J Clin Pract*. 17:585-88.
4. World Health Organization. Quality of Care; definition and Importance.. (2016) Retrieved on 16/04/2019.

5. Claassens MM, Jacobs E, Cyster E, Jennings K, James A, Dunbar, R, Enarson DA, Beyers N. Tuberculosis cases missed in primary health care facilities: Should we redefine case finding? 2013, *Int. J. Tuberc. Lung Dis.* 17, 608–614.
6. Floyd K, Glaziou P, Zumla A, & Raviglione M. The global tuberculosis epidemic and progress in care, prevention, and research: an overview in year 3 of the End TB era. *The Lancet Respiratory Medicine*, (2018), 6(4), 299–314.
7. Federal Ministry of Health, Nigeria (FMOH).. *National health policy 2016: Promoting the health of Nigerians to accelerate socio-economic development.* (2016) Abuja: FMOH.
8. Institute of Medicine. Crossing the quality chasm: A new health system for the 21st century. Washington, DC, USA: *National Academy Press.* (2001)
9. Donabedian A. Evaluating the quality of medical care. *Milbank Quarterly*, (2005) 83 (4), 691–729.
10. Gishu T, Weldetsadik AY & Tekleab AM. Patients' perception of quality of nursing care; a tertiary center experience from Ethiopia. *BMC Nurs* (2019) **18**, 37
11. Beattie M, Lauder W, Atherton I & Murphy D. Instruments to Measure Patient Experience of Health Care Quality in Hospitals: a systemic review protocol. *BioMed Central*: (2014) The open-access publishers.
12. Kraska RA, Weigand M, & Geraedts M. Associations between hospital characteristics and patient satisfaction in Germany. *Health Expectations*, 20(4), 593–600.
13. Jackson JL, Chamberlin J & Kroenke K. Predictors of patient satisfaction. *Social Science & Medicine*, (2016). 52(4), 609–620.
14. Schoenfelder T, Klewer J & Kugler J. Determinants of patient satisfaction: a study among 39 hospitals in an in-patient setting in Germany. *International Journal for Quality in Health Care*, (2011). 23(5), 503–509.
15. Babikako H M, Neuhauser D, Katamba A, & Mupere E. Patient satisfaction, feasibility and reliability of satisfaction questionnaire among patients with pulmonary tuberculosis in urban Uganda: a cross-sectional study. *Health Research Policy and Systems*, (2011) 9(1).
16. Uddin M J, Ashrafun L. Factors determining inpatient satisfaction with hospital care in Bangladesh. *Asian Soc Sci.* (2011) 7(6):15–24.
17. United Nations population estimates and projections of major Agglomerations

18. **University of Port Harcourt Teaching Hospital, Port Hrcourt. Annual Reports-2020**
19. Mohamad AP, Mohsen V & Mitra R. Sample size calculation in medical studies. *Gastroenterol Hepatol Bed Bench*, (2013) 6(1), 14–17.
20. meh S, Gómez-Olivé FX, Kahn K, Tollman SM, &Klipstein-Grobusch K. Relationships between structure, process and outcome to assess quality of integrated chronic disease management in a rural South African setting: applying a structural equation model. *BMC Health Services Research*, (2017) 17(1).
21. Snoek FJ, Skovlund SE, &Pouwer F. Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. *Health and Quality of Life Outcomes*, (2007). 5(1), 69.
22. Ohamaeme MC, Ilika F, Aniagboso CC, Elendu MU, Aniemenal CR, and Eberendu IG. Assessing the Level of Satisfaction of Patients with TB-DOTS Implementation in Rural and Urban Centres of Selected DOTS accredited facilities in Anambra State, Nigeria. *International Journal of Tropical Disease & Health* (2021) 42(5): 42-50.
23. Sani RA, Garba SA, Oyeleke SB, Abalaka ME. Prevalence of pumonary tuberculosis (PTB) in minna and suleja, Niger state, Nigeria. *American Journal of Medicine and medical Sciences*. 2015;5(6):287-291
24. Jaber AA, Khan AH, Sulaiman SA, Ahmad N. Role of socio-demographical factors on tuberculosis outcome in Yemen. *Int J Mycobacteriol*. 2016;51:S20.
25. Jiamsakul A, Lee MP, Nguyen KV, et al. Socio-economic status and risk of tuberculosis: a case-control study of HIV-infected patients in Asia. *Int J Tuberc Lung Dis*. 2018;22(2):179-186.
- 26 Adamu H, Oche MO. Patient Satisfaction with Services at a General Outpatient Clinic of a Tertiary Hospital in Nigeria. *British Journal of Medicine and Medical Research*. 2014(11):2181-2202
26. Tegegn BA, Hailu BZ, Tsegaye BD, Woldeamanuel GG, Negash W. Quality of Care Delivered to Tuberculosis Patients among Public Hospitals in Central Northeast Ethiopia. *Ethiop J Health Sci*. 2020;30(5):671-680.
27. Elsadig M, Mohamed O, Mohamed A, Sawsan A, Khalid M, Hatim S, Waqas S. Patients satisfaction with tuberculosis services of directly observed therapy programs in Gezira state of Sudan. *Arch Clin Infect Dis*. 2014;9(4):1-5.
28. Gezahegn G, Gezahegn T, Mitiku TH, Negussie D. Quality of tuberculosis care in private health facilities of Addis Ababa, Ethiopia. *Tuber Res Treat*. 2014;2014:1- 20.

29. Ugochukwu U. Onyeonoro, Joseph N. Chukwu, Charles C. Nwafor², Anthony O. Meka², Babatunde I. Omotowo³, Nelson O. Madichie², Chidubem Ogbudebe², Joy N. Ikebudu², Daniel C. Oshi², Ngozi Ekeke Evaluation of Patient Satisfaction with Tuberculosis Services in Southern Nigeria. *Health Services Insights* 2015;8 25–33
30. ³¹ Ssenooba W, Kirenga B, Muwonge C, Kyaligonza S, Kasozi S, Mugabe F, Boeree M, Joloba M, Okwera A, PanACEA. Patient satisfaction with TB care clinical consultations in Kampala: a cross sectional study. *Afri Health Sci.* 2016;16(4): 1101-1108.
- 32 Ibrahim L.M., Idris Hadejia IS, Nguku P. Factors associated with interruption of treatment among pulmonary tuberculosis patients in Plateau State, Nigeria. *Pan Afr Med J.* 2014
- 33 Raivio R, Jääskeläinen J, Holmberg-Marttila, D. Decreasing trends in patient satisfaction, accessibility and continuity of care in finnish primary health care - a 14-year follow-up questionnaire study. *BMC Fam Pract.* (2014) 15:98.
- 34 Gebrekidan, G., Tesfaye, G., Hambisa, M. T., & Deyessa, N. Quality of Tuberculosis Care in Private Health Facilities of Addis Ababa, Ethiopia. *Tuberculosis Research and Treatment*, 2014, 1–9.
- 35 Iqbal, A., Bilal, A. Prevalence of Tuberculosis with Clinical Implications in District Sargodha. *J Antivir Antiretrovir.* 2021;13:227.
- 36 Ali SM, Anjum N, Naureen F, Rashid A, Tahir A, Ishaq M et al. Satisfaction Level of Tuberculosis Patients Regarding Their Access to TB Care and Prevention Services, Delivered Through a Public-Private Mix Model in Pakistan. *Healthcare.* 2019 Oct 18;7(4).
- 37 Cazabon D, Alsdurf H, Satyanarayana S, Nathavitharana R, Subbaraman R, Daftary A., & Pai M. Quality of tuberculosis care in high burden countries: the urgent need to address gaps in the care cascade. *International Journal of Infectious Diseases*, (2017). 56, 111–116.16.