

## **Original Research Article**

### **Ecological Adaptation of Endemic *Anthocleista* Species under Moisture Gradient in Parts of Niger Delta, Nigeria**

#### **ABSTRACT**

**Background:** Knowledge of the anatomy of *Anthocleista* species is crucial for understanding how these plants adapt to the environment. **Aim:** This study was aimed at investigating the influence of moisture gradient on the anatomy of four species in the genus *Anthocleista* (*A. djalonesis* A. Chev; *A. liebrechtsiana* De Wild & Th.Dur; *A. nobilis* G.Don; and *A. vogelii* Planch) in light of environment and ecological niche adaptation. **Place of Study:** Alakahia in Obio / Akpor, Rivers State, parts of Niger Delta. **Method:** Conventional classical anatomical techniques for structural sectioning were used. **Result:** Though there are similarities and differences in vascular structure among the species, the study has revealed variance in anatomical responses to moisture gradient ranging from dry mesophytic to mesophytic and wetland conditions of adaptation. The most important and distinct features observed are the presence of sclerenchymatous idioblast, air sacs and sclereidal idioblast. Sclerenchymatous idioblasts are numerous in *A. liebrechtsiana*, few in *A. nobilis* and *A. vogelii* but was lacking in *A. djalonesis*. Therefore the sclerenchymatous idioblast seen in *A. liebrechtsiana*, *A. nobilis*, and *A. vogelii* confirmed these species to be mesophytic to semi-aquatic in their habitat adaptation, while *A. djalonesis* in dry mesophytic adaptation had thicker epidermal layer, multiple hypodermal layer, thicker mesophyll tissues with increased number of palisade layers and thick leaves. The stem and root modification had sclereidal idioblast in abundance distribution in *A. liebrechtsiana*, and *A. nobilis*, moderate in *A. vogelii* and very low in *A. djalonesis*. **Conclusion:** The variation observed in the leaf, petiole, stem and root anatomy was due to environmental variability of anatomical characters with the plant species evolving certain structures and modification to adapt to the niche and environment where they find themselves.

**Key words:** *Idoblast, air sac, leaves, stem, and root.*

## 1.1. INTRODUCTION

One of the most critical features and influences of comparative analysis on plant characteristic variation and similarity in relation to eco-environmental adaptation depends on the degree of relationship of various aspect of function in anatomical characterization of plant (Harvey *et al.*, 1995) as well as phylogenetic relatedness among taxa (Dubuisson *et al.*, 2011). Every plant found in any environmental positions either tropical or sub-tropical region has mechanism for its survival in such place. The adaptation potential may comprise of the plant physiological, and biochemical roles in adapting to the kind of environment found. This makes species partly dependent on any statistical inference, by relating the anatomical traits to ecological preferences which might be alternatively explained by phylogenetic inertia affecting both the similarity of anatomical traits among closely related taxa and the similarity of ecological niches such taxa occupy (Harvey *et al.*, 1995).

The Environment can be regarded as the sum total of all the external conditions (biotic and abiotic factors) affecting the development and life of an organism (Okiwelu and Anyanwu, 2003). These environmental factors do affect the growth, development and distribution of plant in different habitats. Plants species are known to have different requirement of environmental conditions which can directly or indirectly be sources of ecological stress or / and balance for such species (Jegade, *et al.*, 2011). These have made plants to develop a variety of strategies and mechanisms in terms of changes in morpho-anatomical structures to adapt to any changes in their environment. To survive and reproduce, all living organisms must adjust to conditions imposed on them by their environment. Thus Plants species have to respond positively or negatively to changes in their habitat in order to survive (Iwara, *et al.*, 2011). Such conformity between an organism and its environment constitutes what biologist call "Adaptation". Adaptation is an evolutionary process whereby living things are better suited in tolerance to live in their environment or habitat (Donald, *et al.*, 2007). It is a deliberate process of change or compromise which contributes to fitness and survival of a plant in anticipation or in response to external stimuli and stress. Invariably, it is the ability of an organism to adapt and survive in its environment in different ways; involving an increase or / and decrease in temperature, moisture, humidity, wind or light, dark movement (Donald *et al.*, 2007).

The influence of ecological adaptation of organism in an environment, involve the integration of biotic and abiotic factors to the behavioral aspects of species such as plant in the environment (Townsend *et al.*, 2011). Plants species of different families or same family have varied genetic potential to adopt wide range of environmental conditions. This adaptation potential might be due to varying physiological, biochemical and anatomical characteristics (Noman *et al.*, 2012). A number of environmental factors including available moisture are known to influence the anatomy of plant species. Variation in plants growth, distribution and productivity has often been attributed to varying conditions of environmental factors (Edward and Richardson, 2004; Christensen *et al.*, 2007). Though it has been suggested that anatomical features in plants might be more important than morphological features systematically because anatomical features are less susceptible to environmental change. However, it was reported that variation in leaf anatomy might be useful in assessing strong environmental influences on plants (Stace, 1991)

The plants which characteristically grow in certain ecological niches often show a type of structure which is believed to be adapted to that particular environment besides the phenotype. Plant species inhabiting water or dry habitats exhibit a number of special features or adaptations in their

morphology, anatomy and physiology which enables them to survive under such extreme ecological conditions (Verma and Agarwal, 1983). It has been revealed that variation in anatomical structure is dependent on several forces of prevailing conditions in the habitat leading to species adaptation (Niklas, 1985). Knowledge of the anatomy of *Anthocleista* species is crucial for understanding how these plants adapt to the environment. Such plants exhibit considerable variation in their requirements of water in natural soil habitat. The genus *Anthocleista*, Afzel.ex.R.Br., commonly called cabbage tree is a medium size tropical Africa genus composed of medium, small trees or shrubs with soft white wood belonging to the family Loganiaceae (Keay, 1989; Edwin –Wosu *et al.*, 2015). It consist of 50 species, mainly native to tropical African mainland, Madagascar and Mascarene Island; with six (6) of the 50 species commonly found in Nigeria, and economically significance (Keay, 1989). Of the six species in Nigeria, phytogeographical distribution study has revealed four (4) species of common occurrence with preference for diverse ecological adaptation in parts of Niger Delta (Edwin-Wosu and Omara – Achong, 2010; Edwin-Wosu *et al.*, 2015).

In search of understanding the phylogeny of the genus and relationship among the species in parts of Nigeria, several studies have been carried out. Among which includes: the origin and species composition of the genus (Keay, 1989; Backlund *et al.*, 2000); phytomorphological characterization, similarity and variation among species (Sonibare *et al.*, 2007; Edwin-Wosu *et al.*, 2010; Edwin-Wosu, 2012); phytogeographical distribution, habitat adaptation of the species (Edwin-Wosu, and Omara-Achong, 2010; Edwin-Wosu *et al.*, 2015); A Chemotaxonomic study of three species of *Anthocleista* (Sonibare *et al.*, 2007); histochemical localization of tannins in species of *Anthocleista* found in parts of Niger Delta Tropical rainforest of Nigeria (Edwin-Wosu and Ndukwu, 2012). Others include foliar trichome morphology of the species of *Anthocleista* in Parts of the Niger Delta (Edwin-Wosu *et al.*, 2012); stomata morphology of species in parts of Niger-Delta (Edwin-Wosu and Ndukwu, 2012); the antiviral effect of *Anthocleista nobilis* root extract on the biochemical indices of poultry fowls infected with Newcastle disease virus (Ayodele *et al.*, 2013); Phytochemical screening, antimicrobial and antioxidant activity of *Anthocleista djalonesis* and *Anthocleista liebrechtiana* (Luter *et al.*, 2012; Ngbolua *et al.*, 2014); impact of *Anthocleista vogelii* root bark ethanolic extract on weight reduction of induced obesity in male Wister rat (Anyanwu *et al.*, 2013); Morphology and Anatomy of *Anthocleista djalonesis* (Asuzu and Nwosu, 2009); the ecological dynamics and trajectories of bioactive compounds of species in parts of Niger Delta (Edwin-Wosu *et al.*, 2017) and evaluation of diverse potential in the genus *Anthocleista* (Edwin-Wosu and Okafor, 2017). The *Anthocleista* species occurring in the Niger Delta has shown tremendous ecological diversity with a view to adaptation preference among the species ranging from terrestrial dry habitat, mesophytic to swampy habitat (Edwin-Wosu and Omara-Achong, 2010). So far no research has been carried out on the relationship between the environmental adaptation and anatomical modification of the species.

There is therefore the need to investigate the relationship in the anatomical modification of these species with respect to their prevailing habitat. This present study is significant hence it will provide some vital information on this genus, with the view that the variation observed in the anatomy of the species can help understand how they adapt to various ecological niches, determine the influence of environmental and ecological niche adaptation on the petiolar, foliar, stem and root anatomy of *Anthocleista* species found in parts of Niger-Delta. The information obtained is expected to aid in the assessment of taxonomic value of these features as well as their ecological niche adaptation. Also such

useful information could possibly prove as a baseline in taxonomic delimitation of the different taxa and enhance the existing Floristic information on the species and generally add to the already existing information on the taxonomy and the ecographical amplitude of the species.

## 2.0. MATERIALS AND METHODS

### 2.1. Description of study area, location and site

The Niger Delta area is the coastline parts of Nigeria approximately 853 kilometers facing the Atlantic Ocean and lying between latitude  $4^{\circ} 10'$  to  $6^{\circ} 20'N$  and longitude  $2^{\circ} 45'$  to  $8^{\circ} 35'E$ . Geographically the area is the Southern segment of Nigeria, created by myriads of Islands segmented by lagoons and channels, which empty into the Bight of Benin in the East Atlantic Ocean. The Delta is supplied with water by the Rivers Niger and Benue. These Rivers (now joined) break up at Ebu-Otor into the Rivers Nun and Forcados (including their tributaries). The portion of the Niger Delta traversed by the River Nun is the present Bayelsa State, while the present Delta State is traversed by the River Forcados. The Nun River breaks out into many channels and creeks such as Santa Barbara, St. Nicholas, Brass, Nun, Sangana, Fishtown, Koluama, Middleton, Digatoru, Pernnington, Dodo and Ramos Rivers, which empty into the Atlantic Ocean (Alagao, 1999) (Fig. 1).

From the ecological perspective it is that portion of the Southern Nigeria stemming from Northern apex situated at Aboh, bounded in the East and West by the Imo and the Benin Rivers respectively and on the South by the Atlantic Ocean (Fubara *et al.*, 1988). It is the Africa's largest Delta covering some 70,000 square kilometers ( $Km^2$ ), of which one third of the area is made up of wetland (Afolabi, 1988). The region is low lying and not more than 3.0 meters above sea level (Dublin-Green *et al.*, 1999). It possesses myriads of luxuriant landscape of sensitive vegetation and ecosystems ranging from coastal mangrove forest, brackish swamp forests, fresh water swamp forest, barrier island forest and low land rainforest which are attributes of a large river delta in a tropical region (Teme, 2001). The environmental condition of the area has two climatic seasons: a rainy season (April to November) and a dry season (December to March), with an annual rainfall ranging between 1500 and 4000mm (Kurnk, 2004). The mean monthly temperature varies between  $24^{\circ}C$  and  $32^{\circ}C$  throughout the years, thus exhibiting maximum temperature of climatic regimes. Seasonal and latitudinal variations do affect the maximum temperature, diurnal and seasonal ranges, relative humidity, high rainfall pattern, which is comparatively uniform due to the proximity of the region to the Atlantic Ocean (Alagoa, 1999). The soil condition of the area is of sandy silt and clayey, often alkaline and salty, sometimes acidic. The Niger Delta based on ecological context of triangular deltaic basin alluvium formation consists of three core States: Bayelsa, Delta and **Rivers – study location** (Fig.1). Others based on regional and political context include Akwa Ibom, Cross River, and Edo, Abia, Imo and Ondo States.




Fig. 1: The Niger Delta study area showing Rivers State – study location

Rivers State is situated between longitude  $6^{\circ}38'59.414''\text{E}$  and  $6^{\circ}45'14.471''\text{E}$  and latitude  $4^{\circ}21'26.725''\text{N}$  and  $5^{\circ}43'33.358''\text{N}$ . lies on the coastal plain of the Eastern Niger- Delta with its land surface geology of fluvial sediments. The land surface can be grouped into three main divisions: the fresh water; the mangrove swamps and coastal sand ridges zone. These surfaces consist of three main soil groups: the fresh water brown loams and sandy loams, the fluvial marine sediments and Mangrove swamp alluvial soils. The State is characterized with high rainfall, which decreases Northward annually from 4700mm on Coast to about 1700mm in north apex of the State. The rainforest vegetation occupies the upland area; the riverine area is divisible into three hydro-vegetation zones: the Salt water zone; the Beach ridge zone and the Fresh water zone (Online Nigeria Community Portal of Nigeria, 1998). The State houses 23 local council areas including **Obio / Akpor study site** (Fig. 2).

Obio/Akpor Local Council Area situated between long.  $6^{\circ}56'16.932''\text{E}$  &  $7^{\circ}1'11.892''\text{E}$ ; and lat.  $4^{\circ}56'22.948''\text{N}$  and  $4^{\circ}47'0.256''\text{N}$  (Fig. 2) covers an area of 320 square Kilometres, approximately bounded by Ikwerre, Etche and Omuma Council areas in the north, Emohua Council in the west, Degema, Okrika, Eleme and Oyigbo Councils in the east. It is generally flat lowland with a mean elevation range of 15m above sea level. The Sombreiro and New Calabar rivers are the major rivers, with some isolated seasonal fresh water swamps common in the area. The is a mesophytic seasonally flooded terrestrial habitat, dominated with sandy-silt and silty-clay textural soil property and characterized by maximum temperature, relative humidity and maximum rainfall. The vegetation type is a shrubby re-growth, predominated by shrubs, grasses herbs and few trees (Knoepp *et al.*, 2000).





Fig. 3: Odo/Akpor study site showing Alakatia UPTH sampled site




Fig. 4 satellite imagery of study site indicating UPTH species sampled site

### 2.3. Laboratory anatomical study

Fresh specimen of the leaf, petiole, root and stem of collected samples were fixed in FAA 1:1:18 (1 part formalin, 1 part glacial acetic acid, 18 parts of 70% ethanol (v/v)) for 48 hours (Johansen, 1940). The specimen were washed in two changes of distilled water to get rid of the fixative and passed through graded ethanol series in the order 30%, 50%, 70% and 95% ( each for two hours) and finally absolute ethanol (overnight) (Okoli, 1987). Free hand sections were made using a razor blade to cut thin section (1 cell thick) by making horizontal cuts close to the surface. The first sections were discarded; subsequent sections placed in water, held in a glass Petri dish. The sections (leaf and petiole) were stained with 1% methyl blue, root and stem with 1% safranin and methylblue (canter staining), the sections were rinsed with distilled water to get rid of excess stain and then placed on a slide with a drop of glycerine and covered with a cover slip. The prepared slides were viewed under a microscope and photomicrograph was taken from good preparations.

### 3.0. RESULTS

The result of the spatial geographical distribution and herbarium voucher sample deposit are presented in Tables 1 and 2. The anatomical result interpretation of study observations on the leaf, petiole, root and stem of the species are presented below, while the photomicrographs of the sections as shown are presented in Plates 1-16.

**Table 1: GPS co-ordinates of the collected samples.**

Species	Latitude (N)	Longitude (E)	Alt. (Metres)
<i>A. djalonesis</i> A. Chev	4° 53'44.33"	6° 55'52.06"	16
	4°53'43"	6°55'56"	16
<i>A. liebrechtsiana</i> De Wild & Th. Dur	4° 53'37.13"	6° 55'42.15"	11
	4°53'43"	6°55'39"	11
<i>A. nobilis</i> G. Don	4° 53'36.09"	6° 55'45.69"	11
	4°53'37"	6°55'45"	11
<i>A. vogelii</i> Planch	4° 53'36.49"	6° 55'49.36"	13
	4°53'38"	6°55'45"	13

**Table 2: Data for the Herbarium Voucher Samples Deposit**

S/No	Species	Accession No	UPH No
1.	<i>A. vogelii</i>	001	UPH/V/1204
2.	<i>A. djalonesis</i>	002	UPH/V/1205
3.	<i>A. liebrechtsiana</i>	003	UPH/V/1206
4.	<i>A. nobilis</i>	004	UPH/V/1207

### 3.1. Leaf -Midrib Anatomy

The transverse section of the leaf – midrib anatomy has revealed *Anthocleista djalonesis* A.Chev leaf (Plate 1) with two thick layer of epidermis, multiple hypodermis, thick layer of elongated palisade

mesophyll parenchyma, layer of irregular spongy mesophyll parenchyma cell, several crescent shaped and bicollateral vascular bundles with single layer of endodermis. *Anthocleista liebrechtiana* De Wild and Th. Dur (Plate 2) has two layer of epidermal cells, single layer of collenchymatous cells, single thick layer of elongated palisade mesophyll and irregularly shaped thick spongy mesophyll, stellate shaped idioblast in the spongy mesophyll tissue, presence of air sacs (Lacunae) in the leaf lamina and vascular bundles. *Anthocleista nobilis* G.Don (Plate 3) recorded two layer of upper epidermis, a single layer of collenchymatous cells, a single layer of elongated palisade mesophyll and spongy mesophyll cells, several idioblastic sclereids, several bicollateral vascular bundles and fourteen medullary bundles, while *Anthocleista vogelii* Planch (Plate 4) had single layer of epidermal cells, single layer of collenchymatous cells and two layers of elongated palisade mesophyll and irregular layers of spongy mesophyll, crescent shaped vascular bundles, several bicollateral vascular bundles arranged in a concentric ring with eight medullary vascular bundles.





Plate 1: Transverse sections of the Leaf/Midrib of *Anthocleista dialonesis* A.Chev (x10)  
a: Epidermis; b: Hypodermis; c: irregular spongy mesophyll parenchyma; d: vascular bundle (phloem) e: vascular bundle (xylem); and f: endodermis



Plate, 3: Transverse sections of the leaf / midrib of *Anthocleista nobilis* G.Don.  
(x10) a: Epidermis; b: Palisade mesophyll; c: Vascular bundles; d: Endodermis; e: Collenchyma




Plate 4: Transverse section of the midrib of *Anthocleista vogelii*. Planch (x10)  
 a: Epidermis; b: Palisade mesophyll; c: Spongy mesophyll; d: Crescent vascular bundle; e: Bicollateral vascular bundle (Phloem); f: Vascular bundle (Xylem); g: Endodermis

### 3.2. Petiole Anatomy

The petiole anatomy from the various species has indicated *Anthocleista djalonesis* A.Chev (Plate 5) with single layer of epidermal cell, 3 layers of collenchymatous cells, a layer of sclerenchyma cells, spongy parenchyma cells containing a lot of intercellular spaces. Vascular bundles are amphicribal bicollateral arranged in a concentric ring. There are seven medullary vascular bundles scattered within the pith representing vein traces.

*Anthocleista liebrechtiana* De Wild and Th. Dur (Plate 6) transverse section recorded single layer of epidermal cell, 3 layers of collenchymatous cells, single layer of sclerenchymatous cell, numerous stellate shape idioblast and several bicollateral vascular bundles.

*Anthocleista nobilis* G.Don (Plate 7) had single layer of epidermis, two to three layers of collenchymatous cells, one layer of sclerenchyma cell and several bicollateral vascular bundles. sclerenchymatous idioblast are present, some with stellate shape, others with simple or complex branching, while *Anthocleista vogelii* Planch (Plate 8) with it heart shape, recorded single layer of thick epidermal cells, 2-3 layers of collenchymatous cells, a single layer of sclerenchymatous cell, and several bicollateral vascular bundles. It has numerous idioblast, some with stellate shape, others with simple or complex branching.





Plate 5: Transverse sections of the petiole of *Anthocleista djalonesis* A.Chev x10.  
a: Epidermis; b: Collenchyma; c: Sclerenchyma; d: Vascular bundle; e: Medullary bundle  
f: Intercellular space



Plate, 7: Transverse section of the petiole of *Anthocleista nobilis* G.Don x10, a: Epidermis; b: Collenchyma; c: Sclerenchyma; d: Vascular bundle; e: Sclerenchymatous idioblast




Plate 8: Transverse sections of the petiole of *Anthocleista vogelii* Planch (x10).  
a: Epidermis; b: Collenchyma; c: Vascular bundle; d: Sclerenchymatous idioblast

### 3.3. Root and Stem Anatomy

The transverse section of the root and stem anatomy has revealed the internal structure of the four species studied (Plates 9 – 16).

The root of *A. djalonensis*. A. Chev (Plate 9) had two layers of epidermis, cortex with the presence of air chamber, endodermis, pericycle, seven vascular bundles and pseudopith. *Anthocleista liebrechtsiana*. De Wild and Th. Dur., (Plate 10) recorded two layers of epidermis, exodermis, several air chambers, cortex, several sclereid, endodermis, 12 vascular bundles and pseudopith, while *A. nobilis*. G. Don (Plate 11) revealed two layers of epidermis, a layer of exodermis, cortex, several air chambers, endodermis, several vascular bundles, and several astrosclereid and pseudopith. *Anthocleista vogelii*. Planch (Plate 12) recorded two layers of epidermis, cortex, several air chambers, endodermis, pericycle, several vascular bundles and pseudopith.




Plate 9: Transverse section of *A. djalonensis* Root (x10). a: Epidermis; b: Cortex; c: Air chamber; d: Endodermis; e: Pericycle; f: vascular bundles; g: Pseudopith




Plate 10: Transverse section of *A. liebrechtsiana* Root (x10). a: Epidermis; b: Exodermis; c: Air space; d: Cortex; e: Sclereid; f: Endodermis; g: Vascular bundles; h: Pseudopith




Plate 11: Transverse section of *A. nobilis* Root (x10). a: Epidermis; b: Exodermis; c: Cortex; d: Air chamber; e: Endodermis; f: Vascular bundles; g: Pseudopith




Plate 12: Transverse section of the Root of *A. vogelii* (x10). a: Epidermal cell; b: Cortex; c: Air chamber; d: Endodermis; e: Pericycle; f: Vascular bundles; g: Pseudopith

The stem anatomy of *Anthocleista djalensis*, A. Chev (Plate 13) has revealed a thick layer of epidermis, cortex, several astrosclereid, 9 layers of collenchyma, endodermis, several vascular bundle, and pseudopith. *Anthocleista liebrechtsiana*, De Wild and Th. Dur (Plate 14) had a layer of epidermis, layers of collenchyma, endodermis, several vascular bundles, several astrosclereid and pseudopith. *Anthocleista nobilis*.G. Don (Plate 15) recorded pseudopith, several sclereid, vascular bundles, endodermis, collenchyma cells and epidermal cells. *Anthocleista vogelii*. Planch (Plate 16) revealed layers of epidermis, layers of collenchyma, endodermis, several vascular bundles, and few sclereid.




Plate 13: Transverse sections of *A. djalensis* stem (x40). a: Epidermis; b: Cortex; c: Scleried; d: Collenchyma; e: Endodermis; f: Vascular bundles; g: Pseudopith




Plate 14: Transverse sections of *A. liebrechtsiana* stem (x40). a: Epidermis; b: Collenchyma; c: Endodermis; d: Vascular bundle; e: Sclereid; f: Pseudopith




Plate 15: Transverse sections of *A. nobilis* stem (x40). a: Pseudopith; b: Sclereid; c: Vascular bundle; d: Endodermis; e: Collenchyma cells; f: Epidermal cells




Plate 16: Transverse sections of *A. vogelii* stem (x40). a: Epidermal cells; b: Collenchyma cells; c: Endodermis; d: Vascular bundle; e: Pseudopith; f: Sclereid

#### 4. DISCUSSION

Every plant found in any environmental positions either tropical or sub-tropical region has mechanism for its survival in such place. The adaptation potential may comprise of the plant

physiological, and biochemical roles in adapting to the kind of environment found. This makes species partly dependent on any statistical inference, by relating the anatomical traits to ecological preferences which might be alternatively explained by phylogenetic inertia affecting both the similarity of anatomical traits among closely related taxa and the similarity of ecological niches such taxa occupy (Harvey *et al.*, 1995). Plants species are known to have different requirement of environmental conditions which can directly or indirectly be sources of ecological stress or / and balance for such species (Jegede, *et al.*, 2011). To survive and reproduce, all living organisms must adjust to conditions imposed on them by their environment. These have made plants to develop a variety of strategies and mechanisms in terms of changes in morpho-anatomical structures to adapt to any changes in their environment. Thus Plants species have to respond positively or negatively to changes in their habitat in order to survive (Iwara, *et al.*, 2011).

Plants species of different families or same family have varied genetic potential to adopt wide range of environmental conditions. This adaptation potential might be due to varying physiological, biochemical and anatomical characteristics (Noman *et al.*, 2012). A number of environmental factors including available moisture are known to influence the anatomy of plant species. This has informed the study on the ecological adaptation *Anthocleista* species under moisture gradient ranging from dry mesophytic secondary terrestrial forest and mesophytic to wetland condition in parts of Niger Delta.

The result obtained from this investigation has though revealed some similarities and differences among the four species of *Anthocleista* in term of distribution and abundance in the anatomical features adaptation to environmental conditions. It has been suggested that anatomical than morphological features in plants might be more important systematically due to its less susceptibility to environmental change; leaf anatomy appears to vary according to the environment (Johnson, 1980); however, it has been reported that variation in leaf anatomy might be useful in assessing strong environmental influences on plants (Stace, 1991). Though the leaf anatomy has recorded similarities and differences in their vascular structure (Edwin-Wosu and Omoikhabon *in press*) in term of distribution and abundance, the present study observed that anatomical features are generally similar to all the species, which indicate a phylogenetic relatedness of the taxa

Variation in anatomical features of plants has often been attributed to environmental factors (Scheiner and Goodnight, 1984). The most striking feature observed was the presence of sclerechymatous idioblast reaffirming the assertion by Leeuwenberg and Leehourts (1980). The present study observed that sclerenchymatious idioblasts are numerous in *A. liebrechtsiana*, few in *A. nobilis* and *A. vogelii* but lacking in *A. djalonesis*. Earlier study has recorded *A. djalonesis* as a secondary terrestrial forest species with a dry mesophytic habitat adaptation. (Edwin-Wosu *et al.*, 2015). This may be the reason why it lacks sclerenchymatous idioblast, and contain so many intercellular spaces as well as its cuticle being well developed to help prevent excessive water loss. *Anthocleista nobilis* and *A. vogelii* has exhibited greater ecological amplitude hence their capability to establish in various habitat lying along an environmental gradient (Edwin-Wosu *et al.*, 2015). This also corroborate the assertion that landscape are never static, their element are in permanent temporal and spatial flux (Stemier and Kohler, 2003; Brown and Leband, 2006). The numerous sclerenchymatious idioblast and air sacs (Lacunse) observed in *A. liebrechtsiana* shows that it preference for adaptation is a semi-aquatic (Wetland) environment. The species is known for its low ecological amplitude with a localized distribution because of the narrow range of condition on which their growth depend (Edwin-Wosu *et al.*, 2015). Such species

are called “*habitat specialist*” because they have a significant positive association with their habitat or they cannot survive outside their habitat (IUCN, 1994). Specialist species have narrow niche breadth and occur only in a small geographical range where appropriate resources are available.

Presence of sclerenchymatous idioblast and lacunae appears to be characteristic features of species growing in wetland associated with moisture gradient. Observation has shown that despite the species profound preference for tropical and mesophytic environment, they also display disparity in light of their features in restricted occurrence in certain ecological habitats (Edwin-Wosu and Omara-Achong, 2010; Edwin-Wosu *et al.*, 2015). Similar study has indicated *A. djalonesis*, *A. nobilis*, and *A. vogelii* as prevalent and adapted to low land secondary vegetation forest while *A. liebrechtsiana* is prevalent and adapted to fresh water swamp forest zone (Edwin-Wosu and Omara-Achong, 2010). Plants which characteristically grow in certain niches often show a type of structure believed to be adapted to that particular environment; therefore sclerenchymatous idioblast have been seen as an adaptive feature of plants to watery niche (Verna and Agawal 1983).

The presence of sclereide of various shapes (sclerenchymatous idioblast) in leaves and other tissue is the characteristic of hydrophytes. The variation observed in the leaf anatomy was due to environmental variability of anatomical characters. Therefore the sclerenchymatous idioblast seen in *A. liebrechtsiana*, *A. nobilis* and *A. vogelii* confirmed these species to be semi-aquatic while *A. djalonesis* is mesophytic in their habit adaptation. Related study of such environmental influence as low temperature, drought, light and elevation on responses of leaf structures and general ecological trend have been recorded (Fei *et al.*, 1999; Rocas *et al.*, 2001; Li and Bao, 2005). Mesophytes are plants that are exposed to an environment where it is neither too dry nor too wet; they do not need any extreme adaptations. However, the mesophytic adaptation of *A. djalonesis* is associated with thicker epidermal layer, multiple layer of hypodermis, thicker mesophyll tissues with increased number of palisade layers and thicker leaves. This corroborates Rocas *et al.* (2001) where instances of plants in low temperature zones are characterized with smaller leaf area, thicker epidermis, thicker mesophyll tissue and thicker leaves. Similarly, increased number of palisade layers with small cell volume, decreased number of spongy layers with small cell volume and smaller intercellular spaces has been recorded among plants in zones of water shortage (Chartzoulakis *et al.*, 2002). Smaller and thicker leaves; well-developed epidermal cuticles and trichomes; a thicker epidermal layer composed of small cells; the appearance of hypodermal layer; well-developed palisade tissues and tightly arranged spongy tissues have also been recorded among plants in zones of high light intensity (Mendes *et al.*, 2001; Cai and Song, 2001; Rocas *et al.*, 2001). In plants, leaf anatomy might provide insight to assess adaptive variation and ultimately evolutionary change (Stace, 1991).

The anatomical study has also revealed the internal structure of the stem and root of the four species of the genus *Anthocliesta* studied under moisture gradients. The stem epidermis, collenchyma, endodermis, xylem, phloem, parenchyma and sclerenchyma cells are present in abundance distribution in the four species while sclereid is found in abundance distribution in *A. liebrechtsiana* and *A. nobilis*, moderate in the stem of *A. vogelii* and very low distribution in *A. djalonesis* which has a well developed cortex. In the root; epidermis, cortex, endodermis and vascular bundles are present in abundance distribution in the root of the four species, air chamber is present in abundance distribution in the root of *A. liebrechtsiana* and *A. nobilis*, and moderate in distribution in the root of *A. vogelii* and *A. djalonesis*, which were also noted for the presence of pericycle while sclereid was observed in *A. liebrechtsiana*

and *A. nobilis*. Anatomical characters may serve as reliable indicators in the study and understanding of ecological adaptation of living organisms. Therefore anatomical studies of leaf, petiole, stem and root could be used to deduce the influence of environmental and ecological niche adaptation on the anatomy of *Anthocleista* species found in parts of the Niger-Delta.

## CONCLUSION

The result obtained from the anatomical studies of the four *Anthocleista* species shows that the environment and ecological niches where they are found have an influence in their anatomical structures. The variation observed in the anatomical features of the plants can be attributed to environmental factors such as moisture gradient beside other factors not considered in this study. The anatomical features seen in the plants indicate that *Anthocleista* species shows an ecological preference for both a normal terrestrial (dry mesophytic and mesophytic) and wetland (seasonally flooded environment) conditions. Also the plant species tends to evolve structures such as sclerenchymatous idioblast and lacunae to adapt to the niche they occupy.

## REFERENCES

- Afolabi D. The Nigerian Mangrove ecosystem. Third Regional workshop of the Gulf Guinea Large Marine Ecosystem (GOGLME), Lagos Nigeria; 1998.
- Alagoa E.J. The Land and People of Rivers State, Central Niger Delta. Onyema Research Publication, PortHarcourt, Rivers State, Nigeria; 1999
- Anyanwu GO, Oyeneke EC, Udunobin U, Adegbegi AJ. Impact of *Anthocleista vogelli* root bark ethanolic extract on weight reduction in high carbohydrate diet induced obesity in male wistar rats. African Journal of Biochemistry Research, 2013; 7:225-232.
- Asuzu CU, Nwosu MO. Morphology and Anatomy of *Anthocleista djalonesis*. Nigeria Journal of botany, 2009; 22:103-109.
- Ayodele PO, Okonko IO, Odu NN, Bansa A. Antiviral effect of *Anthocleista nobilis* root extract on the biochemical indices of poultry fowls infected with Newcastle disease virus (NDV). Annals of Biological Research, 2013; 3:20-30.
- Backlund M, Oxelman B, Bremer B. American Journal of Botany, 2000; 87: 1029-1043.
- Brown JH, Leband DN. Species imperilment and spatial patterns of development in the United States. Conservation Biology, 2006; 20(1): 239 – 244.
- Cai YL, Song YC. Adaptive ecology of lianas in Tiantong evergreen broad-leaved forest, Zhejiang, China I. leaf anatomical characters. Acta Phytocology Sinica, 2001; 25: 90–98.
- Chartzoulakis K, Patakas A, Kofidis G, Bosabalidis A, Nastou A. Water stress affects leaf anatomy, gas exchange, water relations and growth of two avocado cultivars. Scientia Horticulturae, 2002; 95: 39–50.
- Christensen JH, Hewitson B, Busuico A, Chen A, Gao X, Held I, Jones R, Kolli RK, Kwon WT, Laprise R, Magana RV, Mearns L, Menendez CG, Raisanen J, Rinke A, Sarra A, Whetton P. Regional Climate Protections. In: S Solomon, D Qin, M Manning, Z Chen, M Marquis, KB Avery, M Tignor, HL Miller, (Eds) Climate change: The Physical Science basis, Contribution of Working group I to the Fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, UK and New York; 2007.

- Donald R, Nelson N, Adger W, Katrina B. Adaptations to environmental change: Contributions of a resilient framework. *Annual Review of Environment and Resources*, 2007; 32: 395-419.
- Dublin- Green CO, Awosika LF, Folorunsho R. Climate variability research activities in Nigeria. Nigerian Institute of Oceanography and marine Research, Victoria Island Lagos, Nigeria; 1999.
- Dubuisson JY, Hennequin S, Bary S, Ebihara A, Boucheron- Dubuisson E. Anatomical diversity and regressive evolution in trichomanoid filmy ferns ( Hymenophyllaceae). A Phylogenetic Approach. *Comptes Rendus Biologies* , 2011; 334: 880-895.
- Edward M, Richardson AJ. The impact of climate change on the phenology of the plankton community and trophic mismatch. *Nature*, 2004; 430:881-884.
- Edwin-Wosu NL, Omara-Achong T. Geographical Distribution of species in the genus *Anthocleista* in the Akpabuyo Tropical Rainforest Nigeria. *International Journal of Agriculture*, 2010; 2 (5):128-134.
- Edwin-Wosu NL, Omara-Achong T, Nkang A. Distribution, Habitat adaptation and conservation as integral approach to protection of *Anthocleista* species in Nigeria's Niger-delta landscape. *Asian Journal of Plant Science and Research*, 2015; 5(2): 17-26.
- Edwin-Wosu NL. Phytomorphological Characterization in relation to intraspecific delimitation among members of the tree species in the genus *Anthocleista* found in parts of Niger Delta Tropical rainforest, Nigeria. *European Journal of Experimental Biology*, 2012; 2(6):1962-1973.
- Edwin-Wosu NL, Ndukwu BC. Biosystematic studies in Loganiaceae (Series 2): Histochemical Localisation of Tannins in Species of *Anthocleista* Found In Parts of Niger Delta Tropical Rainforest of Nigeria. *European Journal of Experimental Biology*, 2012; 2(3): 800-806.
- Edwin-Wosu NL, Ndukwu BC. Biosystematic Studies in Loganiaceae (Series 3): Stomatal Morphology In relation to intraspecific delimitation among members of the tree species in the Genus *Anthocleista* found in parts of Tropical Rainforest in Nigeria. *European Journal of Experimental Biology*, 2012; 2(3): 807-813.
- Edwin-Wosu NL, Wejinya C, Omara-Achong T. Biosystematic studies in the Loganiaceae (series 1): Foliar trichome morphology of Tree Species of *Anthocleista* found in Parts of the Niger Delta, Nigeria. *European Journal of Experimental Biology*, 2012; 2(6):1988-2000.
- Edwin-Wosu NL, Jemilat AI, Harry B, Ette Ette E. The Ecological Dynamics and Trajectories of Bioactive Compounds in Plants of the Genus - *Anthocleista* Found In Parts of Niger Delta Ecological Zone, Nigeria. *Global Journal of Pure and Applied Sciences*, 2017; 23: 5-19. DOI: <https://dx.doi.org/10.4314/gjpas.v23i1.2>
- Edwin-Wosu NL, Okafor AC. Evaluation of diverse potentials in the Genus – *Anthocleista*: A short communication on the panacea towards exploiting and optimizing *Anthocleista* species. *Scientia Africana*, 2017; 16(2): 208 – 217.
- Fei SL, Fang JY, Fan YJ, Zhao KL, Liu XJ, Cui KM. Anatomical characteristics of leaves and woods of *Fagus lucida* and their relationship to ecological factors in Mountain Fanjingshan, Guizhou, China. *Acta Botanica Sinica*, 1999; 41: 1002–1009.
- Fubara DMJ, Teme SC, Mgbeke T, Gobo AET, Abam TKS. Master plan design of flood and erosion control measures in the Niger Delta IFERT Technical Report No 1. 1988.
- Harvey PH, Reader AF, Nee S. Why Ecologists need to be phylogenetically challenged. *Journal of Ecology*, 1995; 83: 535-536.

IUCN Red list categories. IUCN Publications, Switzerland, 1994.

Iwara AI, Gani BS, Njar GN, Deeker TN. *Journal of Agricultural Science*, 2011; 2(20):69-75.

Jegede IA, Kunle OF. Ibrahim JA. *Journal of Medicinal Plants Research*. 2011; 5(26): 6136-6139.

**Comment [n1]:** This reference is not used in the text

- Johansen DA. Plant microtechnique. New York: McGraw-Hill, 1940.
- Johnson CT. The leaf anatomy of *Leptospermum* Forst. (Myrtaceae). Australian Journal of Botany, 1980; 28: 77- 104.
- Keay RWJ. Trees of Nigeria. A revised edition. Clarendon Press Oxford, 1989.
- Knoepp JD, Coleman DA, Crossley JR, Clark JS. Biological indices of soil quality and Ecosystem case Study of their uses. Forest Ecology and Management, 2000; 138: 357-368.
- Kurnk P. Customary water loss and practices: Nigeria, 2004.
- Leeuwenberg AZM, Leehours PW. Taxonomy: In A.J.M Leewenberg. Editor-Engler and Prantis. The Natural Plant Families of Angiosperms. Order Gentiales Family Loganiaceae. 28b (1). Dunker and Humblodt, Berlin. pp 8-96. 1980.
- Li FL, Bao WK. Responses of the Morphological and Anatomical structure of the plant leaf to environmental change. Chinese Bulletin of Botany, 2005; 22: 118–127.
- Luter L, Raphael AO, Galadima A, Okoronkwo MU. Phytochemical screening and antimicrobial activity studies of the root extract of *Anthocleista djalonesis*. International Journal of Chemistry, 2012; 4, 4.
- Mendes MM, Gazarini LC, Rodrigues ML. Acclimation of *Myrtus communis* to contrasting Mediterranean light environments-effects on structure and chemical composition of foliage and plant water relations. Environmental and Experimental Botany Journal, 2001; 45: 165-178.
- Ngbolua K, Mubindikila REN, Mpiana PT, Tshibangu DST, Asande MC, Nzongola WK, Baholy R, Fatiany PR. Phytochemical screening, Antibacterial and Antioxidant activities of *Anthocleista liebrechstiana* originated from Democratic Republic of Congo. Journal of Advancement in Medicinal and Life sciences, 2014; 1: 1-6.
- Niger Delta Regional Development Master Plan. Niger Delta Region Land and People. 99pp. 1998.
- Niklas KJ. The Evolution of Tracheid diameter in early vascular plants and its implication on the hydraulic conductance of primary xylem strand. Evolution, 1985; 39:1110-1122.
- Norman A, Hameed M, Ali Q, Aqeel M. Foliar tissue architectural diversity among three species of genus *Hibiscus* for better adaptability under industrial environment. International Journal of Environmental Science, 2012; 2: 2212- 2222.
- Okiwelu SN, Anyanwu AI. Dictionary of Ecology Conservation and Environmental Science: 60pp. First Published in Port Harcourt, 2003.
- Okoli BE. Anatomical studies in the leaf and protract of *Telferia* Hoker (Curcubitaceae). Fedes Repertorium, 1987; 98:231 -236.
- Rocas G, Scarano FR, Barros CF. Leaf anatomical variation in *Alchornea triplinervia* (Spreng) Mull. Arg. (Euphorbiaceae) under distinct light and soil water regimes. Botany Journal of the Linnean Society, 2001;136:231–238.
- Scheiner SM, Goodnight CJ. The comparison of phenotypic plasticity and genetic variation in population of the grass –*Danthonia spicata*. Evolution, 1984; 845-855.
- Sonibare MA, Soladoye MO, Ekine-Ogulana Y. A chemotaxonomic approach to the alkane content of three species of *Anthocleista*. African Journal of Biotechnology, 2007; 6:1516-1520.
- Stace CA. Plant taxonomy and Biosystematics: Contemporary Biology. Edward Arnold, London, 1991.

- Stemier NC, Koliler N. Effect of land scape pattern on species richness, a modeling approach. *Agriculture, Ecosystem and Environment* 2003; 93: 353 – 361.
- Teme SC. Environmental Peculiarities of the Niger Delta in Petroleum Exploration Operations. In: the National Conference of pipeline Vandalisation and Degradation of the Niger Delta Environment. Rivers State Ministry of Environment and Natural Resources of Collaboration with green House Foundation and B. Jean Communication Limited Port Harcourt, Rivers State. 27th -29th Nov. 2001.
- Townsend P, Joirge S, Pearson RG, Anderson RP, Martinez-Meyer E, Nakamura M, Araujo MB. "Species environment- relationships" Ecological niche and geographical distributions MPB-490. Princeton University Press. p82. 2011.
- Verma PS, Agarwal VK. Principles of Ecology in Environmental Biology. Schand and Company Ltd Ram Nagar, New Delhi-110055. pp 591. 1983.

UNDER PEER REVIEW