

## **Isolation and Cryopreservation of *Toxoplasma gondii* isolates from Cat and Chicken from selected households in Thika region, Kenya**

### **Abstract**

There is dearth of information in Africa regarding the genotypic and phenotypic characteristics of the *Toxoplasma gondii* circulating in domestic cats (*Felis catus*) and the intermediate hosts such as chicken. The situation is compounded by lack of collection of well stored isolates. The present study was aimed at creating a cryobank of *T. gondii* bradyzoites, tachyzoites and oocysts. The parasites were isolated from cats and chicken kept in households in Thika region, Kenya. Eight (8) cat fecal samples positive for *T. gondii* oocysts and 38 chicken brain tissues cysts (bradyzoites) were obtained and used for propagation in mice before cryopreservation. For each sample from the cats, (two donor) BALB/c mice were infected orally with  $1 \times 10^4$ /ml of oocysts. From each chicken sample, (two donor) BALB/c mice were infected intraperitoneally with 20-30 tissue cysts. On the third day after infection, tachyzoites were harvested from peritoneal cavity of one donor mice. The other two infected mice were further monitored for eight weeks, euthanized and the brain tissue harvested for toxoplasma cysts which were purified and cryopreserved. From the mice infected with oocysts from the cats samples, 2 (25%) tachyzoites but a higher 8 (100%) isolation was obtained from brain tissue cysts. On the other hand from chicken samples generated 18 (47.3%) tachyzoites and 38 (100%) tissue cysts were obtained. The isolated oocysts (from cats), tachyzoites and tissue cysts (from mice) were cryopreserved using 15% glycerol as cryoprotectant and stored in liquid nitrogen (-196°C). After 6 months of cryopreservation, the viability of the isolates was tested using Trypan blue dye exclusion on manual haemocytometer. Viability (99.5% - 96%) of the cryopreserved samples was maintained for the three toxoplasma stages and there was no significant change ( $p > 0.05$ ) in viability of the parasites before and after cryopreservation. The cryobank will serve as a repository for subsequent studies on molecular and phenotypic characterization of *T. gondii* isolates from Kenya.

**Key words:** chicken, cat, cryopreservation, tissue cyst, oocyst, tachyzoites, Thika

## **1. Introduction**

*Toxoplasma gondii*, which causes toxoplasmosis, infects all warm-blooded animals with felids family being the definitive host while other animals are intermediate hosts. The parasite is mainly transmitted by fecal-oral, carnivorous, and trans-placental routes. The prevalence of toxoplasmosis in human populations varies according to countries, geographical areas and ethnic groups [1]. A recent review also showed that in most African countries there is high-risk area mainly due to the close association of humans and livestock as well as sociocultural practices, poor environmental hygiene [2]. In Kenya, seroprevalence of up to 60% has been reported in epidemiological in different categories of people [3, 4, 5]. Ongoing studies by the current investigators show that in Kenya, just like in other countries, the risk factors associated with the occurrence of toxoplasmosis include keeping of free-ranging domestic cats and chicken which harbor *T. gondii* oocysts and tissue cysts [6, 7].

The phenotypic and genotypic characteristics of *T. gondii* are critical in determination of pathogenesis, virulence, diagnosis and treatment of the ensuing disease. Felids excrete oocysts which sporulate in the environment. The sporozoites are ingested by intermediate hosts through various modes (drinking water, eating raw vegetables). Human may also get infected by consuming raw or undercooked meat containing tissue cysts stages [8]. The tachyzoites in the intermediate hosts are associated with the acute phase of infection while tissue cyst is the encysted form and contains bradyzoites. The latter is also the terminal stage in intermediate hosts and infective to felids.

Currently, most of the information regarding *T. gondii* phenotypic and genotypic characteristics emanates from studies done in Europe and America [9] with little being documented from African region. Isolation of viable *T. gondii* in laboratory is usually maintained by propagation in mice via serial passages a process that faces key challenges on cost, ethical issues based on laboratory animals use and labor intensity [10]. Cryopreservation of isolates provides an alternative to use of laboratory animals [11]. The aim of this study was to create a cryobank of all infective stages of *T. gondii* isolated in Kenya for further studies.

## **2.0 Materials and methods**

### **2.1. Ethical clearance**

Ethical clearance for the studies involving animals was approved by Institutional Animal Care and Use Committee of the Institute of Primate Research, Nairobi (Ethical clearance certificate IRC/21/11). The study also adhered to the ARRIVE guidelines for reporting in vivo animal experiments.

## **2.2. Study area**

The study was carried out in Thika region; (latitude 1° 4' 60 S 37°, longitude 37° 4' 60 E) located in Central Kenya, Kiambu County and covering an area approximately 1,960Km<sup>2</sup>. The region has a tropical climate with an annual rainfall ranging between 500 mm – 1500 mm and an annual average temperature range of 16.4 to 19.8°C. The main economic activity in the area is livestock agriculture. Most homes in this environment keep domestic cats as pets and curiously/traditionally for controlling rodents [6]. Samples from cats and chicken in selected households in a peri-urban setting were collected from Ruiru, Juja and Thika sub-counties.

## **2.3. Laboratory animals**

BALB/c mice were obtained from Institute of Primate and Research (IPR) -Nairobi, breeding colony. The mice were housed in standard shoe box macron cages in groups of five per cage in a BSL2 laboratory at IPR. They were provided with beddings and other environment enrichment items such as tunnels for burrowing. The mice were fed with mice pellets (Mice pellets<sup>®</sup>, Unga Ltd, Kenya) and provided drinking water *ad libitum* as detailed by IPR guidelines. The ambient room environment temperature was maintained at between 22 and 27<sup>0</sup>C.

## **2.4 Isolation of oocysts and tissue cysts from cats and chicken**

The oocysts were obtained and processed from fecal samples of eight (8) cats as described by [12]. In brief, the oocysts were isolated by sugar flotation technique [13] and observed at magnification of x400. The oocysts were confirmed to be *T. gondii* using a polymerase chain reaction (PCR) targeting 529 bp [12, 14].

For tissue cysts isolation, 48 free range chickens were purchased from local farmers and sacrificed by a registered veterinary Laboratory Technician using cervical dislocation method as described by [15]. The tissue cysts were isolated from 38 (79%, 38/48) of the chicken brain tissue. Briefly, the chicken were sacrificed (within 48 hours) and brain removed as previously

described by [16]. The yield on isolated cysts was enumerated using the manual counting method in a haemocytometer [12].

### ***2.5. Propagation of *T. gondii* in mice***

Sporulated oocysts ( $10^4$  in 0.2 ml) from each fecal sample were administered (for expansion/multiplication) by oral gavage into two sets of BALB/c mice per sample. After three days of inoculation the first group of mice was sacrificed using standard CO<sub>2</sub> euthanasia. The tachyzoites were harvested from peritoneal cavity of the mouse using the method described by [17]. The numbers of tachyzoites were determined by manual counting on a haemo-cytometer. The second group of inoculated mice was monitored for eight weeks after which tissue cysts from brain were harvested.

The process was repeated for cysts obtained from chicken brain. In this case, the cysts were diluted using phosphate saline glucose (PSG) into 20-30 cysts per 0.2 ml. The latter dose was injected intraperitoneally into two sets of BALB/c per sample of oocysts from the chicken and harvested as described by [18].

### ***2.6. Cryopreservation of isolated parasites***

A 500µl suspension of oocysts, tachyzoites, and tissue cysts containing bradyzoites were mixed with an equal amount of 500µl 15% glycerol. Aliquots of 200µl for tachyzoites and oocysts and 500 µl for tissue cysts were cryopreserved. The aliquots were cooled in a stepwise manner at -20°C for 4hrs, -80°C overnight and then transferred into liquid nitrogen (-196°C) as described by [19].

### ***2.7. Viability testing***

The cryopreserved tachyzoites, tissue cysts and oocysts were retrieved from liquid nitrogen after 6 months of storage. Following rapid thawing at 37°C in a water bath, samples were then diluted 1:5 in RPMI with 5% FCS, centrifuged and suspended in PBS for viability assessment. Further, the samples were diluted in sterile filtered Trypan Blue dye by preparing 1:1 dilution of the cell suspension using 0.4% Trypan Blue solution. 20µl was loaded on haemocytometer and observed under x 40 eye piece of light microscope live/dead parasites quantified. Motility of tachyzoites was ~~used~~ also used as a confirmatory test to assess their viability.

### ***2.8. Data analysis***

Data was managed and analysed using Ms Excel<sup>®</sup> software (Microsoft, USA). Paired ANOVA test was used to correlate viability of *T. gondii* oocysts, tissue cysts and tachyzoites, before and after cryopreservation.

### 3.0 Results

#### 3.1 Proportion of tachyzoites and tissue cysts harvested from mice

*Toxoplasma gondii* tachyzoites were isolated from 25% (2/8) cat faecal samples. Further, all (100%) mice infected with oocysts survived for the eight (8) weeks after infection. The mice remained asymptomatic throughout the eight weeks of monitoring. However, brain tissue cysts containing bradyzoites were obtained in all eight (8) study mice.

Following infection of the 38 *T. gondii* from chicken tissue samples into BALB/C mice, 18 (47.3%) tachyzoites. The mice remained asymptomatic throughout the eight weeks of monitoring but all 38 (100%) samples from chicken brain cyst led to brain tissue cysts in the infected mice.

#### 3.2 Viability of cryopreserved *T. gondii* isolates

After the cryopreservation for 6 months of the three stages of the parasites the viability of the isolates was done and the results are shown in Table 1.

The cryopreserved parasites (oocysts, bradyzoites and tachyzoites) were not morphologically distinguishable from the fresh ones. The percentage of surviving parasites was similar and high ranging from 96-99.5%. The viability was highest for the oocysts (99.5%). There was no significant ( $p>0.05$ ) difference in viability of the parasites before and after cryopreservation (Table 1).

**Table 1: Viability of *Toxoplasma gondii* after six months cryopreservation**

Parasite Stage	Original host	Mean No. (Parasites/ml) at storage	Mean No. (Parasites/ml) after 6 months cryopreservation	Viability (%) after 6 months cryopreservation
Oocysts	Cat	$7.25 \times 10^2$	$7.21 \times 10^2$	99.5
Tachyzoites	Cat	$2.49 \times 10^7$	$2.49 \times 10^7$	97

Tissue cysts	Cat	$1.60 \times 10^4$	$1.60 \times 10^4$	96
Tissue cysts	Chicken	$1.05 \times 10^7$	$1.05 \times 10^7$	97

#### 4.0 Discussion

Toxoplasmosis is a neglected disease affecting both animals and public health with high risk of morbidity in developing countries [2]. In Africa, limitations in addressing gaps in management of toxoplasmosis study are hindered by lack of repertoire of preserved isolates. This study was geared towards isolation, propagation of isolates collected from cats and chicken hosts from an area known to be endemic of the disease [5, 7].

In the present study, *T. gondii* oocysts from cats were able to generate tachyzoites and brain tissue cysts. However, although all mice injected with oocysts developed the brain tissue cysts, the harvested tachyzoites from peritoneal cavity was quite low. Similarly, the tissue cysts obtained from the chicken also generated fewer tachyzoites than brain tissue cysts in the mice. This could be due to low sensitivity of the peritoneal harvesting method in obtaining all the tachyzoites from the mice. Previous studies have shown tachyzoite formation in peritoneal cavity ranges from a few hours to four days with non-virulent strains taking a longer period than virulent one [20]. Studies done on *T. gondii* cultivation indicate variation in tachyzoite formation depend largely on the genotype of the isolate; the strain of mice used and route of infection are crucial factors to the outcome of infection [13, 21]. In the current study, oocysts generated fewer (25%) tachyzoites compared to tissue cysts from chicken which generated 47.3% tachyzoites in the mice. This could be due to differences in the host source, genotype of parasite and route of infection. Further studies should be undertaken to determine the genotypes of these isolates.

Once the parasites multiply in epithelial cells of small intestine and thereafter in the peritoneal cavity, *T.gondii* have a tropism for other cells with the brain being main target. Thus, in the current study all mice infected either by oocysts or tissue cysts from chicken developed brain tissue cysts which had bradyzoites. This is similar to previous studies which have shown that *T. gondii* primarily targets brain cells ~~such as~~ [22, 23].

Cryopreservation, which was used in this study, has the advantage of maintaining the strains in their original state and not modified as it is common during continuous laboratory animal passage. The latter is costly and laborious [11]. The current study showed that most of the oocysts, tachyzoites and tissue cysts containing bradyzoites were able to maintain their morphology after cryopreservation. This shows that cryopreservation did not significantly affect the integrity of the parasites and could thus produce experimental infections. Higher cryopreservation success (over 90%) was noted in this study. Thus, cryopreservation which was done using 15% glycerol can be used during long term storage of all the three stages of *T. gondii* parasites. Previous studies have shown that glycerol can be used as a cryoprotectant for other cells such as spermatozoa, other mammalian cells and parasites including *Entamoeba histolytica*, *Trichomonas vaginalis*, *Leishmania* spp, *Trypanosoma* spp. [19, 24, 25]. Cryopreservation creates a dehydrating environment during freezing thus preserving the cells from cryo-injuries. Glycerol was used in this study is known to form hydrogen bonds with water molecules thus preventing ice crystals formation which damage cells [26].

Availability of repertoire of isolates of viable *T. gondii* infective stages from developing countries like Kenya will indeed provide a platform to study the relationship between the genotypic and phenotypic characteristics of local isolates of *T. gondii*. Currently most of the information available on *T. gondii* is from Europe, Asia, and America with little documentation from African countries. The cryopreservation method of storage of parasites is able save laboratory workers a considerable amount of the time, effort, and expense involved in frequent serial passage of the parasite in laboratory rodents. Maintenance of *T. gondii* in laboratory

animals has been associated with genome changes characterized by loss of ability to form tachyzoites, gametocyte-forming ability may be lost, and that selection of some mutations by the host animal or growth system was evident [27].

In conclusion this study has been able to isolate and cryopreserve *T. gondii* isolates from cats and chicken from Kenya. The genotypic and phenotypic characterization of these isolates will ensue with an intention of determining their relatedness with isolates from other continents and impact on disease pathogenesis and management.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### **References**

1. G. Pappas, N. Rousso and M.E. Falagas. "Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis," International Journal for Parasitology, Vol.39, pp. 1385–1394, 2009.
2. J.M. Mose, J.M. Kagira, D.M. Kamau, N.W. Maina, N. Maina, and S.M. Karanja, "A review on the present advances on studies of toxoplasmosis in eastern Africa," BioMedical Research International, Vol. 2020, article ID.7135268, pp. 12, 2020.
3. R.H. Cocker, "An immunological survey of toxoplasmosis in a rural Kenyan population" M. Med Dissertation in Human Pathology. *University of Nairobi, Kenya*, 1991.

4. T.R. Bowry, M.E. Camargo, M. Kinyanjui, "Sero-epidemiology of *Toxoplasma gondii* infection in young children in Nairobi, Kenya," Transactions of the Royal Society of Tropical Medicine and Hygiene, Vol. 80.(3), pp. 439-441, 1986.
5. S.K. Thiong'o, J.M. Ichagichu, M. Ngotho, G.O. Aboge, J.M. Kagira, S.M. Karanja, N.N. Maina, "Use of the nested polymerase chain reaction for detection of *Toxoplasma gondii* in slaughterhouse workers in Thika District, Kenya," *South African Medical Journal*, Vol:106 (4), pp. 417-419, 2016.
6. E. Ogendi, N. Maina, J. Kagira, M. Ngotho, G. Mbugua and S. Karanja, "Questionnaire survey on the occurrence of risk factors for *Toxoplasma gondii* infection amongst farmers in Thika District, Kenya," *Journal of the South African Veterinary Association*, Vol.84, (1), pp.191-196, 2013.
7. J.M. Mose, D.M. Kamau, J.M. Kagira, N.M. Maina, M. Ngotho, A.N. Njuguna and S.M. Karanja, "Development of neurological mouse model for toxoplasmosis using *Toxoplasma gondii* isolated from chicken in Kenya," *Pathology Research International*, Vol. 2017, 2017.
8. M.Louis, Weiss and K. Kami, "*Toxoplasma gondii*," Second Edition. Elsevier Ltd publishers, 2014.
9. J.P. Dubey, D.H. Graham, E. Dahl, M. Hilali, A. El-Ghaysh, C. Sreekumar, O.C. Kwok, S.K. Shen and T. Lehmann, "Isolation and molecular characterization of *Toxoplasma gondii* from chickens and ducks from Egypt," *Veterinary parasitology*, Vol.30(114), pp. 89-95, 2003.
10. Z. Huanqin, C. Ying, L. Fangli, L. Man, Y. Xiaoyan, F. Xiaoyin, Z. Ying, H. Bo, H. Shiguang, and H. Lloyd, "Cryopreservation of *Toxoplasma gondii* in infected murine tissues," *Parasitology Research*, Vol.111(6), pp. 2449-2453, 2012.
11. M. Döşkaya, A. Caner, H. Can, S.G. İz, A. Değirmenci, A.Y. Gürüz, "Cryopreservation of *Toxoplasma gondii* tachyzoites and tissue cysts" *Turkish Journal of Parasitology*, Vol. 37(1), pp. 44-46, 2013.
12. A.N. Njuguna, J.M. Kagira, S.M. Karanja, M. Ngotho, L. Mutharia, N.M. Maina, "Prevalence of *Toxoplasma gondii* and Other Gastrointestinal Parasites in Domestic Cats

- from Households in Thika Region, Kenya,” *BioMed Research International*, Vol. 7615810, pp. 6 <https://doi.org/10.1155/2017/7615810>, 2017.
13. J.P. Dubey, “Oocyst shedding by cats fed isolated bradyzoites and comparison of infectivity of bradyzoites of the VEG strain *Toxoplasma gondii* to cats and mice” *Journal of Parasitology*, Vol. **87**(1), pp. 215-219, 2001.
  14. Q. Kong, S. Lu, Q. Tong, D. Lou, C. Rui, Z. Bin, K. Takashi, W. Li-Yong, O. Nobuo and X. Zhou, “Loop-mediated isothermal amplification (LAMP): Early detection of *Toxoplasma gondii* infection in mice,” *Parasites and Vectors*, vol. 5(2), 2012.
  15. AVMA, “2000 report AVMA panel on euthanasia,” *Journal of the American Veterinary Medical Association*, Vol. 218(5), pp. 669-696,
  16. S. Carlos and R. Jack, “Animal Models for *Toxoplasma gondii* Infection,” *Current Protocols in Immunology*, Vol.19(3), pp. 1-19, 1998.
  17. A. Ray, and B.N. Dittel, “Isolation of Mouse Peritoneal Cavity Cells,” *Jove*, Vol. 35(10), 2010.
  18. J.P. Dubey and C.P. Beattie, “*Toxoplasmosis of animals and man*”. CRC Press, Boca Raton, Florida, pp. 220, 1988.
  19. D.B. Lin, K.E. Su, J.C. Yu, “Studies on cryopreservation of *Toxoplasma gondii* and its antigenicity in mice,” *American Journal of Tropical Medicine and Hygiene*, Vol. 53(4), pp.392–396, 1995.
  20. J.P. Dubey, “Bradyzoite-induced murine toxoplasmosis: stage conversion, pathogenesis, and tissue cyst formation in mice fed bradyzoites of different strains of *Toxoplasma gondii*,” *Journal of Eukaryotic Microbiology*, Vol. 44(6), pp. 592–602, 1997.
  21. A.M. Johnson, “Strain dependent route of challenge dependent, murine susceptibility to toxoplasmosis,” *Zeitschrift fur Parasitenkunde*, Vol.70 (3), pp. 303 – 309, 1984.
  22. C.M. Cabral, S. Tuladhar, H.K. Dietrich, E. Nguyen, W.R. MacDonald, T Trivedi, A Devineni and A.A. Koshy, “Neurons are the primary target cell for the brain-tropic intracellular parasite *Toxoplasma gondii*,” *PLoS Pathogen*, Vol. 12(2), 2016.
  23. J.P. Dubey, L.R. Ferreira, M. Alsaad, S.K. Verma, D.A. Alves, G.N. Holland, “Experimental toxoplasmosis in rats induced orally with eleven strains of *Toxoplasma*

*gondii* of seven genotypes: tissue tropism, tissue cyst size, neural lesions, tissue Cyst rupture without reactivation, and ocular Lesions” PLoS ONE, Vol. 11(5), 2016.

24. A.V. Bomhard, A. Elsässer, L. Ritschl, M.S. Schwarz and N. Rotter, “Cryopreservation of endothelial cells in various cryoprotective agents and media - vitrification versus Slow freezing methods,” PloS one, Vol. 11(2), 2016.
25. M. Yuko, K. Panagiotis, U. Shoji, “Cryopreservation of protozoan parasites,” *Cryobiology*, Vol. 48(1), pp. 1-7, 2003.
26. S. Bhattacharya, “Cryoprotectants and Their Usage in Cryopreservation Process,” Intechopen, 2018.
27. I. Literak and I. Rychlik, “Genome changes in the *Toxoplasma gondii* strains during laboratory passages in mice,” Journal of the University of Veterinary and Pharmaceutical Sciences in Brno, Vol. 68(3), pp. 203–208, 1999.

UNDER PEER REVIEW