

## COMPARATIVE EFFECTIVENESS OF WILD AND COMMERCIAL STRAINS OF LACTIC ACID BACTERIA STARTER CULTURE DURING BACKSLOPING FOR YOGHURT PRODUCTION

### Abstract

Yogurt is one of the traditionally fermented dairy products which are usually prepared with lactic acid bacteria cultures containing *Streptococcus thermophilus* and *Lactobacillus bulgaricus* and has gained widespread consumer acceptance as a healthy diet. The roles of exotic starter cultures for dairy milk processing in Nigeria food supply chain cannot be overstated. However, there are concerns over growing food and food-related import dependency in the sub-African countries and Nigeria in particular which have increased during the last few years. The selection of indigenous strains with capacity for backsloping in yoghurt production might reduce importation and improve economy in a way. Yoghurt fermentation was carried out by backsloping method using three commercial (Exotic) formulated cultures of *Lactobacillus bulgaricus* and *Streptococcus thermophilus*; ES07, ES32 and ES14 respectively as well as wild (isolated) formulated culture strains; WS11, WS65 and WS13 respectively and were used to produce yoghurt using backsloping method. Their pH, coagulation, microbial profile and sensory attributes were evaluated after three consecutive batch of backsloping. The overall sensory attributes of yoghurts produced using exotic compared with indigenous strains showed no significant ( $p > 0.05$ ) difference. There was also no significant ( $p < 0.05$ ) variation for all set of yoghurts in term of acidification and coagulation ability. The pH of fermented products using commercial cultures ranged from 4.53-5.32, total plate count  $1.1 \times 10^6$ -  $4.0 \times 10^7$  cfu/ml during the 10 hours fermentation. While pH of yoghurts produced using isolated cultures ranged from 4.35 – 4.72, total plate count  $2.7 \times 10^7$ -  $2.3 \times 10^8$  cfu/ml during the 10 hour fermentation times. The overall acceptability were in the decreasing order; WS65>WS11>ES07>WS13>ES14>ES32. Starter culture formulation using indigenous strains could be further developed genetically for commercial applications because they compete favourably with their exotic counterparts.

**Keywords:** Acidification, Backsloping, Cultures, Importation and Indigenous

### Introduction

There is high demand for yoghurt and fermented milk owing to their numerous health benefits. In recent years, more people are getting aware of the health protections that could be derived from the consumption of controlled-fermented products such as yoghurt. (Bristone *et al.*, 2015). For centuries, the health benefits of the consumption of fermented milk such as yoghurt with its lactic acid bacteria contents reducing the incidence of intestinal toxin, virginal infections, blood pressure and osteoporosis have been reported (Shahnawaz *et al.*, 2013). Fermented products like yoghurt have been known to contain elements with some therapeutic and health maintenance properties (Igbadun *et al.*, 2014). Aside the above mentioned benefits, the presence of *S. thermophilus* and *L. bulgaricus* as the culture organisms in the fermented products like yogurt means the availability of functional probiotic properties (Aryana *et al.*, 2017). The use of Lactic Acid Bacteria strains in dairy starter cultures is highly recommended because of their safety considerations (GRAS status) (Rycken *et al.*, 2018).

Some studies have shown that another beneficial property of LAB is their possession of biomolecules with autophagy triggering capacity which are said to be important in cell recycling for antimicrobial protection and maintenance of epithelial competence (Dinić *et al.*, 2017). Although LAB is wide spread in nature, they are commonly associated with dairy materials where they mostly play the role of acid fermentation. These functions have been harnessed for the commercial production of many dairy products with good sensory characteristics for the benefit of mankind (Bluma *et al.*, 2017). Indigenous Lactic acid bacteria strain isolated from the spontaneous fermented milk referred to as the 'wild' have specific properties that can be harnessed for controlled fermentation of milk (Bluma *et al.*, 2017).

Considering the limited availability of exotic commercial starter cultures in Nigeria, there has been a trend of selection and development of new starter cultures from the

autochthonous habitat for development into viable commercial status with potent traits for backslipping applications and stable shelf life. The effort at using such newly-isolated and well-characterized cultures for use in place of the exotic strains will lead to the production of standard products with predictable characteristics as their exotic peers in terms of both nutrition enhancement and health promoting features.

There is currently limited documentation of the technological consequences of the different traditional techniques employing backslipping for producing controlled fermented products such as yoghurt.

Therefore, the aim of this work was to determine and document the comparative differences between backslipping using commercial strains and those of the isolated cultures during yoghurt production practice. Such indigenous knowledge in traditional fermentation practices utilising backslipping method may enhance further development, up-scaling and utilization of local resources to the level of other renowned fermented milk products.

## Materials and Methods

### Preparation of Glass Wares and Media

Glass wares were oven sterilized by dry heat (Medified Equipment and Scientific DHG-9023A, England) at 160°C for 60 minutes as described by Fawole (1988) and Cheesbrough (2000). Each microbial medium was prepared by dissolving the appropriate powder in 1000 ml (1 litre) of distilled water as described by the manufacturers. Each was autoclaved at 121°C for 15 minutes (Fawole, 1988; Jideani and Jideani, 2006).

### Sample collection

Commercial starter cultures for this research were obtained from standard supermarkets within Abuja metropolis and maintained at 4°C until needed. The locally fermented milk product (Nunu) for the isolation of indigenous lactic acid bacteria strains were collected from the households processing them from around the FCT. The samples were collected in sterile bottles and kept cool until they arrived at the laboratory, where they were kept at 4°C for further use.

### Isolation of Lactic Acid Bacteria

One (1ml) ml aliquot was taken from the collected samples aseptically after thorough homogenization. Serial dilution was then made using sterile peptone water by adding 1ml into 9ml. A 0.1ml aliquot of  $10^{-2}$  and  $10^{-4}$  of the serially diluted samples were aseptically added to M17 and MRS agar for the isolation of LAB (Badis *et al.*, 2004a). In order to prevent the growth of yeast, the agar media were supplemented with  $100\text{mg}^{-1}$  of cycloheximide and were incubated for 5 days at 40°C anaerobically using the Gas Pack system (Merck Anaerocult type A) (Kalavrouzioti *et al.*, 2005). After incubation, discrete colonies were randomly selected and purified on sterile media. The resulting pure strains were kept in two different conditions including at 4°C for MRS and M17-plates and at -20°C for M17 and MRS broths supplemented with 20% glycerol for further use (Mathara *et al.*, 2004).

### Identification of the Bacterial Strains

All the purified bacteria strains were tested for gram reaction, catalase production and spore formation (Harrigan and McCance, 1976). Colonies were characterized on MRS and M17 agar. Only strains with gram positive and catalase negative reactions were finally used for further subjected to further identification (Sharpe, 1979). Growth at different temperatures (10, 15, 37, 40 and 45°C) for 5 days, resistance to 63°C for 30 min (Sherman test), growth in the presence of 2, 3, 4 and 6.5% NaCl and different pHs (4.5 and 6.5) were used for further identification of the strains. Ability to hydrolyse arginine and asculin, utilization of citrate, production of acetone, gas formation from glucose and dextran production from sucrose were also tested according to the method of Samelis *et al.*, 1994. The strains were also tested for fermentation of L-arabinose, D-xylose, galactose, D-fructose, sorbitol, lactose, melibiose, saccharose, D-raffinose, melezitose, mannose and glucose (Tserovska *et al.*, 2002). The growth of bacterial strains at 10, 15, 37, 40 and 45°C was visually confirmed by the changes in turbidity of MRS or M17 broth after 24, 48 and 72 h of incubation. The tolerance of microorganisms to the different levels of salt, pH and heat (60°C) was also visually evaluated. Arginine dihydrolase agar and asculin azid agar (Merck, Germany) were employed to perform the hydrolysis tests. For evaluation of citrate utilization and acetone production, citrate and MR-VP agars (Merck, Germany) were used. MRS or M17 broths containing inverted Durham tubes were used for evaluation of gas production and the production of dextran from sucrose was done in MRS agar (Mayeux *et al.*, 1962).

### Preservation of LAB Strains

In order to prevent loss of properties, the identified strains were stored in skimmed with 30% (v/v) glycerol at 4°C till further use (Badis *et al.*, 2004). Cultures were also kept on MRS agar or M17 agar slant at 4°C and streaked forth-nightly.

## Technological Characterization

### Acidifying activity

The acid-production potential of the strains was measured according to the International Dairy Federation (IDF) standard 306 (Kihal *et al.*, 1996). Fermentation was carried out at

need  
space

need  
a  
space

delete the underline

40°C by preparing 10% milk and introducing 24 hour old cultures at 1% rate. The pH was measured during the 12 hour fermentation period by inserting the cleaned probe of the pH meter in the fermenting medium and determining the values..

**Proteolytic Activity**

The proteolytic activity of the isolates were determined by fermenting 10% skimmed milk for 12 hours at 40°C and recording the coagulation which is either positive (+) or negative (-).

**Biomass production**

The bacteria strains were sub-cultured into MRS broth (100ml) of the medium with 10% of the active culture. The respective growth was monitored by measuring the Optical Density at 600nm (OD600) using a spectrophotometer. The biomass was determined by centrifuging aliquot samples. The dry weight was determined as needed by drying the pellet of samples at 55°C for 24h (Ayad *et al.*, 2004).

**Experimental Design for Yoghurt Production**

In order to fully attain the goal of this research, two sets of milk fermentation experiments were set-up in parallel; one used commercial starter culture strains, viz; co-culture samples of *Lactobacillus bulgaricus* and *Streptococcus thermophilus* (ES07, ES32 and ES14) respectively while the other one used the isolated and purified LAB strains, (WS11, WS65 and WS13) respectively.

The freeze dried samples were prepared from the commercial strains by weigh 0.5g of the lyophilized cultures and dissolved in 5ml of the pasteurized milk. The culture was incubated at 40°C for 18h. Three sets of reconstituted skimmed milk (10%) were homogenised and pasteurized at 65°C for 30 minutes and cooled to 42°C. Each set was inoculated with about 5ml (containing 10<sup>6</sup> CFU/mL) commercial, prepared starter cultures (ES07, ES32 and ES14).

Five (5ml) of isolated cultures (WS11, WS65 and WS13) containing 10<sup>6</sup> CFU/ml which were determined by the initial viable counts. All the set-up was incubated at 40°C until the pH reached 4.5 then kept in the refrigerators set at 4°C. Sample from the fermented samples were used to initiate consecutive batch (2.5%) to get a three-fold sample. During fermentation, the pH, coagulation, aerobic total plate count, flavour and organoleptic properties were measured for each set of the sample. The third-fold fermented sample were compared.

**Organoleptic Analysis**

This analysis was carried out by ten taste panellists who were trained to identify the characteristics of yoghurt without sugar added. They were served with the chilled samples in coded cups and the respective records were made after tasting. The panellists were asked to rank the coded samples on the basis of their quality attributes made up of; consistency, flavour, taste, colour, texture and overall acceptance using nine-point hedonic scale, expressing degree of liking or disliking as in the questionnaire (Igbadul *et al.*, 2014).

= 2210 symbol

remove space  
if small letter

**Results**

**Table 1: pH, Proteolytic ability and Total Plate Count of the third-fold fermented Yoghurt using Commercial and Isolated LAB strains**

Parameter	Commercial Cultures			Isolated Cultures		
	ES07	ES32	ES14	WS11	WS65	WS13
pH (EPV)	4.53	5.32	5.21	4.42	4.35	4.72

organise the table's components

Proteolysis      +      +      +      ++      ++      +


Total Plate Count       $4.0 \times 10^7$      $3.4 \times 10^6$      $1.1 \times 10^6$        $2.3 \times 10^8$      $1.2 \times 10^8$      $2.7 \times 10^7$

EPV = End point value; += Coagulation

*^ = space bar*

Table 1 shows the pH, Proteolytic ability and total plate count of both the commercial and isolated LAB strains recorded at the end of a third-fold milk fermentation (Backsloping). The lowest pH (4.35) was recorded in the yoghurt culture fermented with isolated lactic acid bacteria culture (WS65) while the highest pH (5.32) was recorded in yoghurt fermented with bacteria culture (5.32). Proteolytic activity (Coagulation rate) occurred fastest in yoghurt products produced with cultures WS65 and WS11. Total aerobic plate count of yoghurt culture were recorded overall higher in yoghurt cultures fermented with isolated lactic bacteria strains than in the commercial strains. The highest being  $2.3 \times 10^8$  CFU/mL (WS11) while the lowest was  $1.1 \times 10^6$  CFU/mL (ES14).

*the graph should be only shown*



*why attach the excel style*

Fig.1: The Percentages Acceptance of the Yoghurt Samples Produced by the Respective Cultures

Figure 1 shows the mean percentage acceptance of the yoghurt produced by the various lactic acid bacteria cultures. The highest being WS65 (84%) while the lowest was ES32 (50%).

*word*



*picture was distorted not the common right picture*

A. Soft, smooth, pudding-like, firm textured Product- Consistent Throughout Backsloping Batches.

*remove bold and capitalized*



**Fig.2: COMPARATIVE BACKSLOPING ABILITY OF SOME LACTIC ACID BACTERIA STRAINS**

the author should not compress the picture it is distorted

### Discussions

The average acidification activity of the isolated lactic acid bacteria culture was better than their commercial counterparts. As a result, there was a faster coagulation of pasteurised milk protein thereby producing yoghurt with better quality as elucidated by the panelists' results. This is in agreement with the reported work on other product but same principle by Holzapfel (2002) that various carriers such as porous material of gourd have been used to achieve backsloping. That 'inoculation belt' was used for the initiation of *pito* beer fermentation in work carried out in Ghana. And that in Egypt *khisk* is prepared using cow/camel milk and wheat where one-third (w/w) of dried *khisk* is used as a starter inoculum. These practices in the preparation, preservation and distribution of inoculums according to the research could be used as a basis for the development of starters for other fermented foods.

Also, the high acid production means that the product could be better preserved thereby giving it longer shelf life because acid content of food matrix reduce the population spoilage microorganisms. It is said that within a spoiling food, there is often a succession of different populations that rise and fall as different nutrients become available or are exhausted. Notably, some microbes, such as lactic acid bacteria, secrete compounds such as acid that inhibit competitors (Gram, *et al.*, 2002). The aerobic total plate count recorded higher values in products fermented with indigenous strains. The implication is that such strains could be better as probiotic cultures since they would be delivered into the gastrointestinal tract in larger concentrations resulting in better and more colonization for enhanced activity. According to the Food and Agriculture Organisation of the United Nations (FAO) and the World Health Organisation (WHO), (2014), probiotics are defined as live microorganisms, which when administered in adequate amounts, confer a health benefit on the host.

your thought should have other support references

### Conclusions

From the data and observations recorded from our study, it can be concluded that backsloping method could be used for large-scale production of yoghurt because there was no significant difference in the performance of the backsloped lactic acid bacteria cultures as compared with the commercial strains. In fact, the yoghurts produced with indigenous strains showed better overall acceptance over the ones produced with commercial strains.

### Recommendations

This old traditional method of fermentation should be employed/upscaled for large scale yoghurt production without delay and with utmost assurance of predictable success.

### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company

rather it was funded by personal efforts of the authors.

## References

Aryana K.J., Olson D.W. (2017): A 100-Year Review: Yogurt and other cultured dairy products. *J. Dairy Sci.* 2017;100:9987–10013.

[Ayad EHE, Nashat S, El-Sadek N, Metwaly H, El-Soda M \(2004\) Selection of wild lactic acid bacteria isolated from traditional Egyptian dairy products according to production and technological criteria. \*Food Microbiology\* 21: 715-725.](#)

Badis, A., D. Guetarni, B. Moussa-Boudjema, D.E. Henni, M.E. Tornadijo and M. Kihal, 2004. Identification of cultivable lactic acid bacteria isolated from Algerian raw goat's milk and evaluation of their technological properties. *Food Microbiol.*, 21: 343-349.

Bluma A., Ciprova I., Sabovics M. The influence of non-starter lactic acid bacteria on swiss-type cheese quality. *Agric. Food.* 2017;5:34–41.

Bristone C, Badau MH, Igwebuike JU, Igwegbe AO (2015). Production and evaluation of yoghurt from mixtures of cow milk, milk extract from soybeans and tiger nut. *World J. Dairy and Food Sci*, 10(2), 159 - 169.

Dinić M., Lukić J., Djokić J., Milenković M., Strahinić I., Golić N., Begović J. *Lactobacillus fermentum* Postbiotic-induced Autophagy as Potential Approach for Treatment of Acetaminophen Hepatotoxicity. *Front. Microbiol.* 2017;8:594.

Fawole, O. O.\* , Oyelese, O. A. and Etim, E. U. (2018): ORGANOLEPTIC AND CHEMICAL ASSESSMENT OF TWO FROZEN MARINE FISHES OBTAINED FROM MARKETS IN FOUR AGRICULTURAL ZONES OF OYO STATE, NIGERIA; *Ife Journal of Science* vol. 20, no. 2 (2018).

Food and Agriculture Organization of the United Nations/World Health Organization FAO/WHO Health and Nutritional Properties of Probiotics in Food including Powder Milk with Live Lactic Acid Bacteria. 2001. [(accessed on 20 January 2014)]

Gram, L. Ravn, M. Rasch, J.B. Bruhn, A.B. Christensen and M. Givskov. *Int. J. Food Microbiol.*, 2002, 78: 79–97.

Harrigan, W.F. and M.E. MaCance, 1976. *Laboratory Methods in Food and Dairy Microbiology*. Revised Edn., Academic Press, New York, pp: 33-200

Igbadul B, Shember J, Amove J (2014). Physicochemical, microbiological and sensory evaluation of yoghurt sold in Makurdi metropolis. *African J, Food Sci. and Tech.* 5(6), 129 - 135.

Holzappel WH, Appropriate starter culture technologies for small-scale fermentation in developing countries. *Int J Food Microbiol* 75: 197– 212(2002).

Johansen E. Use of Natural Selection and Evolution to Develop New Starter Cultures for Fermented Foods. *Annu. Rev. Food Sci. Technol.* 2018;9:411–428.

Kalavrouzioti, I., M. Hatzikamari, E. Litopoulou-Tzanetaki and N. Tzanetakis, 2005. Production of hard cheese from caprine milk by the use of two types of probiotic cultures as adjuncts. *Int. J. Dairy Technol.*, 58: 30-38.

[Kihal M, Prevost H, Lhotte ME, Huang DQ, Diviès C \(1996\) Instability of plasmid-encoded citrate permease in J Appl Microbiol 22: 219-223.](#)

Mathara, J.M., U. Schillinger, P.M. Kutima, S.K. Mbugua and W.H. Holzappel, 2004. Isolation, identification and characterization of the dominant microorganisms of *kule naoto*: The Maasai traditional fermented milk in Kenya. *Int. J. Food Microbiol.*, 94: 267-278.

Mayeux, J.V., W.W.E. Sandine and P.R. Elliker, 1962. A selective medium for detecting *Leuconostoc* organisms in mixed strain starter cultures. *J. Dairy Sci.*, 45: 655-656.

Rychen G., Aquilina G., Azimonti G., Bampidis V., Bastos M.d.L., Bories G., Chesson A., Cocconcelli P.S. (2018): Flachowsky G., Gropp J., et al. Guidance on the characterisation of microorganisms used as feed additives or as production organisms. *EFSA J.* 2018;16:e05206.

Samelis, J., F. Maurogenakis and J. Metaxopoulos, 1994. Characterisation of lactic acid bacteria isolated from naturally fermented Greek dry salami. *Int. J. Food Microbiol.*, 23: 179-196.

Shahnawaz M, Sheikh SH, Akbar ZA (2013). Physicochemical analysis of desi yoghurts produced by the local community in Gilgit District. *African J. Food Sci.* 7(7), 183 - 185.

Sharpe, M.E., 1979. Identification of the Lactic Acid Bacteria. In: *Identification Methods for Microbiologists*, Skinner, F.A. and D.W. Lovelock (Eds.). Academic Press, London, pp: 233-259

Tserovska, L., S. Stefanova and T. Yordanova, 2002. Identification of lactic acid bacteria isolated from Katyk, goats milk and Cheese. *J. Cult. Collect.*, 3: 48-52.