

Original Research Article

CHARACTERIZATION OF ACETIC ACID BACTERIA ISOLATED FROM SOME SELECTED WASTE PROTEIN SOURCE

ABSTRACT

Background: AABs are an essential and diverse group of bacteria involved in producing fermented foods and beverages, primarily known for their production of acetic acid (ethanoic acid) used in [the production of vinegar-making](#).

Aim: This study aims to isolate Acetic Acid Bacteria from protein waste and quantify the acetic acid produced; it is research carried out as a means of protein waste management/recycling to conserve the environment/society at large while producing the value-added product; acetic acid from waste protein sources (fish, milk, meat, egg, soy milk, beans and unpasteurized milk).

Methodology: The samples were homogenized in sterile distilled water, followed by serial dilution and inoculation on GYC agar using the spread plate method with an inoculum size of 0.1 ml. The plates were incubated at 30°C for 48 hours. Several biochemical tests, including oxidase, catalase, and Gram stain, were ~~carried out done~~ on the isolates. The acetic acid production rate was estimated using the YGEA medium.

Results: Twenty-five (25) (white, creamy and yellow pale-coloured) well-isolated colonies were selected to be analyzed. Out of 25 isolates, fourteen (14) suspected acetic acid bacteria because they produced clear zones around their colonies in the selective GYC culture medium. Of the 14 analyzed isolates, six (6) were Gram-negative rods. Presumptive identification of the acetic acid bacterial isolates included *Acetobacter aceti* and *Gluconobacter oxydans*. Morphologically identified and biochemically confirmed acetic acid bacterial isolates were then further checked for their abilities to produce acetic acid by titration in YGEA (Yeast, Glucose, Ethanol, Acetic acid) medium. Acetic acid production rate varied among the isolates. Among three (3) acetic acid-producing strains, isolates from fish 2 showed the highest acetic acid production (24.34 g/100ml) rate at 37°C, while those from milk 1 and meat 2 showed 23.96 g/100ml and 23.46 g/100ml, respectively.

Conclusion: This study is based on the importance of AAB in converting low-value protein waste to value-added products such as acetic acid, ascorbic acid, cellulose and others. This is due to their ability to use protein waste as a substrate.

Keywords: *Acetobacter aceti*, *Acid Bacteria*, *Gluconobacter oxydans*, *Protein Waste*

ABBREVIATION

AAB: Acetic acid bacteria

GYC: Glucose yeast calcium carbonate

YGEA: Yeast glucose ethanol acetic acid

1. INTRODUCTION

Louis Pasteur first described acetic acid bacteria (AAB) as vinegar bacteria. They constitute a diverse group of bacteria known for their importance in producing fermented foods and beverages, especially in the production of acetic acid (ethanoic acid) in vinegar making [1]. AAB are obligatory aerobes usually found in sugary, acidic and alcoholic niches belonging to the Alphaproteobacteria class, Rhodospirillales order, and Acetobacteraceae family [2]. They are known for oxidizing carbohydrates, alcohols and sugar

Comment [ENA1]: This should be incorporated into the Background as a justification for the study.

Comment [ENA2]: What presumptive method was used? Presumptive identification can only be at genus level.

alcohols (polyols) into their corresponding organic acids, aldehydes or ketones [3]; this process is known as oxidative fermentation. Acetic acid bacteria are motile, ellipsoidal to rod-shaped gram-negative bacteria occurring singly, pairs or chains. They move with the aid of either polar or peritrichous flagella and do not form endospores as defensive resistance. They are obligate aerobes with oxygen as the terminal electron acceptor in the electron transport chain. They grow at an optimum pH of 5-6.5 but can also grow at lower pH values between 3-4. They are catalase positive and oxidase negative with sizes ranging from 0.4-1 µm wide and 0.8-4.5 µm long. They are mesophiles having an optimum growth temperature between 25 and 30°C. They produce different polysaccharides and pigments in solid cultures [4,5].

An important characteristic that makes AAB desired industrially is its ability to oxidize ethanol to acetic acid. Two membrane-bound dehydrogenases catalyze this process; pyrroloquinoline quinone (PQQ)-dependent alcohol dehydrogenase (ADH) and membrane-bound aldehyde dehydrogenase (ALDH), located near ADH [6,7]. On acetic acid production, some genera of AAB further oxidize acetic acid to CO₂ and H₂O. This property is known as over-oxidation of ethanol and is customarily used to distinguish *Gluconobacter*, which does not have the capability. Over-oxidation depends on the medium composition, mainly when bacteria use ethanol [8].

Formatted: Font: Italic

Aside from the use of AAB in the production of vinegar making and fermented foods, they are harnessed industrially to produce relevant compounds, primarily ascorbic acid (vitamin C) and miglitol (antidiabetic drug) [9]. They also have an essential role in cocoa fermentation [10]. Also, species of the genera *Acetobacter* and *Gluconacetobacter* are used in agriculture, where nitrogen fixation is attributed to strains of *G. diazotrophicus*. Furthermore, some genera of AAB are known for their production of exopolysaccharides (EPS) and bacterial cellulose. The absence of hemicellulose and lignin in bacterial cellulose offers an advantage over conventional cellulose [11]. Gluconic acid, sorbose and ascorbic acid are also products of AAB metabolism. Due to the importance of AAB metabolic products, many kinds of research have been carried out. However, most researchers focus on their isolations from sugary substrates and alcohol. In contrast, this study emphasizes identifying acetic acid bacteria isolated from waste protein waste as waste management. In addition to waste management, this study aims to isolate

Formatted: Font: Italic

Comment [ENA3]: This sentence is confusing.

AAB from protein wastes like milk, fish, egg and meat, estimate the amount of acetic acid produced, and show the transformation of waste (protein waste) to wealth (acetic acid).

2. MATERIALS AND METHODS

2.1. Study area

Umuahia is the capital city of Abia State in South-Eastern Nigeria, coordinating latitude 5° 32'N and longitude 7° 29'E on the Northern and Eastern hemispheres respectively, located along the railroad that lies between Port-Harcourt to the South and Enugu city to its North.

2.2. Media Used

Glucose yeast extract calcium carbonate medium (GYC), peptone water, yeast glucose ethanol acetic acid medium, triple sugar iron agar (TSI), Simmon citrate agar, nutrient agar. All media were prepared according to manufacturers' instructions.

Comment [ENA4]: Is it necessary to state the media used? The author can just indicate in the methods which media was used at which point. Also indicating the manufacturer and country of origin.

2.3. Sample Description and Sampling Procedure

The samples used are waste protein products (fish, milk, meat, egg, soy milk, beans and unpasteurized milk). Two different samples of each protein waste listed ~~in the bracket~~ were collected randomly from ~~different areas in Umudike, Umuahia~~. Each sample was collected aseptically and handled carefully to ensure no change between sampling and analysis. The seven samples were taken to the laboratory, where the culture was done immediately.

Comment [ENA5]: The author should be a little more specific as to the collection points. Were samples taken from markets, restaurants, or waste sites?

2.4. Preparation of Samples and Isolation of Acetic Acid Bacteria

~~For~~ the isolation of acetic acid bacteria (AAB), each sample (1 ml/ 1g) was suspended in 9 ml sterilized distilled water and ~~homogenized mixed thoroughly~~. Serial dilutions (10^{-3} & 10^{-7}) were prepared, and 0.1ml of each diluted sample was spread on glucose yeast extract CaCO_3 (GYC) medium plates. All plates were incubated aerobically for 2- 5 days at 30°C, and colonies with clear zone were isolated.

Comment [ENA6]: This is not clear. Were these the only dilutions prepared? Or they were the only dilutions plated?

2.5. Purification and Maintenance of the Microbial Isolates

~~The streak plate method obtained P~~ure cultures of the AAB ~~were obtained isolates~~ by sub-culturing in fresh GYC medium. The final cultures containing discrete colonies were maintained by transferring onto

McCartney bottles containing agar slants, and they were stored in the refrigerator at 4°C for further analysis.

2.6. Identification of AAB Using Biochemical Tests

Acetic acid bacteria species, after isolation, were identified based on microbiological, cultural and biochemical properties as described by Cheesbrough [12]. Biochemical tests include Gram's Staining, catalase test, oxidase test (cytochrome oxidase test), sugar fermentation test, indole test, citrate utilization test, and hydrogen sulfide test.

2.7. Endospore Staining

A smear of the organism was made on the slide, then saturated with malachite green stain solution and steamed for 5 minutes keeping blotting paper moist and adding more dye as required. Then the slide was washed under the tap and then counterstained with 0.5 safranin for 30 seconds, washed again and blotted dry and then observed under the microscope. Endospores stain bright green while vegetative cells stain pink.

Comment [ENA7]: What is the purpose of this test? The AAB are not endospore formers.

2.8. Acetic Acid Production

~~Selected colonies of AAB were diluted with peptone water until an OD₆₀₀ of 0.5 was achieved. The selected colonies from GYC were transferred to peptone water until~~ OD₆₀₀ of 0.5 was achieved. Inoculum size of 4% from the preceding peptone water was cultured in yeast glucose ethanol acetic acid medium for acetic acid production. The production medium was aerated, and samples were taken at 48 hours intervals.

Comment [ENA8]: How was this determined? This sentence is not clear.

Comment [ENA9]: Over how many hours in total? How was the medium aerated?

2.8.1. Estimation of acetic acid: 5ml of the culture was mixed with 20ml of distilled water. 3-5 drops of phenolphthalein were added. The solution was titrated against 0.05N NaOH. The amount (g) of acetic acid produced in 100ml of the medium was calculated using the formula:

Comment [ENA10]: Is there a reference for this methodology used?

Acetic acid (g/100ml) = volume of NaOH (ml) used in titration * 0.03 * 20

3. RESULTS

3.1 Macroscopic Characteristics of Isolates

Colony morphologies of AAB isolates grown on GYC media are listed in Table 1. ~~The cultural characteristics observed on the GYC medium is shown in Table 1.~~ Different colonies were observed with different shapes as below.

Table 1: Colony Morphology of AAB on GYC media. ~~Cultural appearance of colonies.~~

Sample	Appearance on agar
Raw milk 1	Large creamy/ milk coloured colonies were observed.
Raw milk 2	Creamy colonies with a clear zone around them.
Beans 1	Creamy mucoid colonies with clustered growth.
Beans 2	No growth appeared.
Meat 1	Creamy large elevated colonies with a clear zone around the colonies.
Meat 2	Large creamy convex colonies observed.
Milk 1	Creamy large elevated colonies were observed.
Milk 2	Creamy elevated colonies were observed.
Fish 1	Creamy mucoid colonies with clustered growth.
Fish 2	Tiny creamy colonies and yellow colonies were observed.
Egg 1	Large creamy elevated colonies appeared on plates.
Egg 2	Yellow colonies and creamy colonies observed.
Soy milk 1	Creamy large elevated colonies with a smooth edge were observed.
Soy milk 2	No growth observed.

Comment [ENA11]: Was there just one colony morphology per sample?

Comment [ENA12]: Maybe a more appropriate terminology than "milk coloured" can be used.

3.2 Growth of Colonies Per Time

Table 2 shows the time for visible growth to be noticed on each media plate. There was no growth on all plates till after 48 hours. However, colonies appeared at 72 hours on some medium plates while there was no growth on two plates (Beans 2 and Soy milk 2) even after 144 hours.

Table 2: Growth of colonies per time

Sample No	Growth of Colonies/ Time					
	24 hrs	48 hrs	72 hrs	96 hrs	120 hrs	144 hrs
Raw milk 1	No	No	Yes	Yes	Yes	Yes
Raw milk 2	No	No	Yes	Yes	Yes	Yes
Beans 1	No	No	Yes	Yes	Yes	Yes
Beans 2	No	No	No	No	No	No
Meat 1	No	No	Yes	Yes	Yes	Yes
Meat 2	No	No	Yes	Yes	Yes	Yes
Milk 1	No	No	Yes	Yes	Yes	Yes
Milk 2	No	No	Yes	Yes	Yes	Yes
Fish 1	No	No	Yes	Yes	Yes	Yes
Fish 2	No	No	Yes	Yes	Yes	Yes
Egg 1	No	No	Yes	Yes	Yes	Yes
Egg 2	No	No	No	Yes	Yes	Yes
Soy milk 1	No	No	No	Yes	Yes	Yes
Soy milk 2	No	No	No	No	No	No

3.3 Biochemical Characterization Tests for AAB

Isolates with a clear zone around their colonies were presumed to be AAB and subjected to further biochemical tests for confirmation. Table 3. shows the reaction of selected isolates to biochemical tests, including catalase test, oxidase test, Gram stain, H₂S test, Gas production, Indole, and Citrate test.

Table 3: Characterization tests for acetic acid bacteria (AAB)

Sample	Catalase	Oxidase	Gram stain	Cell morphology	H ₂ S	Gas	Indole	Citrate
Raw milk 1	+	-	+	Cocci	-	-	-	+
Raw milk 2	+	+	-	rod	-	-	-	-
Beans 1	-	-	+	rod	-	-	-	-
Meat 1	+	-	+	Rod	+	-	-	+
Meat 2	+	-	-	rod	-	+	-	+
Milk 1	+	-	-	Rod	-	+	-	+
Milk 2	+	-	+	cocci	+	-	-	+
Fish 1	+	+	-	Rod	-	+	-	+
Fish 2	+	-	-	rod	-	-	-	+
Egg 1	+	+	-	Rod	+	-	-	+
Egg 2	+	-	+	Cocci	-	-	-	-
Soy milk 1	+	-	+	rod	-	-	-	-

+ = positive, - = negative.

3.4 Estimation of Acetic Acid Production.

Isolates whose biochemical characteristics agree with **Bergey's manual of determinative bacteriology** were cultured using Yeast glucose ethanol acetic acid (YGEA), and the amount of acetic acid produced by each isolate was estimated by titration. Tables 4, 5, and 6 shows the amount of acetic acid produced in g/100ml

Comment [ENA13]: Which particular characteristics were selected? These should be indicated in the manuscript.

Table 4: Estimation of acetic acid produced in milk 1

Initial (ml)	Final (ml)	(ml)
0.00	40.30	40.30
0.00	39.70	39.70
0.00	39.80	39.80

$$40.30 + 39.70 + 39.80 = 119.80; 119.80/3 = 39.93$$

$$\text{Acetic acid (g/100ml)} = \text{volume of NaOH (ml) used in titration} * 0.03 * 20$$

$$\text{Acetic acid (g/100ml)} = 39.93 * 0.03 * 20 = 23.96$$

Table 5: Estimation of acetic acid produced in fish 2.

Initial (ml)	Final (ml)	(ml)
0.00	40.00	40.00
0.00	41.30	41.30
0.00	40.40	40.40

$$40.00 + 41.30 + 40.40 = 121.70; 121.70/3 = 40.57$$

$$\text{Acetic acid (g/100ml)} = 40.57 * 0.03 * 20 = 24.34$$

Table 6: Estimation of acetic acid produced in meat 2

Initial (ml)	Final (ml)	(ml)
0.00	39.00	39.00
0.00	39.50	39.50
0.00	38.80	38.80

$$39.00 + 39.50 + 38.80 = 117.30; 117.30/3 = 39.10$$

$$\text{Acetic acid (g/100ml)} = 39.10 \times 0.03 \times 20 = 23.46$$

4. DISCUSSION

Just as AAB is commonly isolated from a diversity of AAB in different fruits, vegetables, soil, water, and carbohydrate sources [13, 14], this study shows that protein-rich foods harbour AAB and can be used as a substrate to produce other value-added products. In the present study, acetic acid bacteria that comprise a large group of obligate aerobic Gram-negative bacteria commonly found in association with various sugary materials were isolated and characterized from the various protein wastes in Umudike, Umuahia. For this reason, various protein waste samples were targeted. A total of 25 (white, creamy and yellow pale coloured) colonies on GYC agar plates were derived from protein waste samples collected from various markets of Umudike (Table 1). Out of 25 isolated colonies, 14 suspected acetic acid bacteria were screened based on their morphological, cultural and biochemical properties. The fourteen strains were primarily screened and detected as acetic acid bacteria because they were small, white, raised and produced clear zones around their colonies in the selective GYC culture medium. The property of clear zones around their colonies agrees with the findings of Trcek [15] and Assietta *et al.* [16]. However, because clear zone can be produced by other strains such as lactic acid bacteria [14], further biochemical tests were carried out to confirm the presence of AAB. The theory of using CaCO_3 in the media was that if the grown bacteria on the media are acetic acid bacteria, the AAB will produce acetic acid to react with CaCO_3 to form Calcium acetate ($\text{Ca}(\text{CH}_3\text{COO})_2$), water and carbon dioxide. The calcium acetate is more soluble in water than CaCO_3 with a solubility of 34.7 g/100 mL (20 °C), which made the clear zones around AAB colonies [17].

Comment [ENA14]: Can the author clarify this statement? AAB stands for Acetic acid bacteria, I presume?

Examination using Gram staining revealed that 6 isolates were Gram-negative rods. Following Gram stain, biochemical tests such as oxidase, catalase, Indole, and hydrogen sulfide were performed using the selected 6 Gram-negative bacterial isolates according to the standard guideline of Bergey's manual of determinative bacteriology [18]. Several biochemical characterizations, including negative oxidase and positive catalase tests (Table 3), suggested that our isolates from 3 samples (milk 1, fish 2 and meat 2) are acetic acid bacteria. These characteristics agree with those of Gullo M. and Giudici, P. [19], Beheshti and Shafiei [20], Farzana *et al.* [21] and Zahoo *et al.* [22].

Morphologically identified and biochemically confirmed acetic acid bacterial isolates were then further checked for their abilities to produce acetic acid by titration in YGEA (Yeast, Glucose, Ethanol, Acetic acid) medium. Acetic acid production rate varied among the isolates at different incubation periods. Three acetic acid-producing strains isolated from fish 2 showed the highest acetic acid production (24.34 g/100ml) rate at 37°C (Table 5). In contrast, those from milk 1 and meat 2 showed 23.96 g/100ml and 23.46 g/100ml, respectively this is similar to the findings of Lin *et al.* [23] and Beheshti and Shafiei [20] but contrary to research done by Farzana *et al.* [21], this could be because they used a different substrate (decomposed Fruits) in their study.

5. CONCLUSION

Most previous studies show that acetic acid bacteria are commonly associated with various kinds of sugary materials and alcohol. However, AABs were isolated and characterized from waste protein foods such as fish, meat, and milk in this study. This shows that these protein waste sources can serve as a substrate for AAB growth. In these findings, these wastes can be converted to wealth by applying AAB to convert many low-cost protein wastes to high-value products, including acetic acid, bacterial cellulose, vitamin C and other exopolysaccharides, industrially important. This is also important in waste management and environmental conservation.

REFERENCES

1. Hutkins RW. Vinegar fermentation. *Microbiology and technology fermented foods*. 2006; 397-417). Ames, IA: Blackwell publishing.

2. Kersters K, Lisdiyanti P, Komagata K, Swings J. The family Acetobacteraceae: The genera *Acetobacter*, *Acidomonas*, *Asaia*, *Gluconacetobacter*, *Gluconobacter*, and *Kozakia*. *The Prokaryotes*. 2006; 5:163–200.
3. Taban BM, Saichana, N. Physiology and biochemistry of acetic acid bacteria. In I. Y. Sengun (Ed.), *Acetic acid bacteria: Fundamentals and food application* 2017; 71-91. Boca Raton, FL: CRC press.
4. Sengun IY, Karabiyikli S. Importance of acetic acid bacteria in food industry. *Food control*. 2011; 22(5): 647-56.
5. Wang B, Shao Y, Chen F. Overview on mechanisms of acetic acid resistance in acetic acid bacteria. *World J. Microbio. Biotech*. 2015; 31 (2): 255-63.
6. Adachi O, Ano Y, Toyama H, Matsushita K. Biooxidation with PQQ- and FAD dependent dehydrogenases. In: Schmid RD, Urlacher VB, editors. *Modern biooxidation: Enzymes, reactions and applications*. Weinheim, Germany: Wiley-VCH Verlag GmbH & Co. KGaA. 2017.
7. Yakushi T, Matsushita K. Alcohol dehydrogenase of acetic acid bacteria: Structure, mode of action, and applications in biotechnology. *Applied Microbio. Biotech*. 2010; 86(5):1257–65.
8. Gullo M, Verzelloni E, Canonico M. Aerobic submerged fermentation by acetic acid bacteria for vinegar production: Process and biotechnological aspects. *Process Biochemi*. 2014;49(10):1571–9.
9. Shinjoh M, Toyama H. Industrial application of acetic acid bacteria (vitamin C and others). In K. Mutsushita, H. Toyama, N. Tonouchi, & A. Okamoto-Kainuma (Eds.), *acetic acid bacteria: Ecology. physiol*. 2016; 321-338. Tokyo, Japan: *Springer Nature*.
10. Pothakos V, Illeghems K, Laureys D, Spiteals F, Vandamme P, De Vuyst L. Acetic acid bacteria in food and beverage ecosystems. In K. Matsushita, H. Toyama, N. Tonouch, & A. Okamoto-Kainuma (Eds.), *Acetic acid bacteria: Ecology. physiol*. 2016; 73-99. Tokyo, Japan: *Springer Nature*.
11. Dagbagli S, Goksungur Y. Exopolysaccharide production of acetic acid bacteria. In I. Y. Sengun (Ed.), *Acetic acid bacteria: Fundamentals and food application*. 2017; 120-141). Boca Raton, FL: CRC Press.

12. Cheesbrough M. *District Laboratory Practice in Tropical Countries*, 2, 2006; 182-186. Cambridge University Press.
13. Sharafi SM, Rasooli I, Beheshti-Maal K. Isolation, characterization and optimization of indigenous acetic acid bacteria and evaluation of their preservation methods. *Iranian Journal of Microbiology*. 2010; 2(3):38-45.
14. Giudici P, Gullo R. A theoretical model to predict the age of traditional balsamic vinegar. *J. Food Engin.* 2008; 82:121-127.
15. Trcek. Quick identification of acetic acid bacteria based on nucleotide sequences of the 16S-23S rDNA internal transcribed spacer region and the PQQ-dependent alcohol dehydrogenase gene. *Systematic Appld Microb.* 2015; 28(8):735-745.
16. Assietta O, Somda MK, Ouattara AT, N'doye B, Traore AS, Aboubakar S. Molecular identification of acetic acid bacteria isolated from fermented mango juices of Burkina Faso: 16S rRNA gene sequencing. *Afri. J. Bioech.* 2019; 18(29); 766-773.
17. Gordon A, Tristan F. *SI Chemical Data Book* (4th ed.). John Wiley & Sons Australia, Ltd, 2007; 60, 156-170.
18. Goodfellow M, Peter K, Busse H, Trujillo ME, Ludwig W, Suzuki KI, Parte A. *Bergey's Manual of Systematic Bacteriology*. 2012; 5, The Actinobacteria, Springer, New York.
19. Gullo M, Giudici P. Isolation and selection of acetic acid bacteria strains for traditional balsamic vinegar. *Industrie Delle Bevande*, 2006; 35, 345-350.
20. Beheshti Maal, K, Shafiei, R. Isolation and Characterization of an Acetobacter Strain from Iranian White-Red Cherry as a Potential Strain for Cherry Vinegar Production in Microbial Biotechnology. *Asian Journal of Biotechnology*. 2010; 2, 53-59.
21. Farzana D, Fahmida A, Ali AT. Screening of Acetic Acid Producing Microorganisms from Decomposed Fruits for Vinegar Production. *Scientific research publishing*. 2015; 5, 291-297
22. Zahoor T, Siddique F, Farooq U. Isolation and characterization of vinegar culture (*Acetobacter aceti*) from indigenous sources. *British Food J.* 2006; 108, 429-439.

23. Lin SP, Huang YH, Hsu KD, Lai YJ, Chen YK, Cheng KC. Isolation and identification of cellulose-producing strain *Komagataeibacter intermedius* from fermented fruit juice. *Carbohydrate Polymers*. 2016; 151:827–33.

UNDER PEER REVIEW