

Original Research Article

THYROID HORMONES AND RED CELL INDICES OF HIV SUBJECTS AT ENUGU

ABSTRACT

Background: Thyroid hormones act on nearly every cell in the body including red blood cells. Thyroid hormones are also important to proper development and differentiation of all cells of the human body. Human Immunodeficiency Virus (HIV) is a virus that destroys certain cells in the immune system. The aim of this study was to determine free triiodothyronine (FT3), free thyroxine (T4) and red cell indices level of HIV subjects attending clinic at Enugu State University of Science and Technology Teaching Hospital.

Place and Duration of Study: Human Immunodeficiency Virus (HIV) clinic and Haematology Laboratory both of Enugu State University of Science and Technology Teaching Hospital, between August and October 2021.

Methodology: Sixty HIV subjects and thirty apparently healthy individuals within the age range of 30 - 60 years were recruited for this study. Free triiodothyronine (FT3), free thyroxine (FT4), hemoglobin (HGB), packed cell volume (PCV), red blood cell count (RBC), mean cell volume (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), red cell distribution width standard deviation (RDW-SD), red cell distribution width coefficient of variation (RDW-CV) were determined in HIV subjects and apparently healthy individuals.

Results: There was significant difference in MCV (125.79 ± 7.57 vs 78.89 ± 5.71 fl), MCH (23.17 ± 2.10 vs 27.21 ± 3.68 pg), MCHC (25.08 ± 2.74 vs 30.09 ± 3.27 g/dl), RDW-SD (61.52 ± 7.55 vs 46.23 ± 5.87 fl), RDW-CV (23.06 ± 5.78 vs 43.87 ± 1.28 fl) and PCV (39.38 ± 3.06 vs 31.75 ± 2.44 %) between patients on antiretroviral therapy (ART) and ART-naïve patients respectively. There was significant difference in T4 (8.29 ± 1.29 vs 6.12 µg/dl), MCV (78.89 ± 5.71 vs 84.47 ± 3.79 fl), MCH (27.21 ± 3.68 vs 29.20 ± 2.27 pg), RDW-CV (43.87 ± 1.28 vs 12.51 ± 1.08 fl) and PCV (31.75 ± 2.44 vs 37.63 ± 2.95 %) on ART-naïve patients and apparently healthy subjects respectively. There was a positive correlation between FT3 and MCH in HIV patients on ART ($p = 0.016$).

Conclusion: In people taking antiretroviral therapy, mean cell haemoglobin, mean cell hemoglobin concentration and red cell distribution width coefficient of variation significantly reduced, as well as raised mean cell volume and red cell distribution width standard deviation. This study also observed reduced mean cell haemoglobin, mean cell haemoglobin concentration and mean cell volume in ART-naïve subjects.

Keywords: Triiodothyronine, thyroxine, antiretroviral therapy, red cell distribution width

1. INTRODUCTION

Thyroid hormones have great effects on numerous physiologic processes, including development, growth and metabolism. Countless effects of thyroid hormone have been described by study of excess and

deficiency states. Thyroid hormones cause many metabolic activities in tissues, which lead to an increase in basal metabolic rate. One effect of this activity is to raise body heat production, which appears to occur, slightly in part, from raised oxygen consumption and rates of adenosine triphosphate hydrolysis [1]. Human immunodeficiency virus (HIV) infection is one of the most critical diseases globally, with about 940,000 deaths in 2017 [2]; which lead to decline in CD4 T lymphocytes cells, leading to immune system failure, susceptibility to opportunistic infections and cancer development. HIV diagnosis has improved over the years because of antiretroviral regimens [3]. Several disease conditions caused by HIV have been recently reported; such as anomalies of pancreas, endocrine and gonadal metabolisms [4, 5]. Studies have also reported antiretroviral medications induce complications such as hypercholesterolaemia, lipodystrophy and glucose intolerance [6]. In HIV patients, manifestations of thyroid problems which include functional disturbances and abnormal changes have been reported. Mechanism behind thyroid disease in HIV patients has been unclear. But some explanation such as opportunistic diseases induce reversible thyroid organ damage has been made [7]. Thyroid hormones are important for different metabolic activities in the body. Thyroid dysfunction can decrease the quality of life in HIV patients, hence there is need to carry out thyroid function screening test in HIV patients during treatment. Red blood cell (RBC) indices, such as the mean cell hemoglobin concentration (MCHC), mean cell volume (MCV) and mean cell hemoglobin (MCH), haemoglobin level and red blood cell count are indicators of systemic iron status [8, 9]. Exact measures of red blood cell size such as mean cell volume, which also depend on reduction-oxidation homeostasis and normal biosynthesis of red blood cell membrane lipids, may also show mitochondrial dysfunction [10,11]. In HIV infection, anemia has been reported in individuals with stunted viremia during highly active antiretroviral therapy, with or without macrocytosis [12, 13]. Therefore, the aim of this study was to determine free triiodothyronine (FT3), free thyroxine (T4) and red cell indices level of HIV subjects taken ART treatment, not yet on ART and to correlate FT3, FT4 and red cell indices.

Comment [DEU1]: leads

Comment [DEU2]: leads

Comment [DEU3]: resulting in

Comment [DEU4]: taking

2. METHODOLOGY

The case control study was carried at Enugu State University of Science and Technology Teaching Hospital from August to October 2021. Approval from Ethical Committee was obtained prior to the study. The subjects were selected by simple random sampling method. After obtaining informed consent, the subjects were recruited for both groups. Patients who were known HIV for at least 6 months were included in this study. These patients were of either gender and between the ages of 30-60 years and ART duration of 1-15 years. The subjects were grouped into three; Group one were HIV positive patients taken ART, group two were HIV positive patients not yet on ART and group three were apparently healthy individuals. Highly active antiretroviral therapy treatment duration was grouped as less than 5 years, 6-10 years and more than 10 years. The drugs are Tenofovir, Lamivudine, Dolutegravir. Blood samples were collected via venepuncture. Free triiodothyronine and free thyroxine were determined by competitive solid-phase enzyme linked immunosorbent assay. Red cell indices were analyzed by automated haematology analyzer (Mindray/BC-5150). The data were evaluated in statistical programme SPSS version 21.0. Student's t-test and one-way ANOVA were applied with mean \pm standard deviation. The association between FT3, FT4 and red cell indices was determined by Pearson correlation. P- Value \leq 0.05 was considered as statistical significant.

Comment [DEU5]: confirmed HIV positive

Comment [DEU6]: taking

3. RESULT

Table 1 showed mean \pm SD of free triiodothyronine (FT3), free thyroxine (FT4), hemoglobin (HGB), packed cell volume (PCV), red blood cell count (RBC), mean cell volume (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), red cell distribution width standard deviation (RDW-SD), and red cell distribution width coefficient of variation (RDW-CV) of HIV subjects on ART, HIV ART-naïve subjects, control subjects using one way ANOVA. HIV subjects on ART have significantly decreased MCH (23.17 \pm 2.10 vs 27.21 \pm 3.68 pg), MCHC (25.08 \pm 2.74 vs 30.09 \pm 3.27 g/dl) and RDW-CV (23.06 \pm 5.78 vs 43.87 \pm 1.28 %) compared with ART-naïve subjects ($p < 0.001$, < 0.001 and < 0.001 respectively). Again, HIV subjects on ART have significantly decreased MCH (23.17 \pm 2.10 vs 29.20 \pm 2.27 pg), MCHC (25.08 \pm 2.74 vs 33.70 \pm 2.39 g/dl) and significantly increased MCV (125.79 \pm 7.57 vs 84.47 \pm 3.79 fl), RDW-SD (61.52 \pm 7.55 vs 44.23 \pm 2.16 fl) and RDW-CV (23.06 \pm 5.78 vs 12.51 \pm 1.08 %) compared with control subjects ($p < 0.001$, < 0.001 , < 0.001 and < 0.001 respectively). Antiretroviral therapy-naïve subjects have significantly decreased MCV (78.89 \pm 5.71

vs84.47± 3.79 fl), MCH (27.21 ± 3.68 vs29.20 ± 2.27 pg), MCHC (30.09 ± 3.27 vs33.70 ± 2.39 g/dl), PCV (31.75 ± 2.44 vs37.63 ± 2.95 %) and significantly increased RDW-CV (43.87 ± 1.28 vs12.51 ± 1.08 %), FT4 (8.29 ± 1.93 vs6.12 ± 1.51 µg/dl) compared with control subjects (p=<0.001, 0.04, <0.001, <0.001, <0.001 and 0.03respectively).Table 2 showed mean ± SD of Free triiodothyronine(FT3), free thyroxine (FT4), hemoglobin (HGB), packed cell volume (PCV), red blood cell count (RBC), mean cell volume (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), red cell distribution width standard deviation (RDW-SD), and red cell distribution width coefficient of variation (RDW-CV) of drug duration of < 5years, 6-10year and > 10 year for HIV subjects on ART using one way ANOVA.HIV patients that have taken ART treatment for more than 10 years has significantly increased MCV (132.40 ± 2.30 vs122.05 ± 7.67 fl) compared with those that are still less than 5 years (p=0.001). Packed cell volume of HIV patients that have taken ART for more than 10 years were significantly lower (36.92 ± 1.28 vs41.45 ± 1.44 %) (p=<0.001) compared with those that are between 6-10 years. Table 3 showed correlation between red cell indices and thyroid hormones in HIV subjects on ART using Pearson correlation. Mean cell hemoglobin significantly correlated with free triiodothyronine (p=0.016) in HIV subjects on ART.Table 4 showed correlation between red cell indices and thyroid hormones of HIV subjects not yet on ART treatment using Pearson correlation. However, no significant correlation was found.

Comment [DEU7]: Should take a new paragraph

Comment [DEU8]: had

Comment [DEU9]: who were

Comment [DEU10]: who had

Comment [DEU11]: who were

Comment [DEU12]: should take a new paragraph

Comment [DEU13]: should take a new paragraph

Table 1: Mean values of FT3, FT4, HGB, PCV, RBC, MCV, MCH, MCHC, RDW-SD, RDW-CV of HIV subjects on ART, ART-naïve subjects and control subjects

	FT3(ng/ml)	FT4(µg/dl)	HGB(g/dl)	PCV(%)	RBC(x10 ¹² /l)	MCV(fl)	MCH(pg)	MCHC(g/dl)	RDW-SD(fl)	RDW-CV(%)
HA	1.23±	7.38±	12.82±	39.38±	4.22±	125.79±	23.17±	25.08±	61.52±	23.06±
N=30		0.44	1.89	0.82	3.06	0.54	7.57	2.10	2.74	7.55
AN	1.17±	8.29±	10.63±	31.75±	3.53±	78.89±	27.21±	30.09±	46.23±	43.87±
N=30		0.42	1.93	0.86	2.44	0.26	5.71	3.68	3.27	5.87
C	1.23±	6.12±	12.61±	37.63±	4.27±	84.47±	29.20±	33.70±	44.23±	12.51±
N=30		0.54	1.51	0.97	2.95	0.53	3.79	2.27	2.39	2.16
F (P) value	0.18	3.52	56.27	59.74	24.03	567.07	36.79	70.66	83.91	81.79
		(0.84)	(0.03)	(<0.001)	(<0.001)	(<0.001)	(<0.001)	(<0.001)	(<0.001)	(<0.001)
HA vs AN	0.85	0.17	<0.001**	<0.001**	<0.001**	<0.001**	<0.001**	<0.001**	<0.001**	<0.001**
HA vs C	1.00	0.83	0.65	0.07	0.91	<0.001**	<0.001**	<0.001**	<0.001**	<0.001**
AN vs C	0.87	0.03*	<0.001**	<0.001**	<0.001**	0.04*	<0.001**	0.20	<0.001**	<0.001**

Abbreviations

HA= HIV subjects on ART

AN= ART naïve subjects

C= control

Table 2: Mean values of FT3, FT4, HGB, PCV, MCV, MCH, MCHC, RDW-SD, RDW-CV of drug duration of < 5years, 6-10year and > 10 year for HIV subjects on ART

	FT3(ng/ml)	FT4(µg/dl)	HGB(g/dl)	PCV(%)	RBC(x10 ¹² /l)	MCV(fl)	MCH(pg)	MCHC(g/dl)	RDW-SD(fl)	RDW-CV(%)
<5yrs	1.17±	7.29±	12.98±	39.67±	4.18±	122.05±	23.22±	25.06±	59.99±	21.56±
N=14		0.56	2.50	0.92	3.21	0.59	7.67	2.25	2.89	10.14
6-10yrs	1.23±	7.29±	12.95±	41.45±	4.29±	127.55±	22.78±	24.99±	61.64±	23.58±
N=11	0.28	1.24	0.56	1.44	0.58	6.53	1.24	2.70	4.30	4.45

>10yrs	1.38±	7.86±	12.10±	36.92±	4.15±	132.40±	23.88±	2.32±	65.60±	24.80±
N=5	0.38	1.20	0.75	1.28	0.41	2.30	3.27	2.97	1.82	4.02
F (P) value	0.40	0.18	2.54	5.99	0.16	5.08	0.46	0.02	1.02	1.66
	(0.67)	(0.84)	(0.09)	(0.01)	(0.85)	(0.01)	(0.64)	(0.98)	(0.37)	(0.21)
<5yrs vs 6-10yrs	0.94	0.17	0.99	0.18	0.88	0.15	0.81	0.10	0.85	0.43
<5yrs vs>10yrs	0.64	0.83	0.15	0.04*	0.99	0.001*	0.91	0.98	0.15	0.31
6-10yrs vs>10yrs	0.72	0.88	0.14	<0.001**	0.85	0.11	0.76	0.98	0.06	0.85

Table 3: Pearson correlation coefficient between red cell indices and thyroid hormones in HIV subjects on ART

HGB(g/dl)	PCV(%)	RBC(x10 ¹² /l)	MCV(fl)	MCH(pg)	MCHC(g/dl)	RDW-SD(fl)	RDW-CV(%)
FT3							
r	0.157	0.190	0.063	0.101	0.43	0.123	0.152
p value	0.406	0.315	0.739	0.596	0.016*	0.516	0.424
FT4							
r	0.134	0.150	-0.019	0.107	0.350	0.053	0.201
P value	0.481	0.429	0.920	0.573	0.058	0.783	0.286

Table 4: Pearson correlation coefficient between red cell indices and thyroid hormones in ART- naïve subjects

HGB (g/dl)	PCV (%)	RBC (x10 ¹² /l)	MCV (fl)	MCH (pg)	MCHC (g/dl)	RDW-SD (fl)	RDW-CV (%)
FT3							
r	-0.097	-0.078	-0.090	-0.238	-0.133	-0.237	0.296
p value	0.609	0.681	0.638	0.204	0.484	0.207	0.112
FT4							
r	-0.188	-0.181	-0.298	-0.246	-0.233	0.329	0.221
P value	0.319	0.337	0.110	0.110	0.190	0.215	0.076

Discussion

This study investigates the thyroid hormones and red blood cell indices of 60 HIV- positive adult on ART and ART-naïve. Thus, the commonest findings found in the current study were increased RDW-SD, RDW-CV and MCV in ART subjects. Also in ART- naïve subjects decreased MCV, MCH, MCHC and increased RDW-CV and free thyroxine were found. Red blood cell distribution width (RDW), as well as different type hereof (red cell distribution width standard deviation and red cell distribution width coefficient of variation), is a measure of the range of variation of RBC volume that is reported as part of a typesstandard complete blood count [14]. The average volume of red blood cells is 80-100 femtoliters, but in healthy human blood, individual cell volumes vary. Certain disease conditions, however, give rise to a significantly increased variation in cell size. Higher Red blood cell distribution width value shows greater variation in size [15]. Normal value of RDW-CV in RBC is 11.5-15.4% [16]. In anaemia patients, Red cell distribution width test results are often used together with MCV results to determine the possible causes of the anaemia [17]. Folate and vitamin B₁₂ deficiency anemia normally presents with high RDW and high MCV. Study had shown that micronutrient concentrations are lower among patients with HIV wasting syndrome [18]. Antiretroviral therapy is generally recommended for patients with HIV-related opportunistic infections or a CD4 count < 200 cells/μL. Antiretroviral therapy restores immunologic function [19], but does not put an end to weight loss and wasting [20, 21]. In this study increased levels of MCV, RDW-SD and RDW-CV were found in HIV patient on ART. This finding is consistent with the study done in North India which reported macrocytic anaemia in patients taking ART [22]. Mean cell haemoglobin (MCH)

Comment [DEU14]: investigated

Comment [DEU15]: delete

Comment [DEU16]: types

Comment [DEU17]: types standard

Comment [DEU18]: delete

value refers to the average quantity of haemoglobin present in a single red blood cell. Mean cell haemoglobin value is connected to two other values, MCV and MCHC. At the same moment, MCH, MCV and MCHC are sometimes called red blood cell indices. Mean cell haemoglobin value calculated below 27.5 pg is considered low MCH. A low MCH value shows the presence of iron deficiency anaemia [23]. In this study decreased levels of MCH, MCHC and MCV were found in ART naïve subjects. This finding is consistent with the study done in Indonesia that found significantly lower MCH, HGB and PCV in subjects with $CD4 \leq 350mm^3$ [24]. Thyroid dysfunction has been reported in HIV patients [25]. In the present study, free thyroxine significantly increased in ART-naive subjects and MCH significantly correlated with free triiodothyronine. This finding is nearly identical with study done in India that reported significant positive correlation between thyroid stimulating hormone and CD4 count [26]. The cause of thyroid dysfunction in HIV subjects are not yet clear, but hypotheses involve coexisting infections, autoimmune disease, damage by opportunistic infections, and drug reactions [27]. Thyroid dysfunctions are associated with disease progression, such as severe immunosuppression and increase viral load [28].

Comment [DEU19]: this finding is similar with the study

Comment [DEU20]: with the study

Conclusion

In people taking antiretroviral therapy, mean cell haemoglobin, mean cell hemoglobin concentration and red cell distribution width coefficient of variation significantly reduced, as well as raised mean cell volume and red cell distribution width standard deviation. This study also observed reduced mean cell haemoglobin, mean cell haemoglobin concentration and mean cell volume in ART- naïve subjects. We suggest further study that will involve larger study population and vitamin B12, folate and iron measurement in HIV patients taking antiretroviral therapy.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Mullur R, Liu YY, Brent GA. Thyroid hormone regulation of metabolism. *Physiological Reviews*. 2014; 94:355–382.
2. GBD HIV Collaborators. Global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2017, and forecasts to 2030, for 195 countries and territories: a systematic analysis for the Global Burden of Diseases, Injuries, and Risk Factors Study 2017. *Lancet HIV*. 2019;6(12):e831–59.
3. Oliva-Moreno J, Trapero-Bertran M. Economic impact of HIV in the highly active antiretroviral therapy era – reflections looking forward. *AIDS Rev*. 2018;20:226–235.
4. Noureldeen AF, Qusti SY, Khoja GM. Thyroid function in newly diagnosed HIV-infected patients. *Toxicol Ind Health* 2014;30:919-25
5. Khandelwal D, Tandon N. Overt and subclinical hypothyroidism: who to treat and how. *Drugs*. 2012;72:17-33.
6. Lake JE, Stanley TL, Apovian CM, Bhasin S, Brown TT, Capeau J, Currier JS, Dube MP, Falutz J, Grinspoon SK, Guaraldi G, Martinez E, McComsey GA, Sattler FR, Erlandson KM. Practical Review of Recognition and Management of Obesity and Lipohypertrophy in Human Immunodeficiency Virus Infection. *Clin Infect Dis*. 2017; 64(10):1422-1429.
7. Calza L, Manfredi R, Chiodo F. Subclinical hypothyroidism in HIV-infected patients receiving highly active antiretroviral therapy. *J Acquir Immune Defic Syndr* 2002;31:361-363.

8. Addo OY, Yu EX, Williams AM, Young MF, Sharma AJ, Mei Z, Kassebaum NJ, Jeffers MED, Suchdev PS. Evaluation of Hemoglobin Cutoff Levels to Define Anemia Among Healthy Individuals. *JAMA Netw. Open* 2021;4, e2119123.
9. Gonzales GF, Tapia V, Vásquez-Velásquez C. Changes in hemoglobin levels with age and altitude in preschool-aged children in Peru: The assessment of two individual-based national databases. *Ann. N. Y. Acad. Sci.* 2021;1488: 67–82.
10. Choque-Quispe BM, Alarcón-Yaquette DE, Paredes-Ugarte W, Zaira A, Ochoa A, Gonzalez GF. Is the prevalence of anemia in children living at high altitudes real? An observational study in Peru. *Ann. N. Y. Acad. Sci.* 2020;1473: 35–47.
11. Manckoundia P, Barben J, Asgassou S, Putot A, Konaté A. Iron deficiency in adults: To understand what biological evaluation should be carried out. *Rev. Med. Liege* 2020;75: 791–796.
12. Manckoundia P, Konaté A, Hacquin A, Nuss V, Mihai AM, Vovelle J, Dipanda M, Putot S, Barben J, Putot A. Iron in the General Population and Specificities in Older Adults: Metabolism, Causes and Consequences of Decrease or Overload, and Biological Assessment. *Clin. Interv. Aging* 2020;15: 1927–1938.
13. Sharma AJ, Addo OY, Mei Z, Suchdev PS. Reexamination of hemoglobin adjustments to define anemia: Altitude and smoking. *Ann. N. Y. Acad. Sci.* 2019;1450: 190–203.
14. Lippi G, Plebani M. Red blood cell distribution width (RDW) and human pathology. One size fits all. *Clin Chem Lab Med* 2014;52:1247-9.
15. Lippi G, Salvagno GL, Guidi GC. Red blood cell distribution width is significantly associated with aging and gender. *Clin Chem Lab Med* 2014;52:e197-9
16. Hoffmann JJ, Nabbe KC, van den Broek NM. Effect of age and gender on reference intervals of red blood cell distribution width (RDW) and mean red cell volume (MCV). *Clin Chem Lab Med* 2015;53:2015-9.
17. Gang L, Lifang W. Association of the elevated red blood cell distribution width with the risk of developing diabetes mellitus. *Intern Med* 2016;55:1959-1965.
18. Semba RD, Shah N, Strathdee SA, Vlahov D. High prevalence of iron deficiency and anemia among female injection drug users with and without HIV infection. *J Acquir Immune Defic Syndr.* 2002; 29:142-144
19. Owiredu WK, Quayle L, Amidu N, Addai-Mensah O. Prevalence of anaemia and immunological markers among Ghanaian HAART-naïve HIV-patients and those on HAART. *Afr Health Sci* 2011;11:2-15.
20. Agarwal D, Chakravarty J, Chaube L, Rai M, Agrawal NR, Sundar S. High incidence of zidovudine induced anaemia in HIV infected patients in eastern India. *Indian J Med Res* 2010;132:386-389.
21. Meidani M, Rezaei F, Maracy MR, Avijgan M, Tayeri K. Prevalence, severity, and related factors of anemia in HIV/AIDS patients. *J Res Med Sci* 2012;17:138-142
22. Panwar A, Sharma S C, Kumar S, Sharma A. A study of anemia in human immunodeficiency virus patients: Estimating the prevalence, analyzing the causative effect of nutritional deficiencies, and correlating the degree of severity with CD4 cell counts. *Med J DY Patil Univ* 2016;9:312-318
23. Enawgaw B, Alem M, Addis Z, Melku M. Determination of hematological and immunological parameters among HIV positive patients taking highly active antiretroviral treatment and treatment naïve in the antiretroviral therapy clinic of Gondar University Hospital, Gondar, Northwest Ethiopia: a comparative cross-sectional study. *BMC Hematol.* 2014;14(1):1–7.
24. Lumbanraja SN, Siregar DIS. Association between red blood cell indices and CD4 count in HIV-positive reproductive women. *IOP Conference Series Earth and Environmental Science.* 2018; 125(1):012027. DOI:10.1088/1755-1315/125/1/012027
25. Jain G, Devpura G, Gupta BS. Abnormalities in the thyroid function tests as surrogate marker of advancing HIV infection in infected adults. *J Assoc Physicians India.* 2009;57:508–10.
26. Satyakumar T, Sunibala K, Tomba YS, Ranbir LS, Amuba MS, Salam R. Thyroid dysfunction in human immunodeficiency virus-infected children and its correlation with CD4⁺ T lymphocyte count. *Indian J Endocrinol Metab.* 2015; 19(2): 272–276.

27. Chiarelli F, Galli L, Verrotti A, di Ricco L, Vierucci A, de Martino M. Thyroid function in children with perinatal human immunodeficiency virus type 1 infection. *Thyroid*. 2000;10:499–505
28. Hirschfeld S, Laue L, Cutler GB, Jr, Pizzo PA. Thyroid abnormalities in children infected with human immunodeficiency virus. *J Pediatr*. 1996;128:70–4.

UNDER PEER REVIEW