

MR Imaging of Pancreatic Lesions

ABSTRACT

Aim: The aim of this work was to evaluate the role of MRI in differentiating between benign and malignant pancreatic lesions and its correlation with histopathological results as the reference standard.

Patients and Methods: This MRI study included 30 patients, 17 females and 13 males with a mean age 50 years. Sixteen patients had malignant masses (14 patients were adenocarcinoma, one patient was lymphoma and one patient was metastasis) and 14 patients had benign masses (7 patients were pancreatic pseudocysts, two patients were pancreatic abscesses, three patients were simple cysts and two patients were focal pancreatitis). The main clinical symptom was abdominal pain and most of masses were located in the head of the pancreas.

Results: In our study, 25 cases of the 30 patients showed increased intensity at T2-weighted images. Most of malignant cases showed low or equal intensity on T1- and high intensity on T2-weighted images compared to normal pancreatic parenchyma. In our study, DW-MRI was performed on all subjects at b-values of 500 and 1000 s/mm². Benign pancreatic masses as pancreatic pseudocyst, simple cyst and abscess show low signal intensities on DWI, however malignant pancreatic masses as adenocarcinoma, lymphoma and metastasis show high signal intensities on DWI with a cut-off value of 1.5 x10⁻³ s/mm² for the differentiation of benign from malignant pancreatic masses by b-value 1000 s/mm² with the sensitivity, specificity, PPV, NPV& p value were 100%, 83.33%, 100%, 88.88% and <0.001 respectively.

Conclusion: MRI plays an important role in the diagnosis of different pancreatic lesions and can assess the neoplastic pancreatic lesions with accurate detection of extension, nodal involvement and hepatic metastatic lesions. It also has a major role in differentiation between benign and malignant pancreatic lesions by the aids of DWI.

Keywords: MR Imaging, Pancreatic Lesions

1. INTRODUCTION:

The pancreas is a non-encapsulated retroperitoneal organ composed of a head, uncinate process, neck, body, and tail [1].

Pancreatic abnormalities may manifest as pancreatic enlargement (tumor, acute pancreatitis), pancreatic atrophy (cystic fibrosis, chronic pancreatitis), cystic lesions (pseudocysts, congenital simple cysts, autosomal dominant polycystic renal disease, von Hippel-Lindau disease, cystic fibrosis, cystic neoplasms), or fatty replacement (cystic fibrosis, Shwachman-Diamond syndrome, obesity) [2].

Imaging modalities for pancreas include plain radiography, contrast studies (barium meal studies, PTC / Endoscopic retrograde cholangiopancreatography (ERCP), Angiography), ultrasonography, CT Scanning, Magnetic resonance imaging (MRI) including Magnetic resonance [angiography angiogram](#) (MRA) and Magnetic resonance cholangiopancreatography (MRCP) [3].

Computed tomography plays a primary role in the diagnosis and imaging workup of patients with pancreatic diseases. However, advanced magnetic resonance imaging (MRI) techniques offer images of the pancreas with excellent contrast resolution in a reasonable examination time [4].

MRI plays a triple role in the evaluation of the pancreas: Diagnosis, staging and detection of complications. The role of MRI has increased, especially in the imaging patients with suspected pancreatic neoplasms [5].

Technical innovation in MRI, such as use of phased-array coils, allows improved spatial resolution and faster T1- and T2-weighted sequences for imaging the entire upper abdomen in a single breathhold and providing cross-sectional images of pancreatic parenchyma [6].

Magnetic resonance cholangiopancreatography (MRCP) is used for noninvasive work-up of patients with pancreaticobiliary disease. MRCP is comparable with invasive endoscopic retrograde cholangiopancreatography (ERCP) for diagnosis of extrahepatic bile duct abnormalities [7].

Diffusion-weighted MRI (DWI) is a technique in which phase defocusing and refocusing gradients are used to evaluate the rate of microscopic water diffusion within tissue. Quantitative measurements of the diffusivity of water are described by the apparent diffusion coefficient (ADC) [8].

DWI is becoming an important noninvasive technique in the characterization of biologic tissues based on their water diffusion properties, especially those with high b value. A few recent reports have suggested that DWI with single-shot echo-planar imaging may be helpful in the detection of pancreatic adenocarcinoma [9].

2. PATIENTS & METHODS

This prospective study included 30 patients with pancreatic lesions and their ages ranged from 30 years to 70 years with a mean age 51 years. They were referred from Surgery, Internal Medicine and Oncology Departments to Radiodiagnosis and Medical Imaging Department in our institute. Conventional MRI examinations were performed for all patients. The results of the 30 patients were histopathologically correlated in 21 cases and clinical follow up was carried out for the remaining 9 patients.

Inclusion criteria:

- Patients who were suspected or known to have pancreatic disease regardless to gender.
- Patients who agreed to join the study according to the ethical considerations and a written informed consent was obtained from all the included patients.

Exclusion criteria:

- All patients with relative or absolute MRI contraindications:
 - Patients with a heart pacemaker or cardiac defibrillator.
 - Patients who have an aneurysm clip in their brain.
 - Patients who have cochlear implant.
 - Patients who have implanted neural stimulator.
- Patients with poor renal functions.
- Patients unable to tolerate baseline MRI scan or scan not of adequate quality for analysis (e.g. too much movement artifacts).

Patients were subjected to the following:

1. Clinical assessment:

- Detailed meticulous history taking.
- Clinical examination:
- Laboratory investigations:

2. Radiological assessment:

a. Conventional MR imaging:

Instructions and preparation of the patients:

- Ferromagnetic materials were removed.
- The method of examination was fully explained to all patients before imaging to obtain their consent.

b. Diffusion Weighted Magnetic Resonance Imaging (DW-MRI) and apparent diffusion coefficient (ADC) maps: Were performed for all patients.

3. Histopathological correlation:

Histopathological correlation between diffusion weighted magnetic resonance readings and histopathological sections was performed in 21 cases where biopsy was done either by:

- US guided fine needle aspiration: in 3 cases.
- Postsurgical histopathology: in 14 cases.
- Core biopsy: in 4 cases.

Technique of examination:

- ☑ Patients were positioned in supine position and were instructed to hold breath to reduce respiratory artifacts during the examination. Circularly polarized body coil was placed over the upper abdomen.
- ☑ A fast scout scan in sagittal, axial and coronal planes was obtained.
- ☑ All patients with pancreatic masses underwent DW-MRI. This was obtained in axial plane and performed without injection of contrast material. DW-MRI was obtained using a multi-section single shot spin echo EPI sequence (TR/TE/NEX: 3395/100ms/1) with diffusion sensitivities of b-values = 0,500 and 1000 s/mm².
- ☑ The diffusion gradients were applied sequentially in the three orthogonal directions. Sections of 2-4 mm thickness, inter-slice gap of 1mm, a 230-255 mm FOV, and a 256 x 256 matrix were used with average scan time of 35s.
- ☑ ADC maps were formed automatically by the device, a circular regions of interest (ROI) ranging from 10 to 40 mm² according to the size of the mass, were placed in the center of the lesion for each b-value (same slice location).

HOW DID YOU :[1U]Comment
30ARRIVE AT THE SAMPLE SIZE OF
WHAT IS THE STUDY PERIOD OR
DURATION

- ☒ ADC value was obtained with b-values 500 and 1000 s/mm².
- ☒ The ADC values are expressed in square millimeters per second.

Statistical Analysis

The following statistical methods were used for analysis of results of the present study. Data were checked, entered and analyzed using SPSS version 21 for data processing and statistics.

A. Descriptive statistics:

Data were expressed as number and percentage for qualitative variables and mean ± standard deviation for quantitative one.

Data were summarized using:

1. **The arithmetic mean (\bar{X})** as an average describing the central tendency of observations.
2. **The standard deviation (SD)** as a measure of dispersion of the results around the mean.

B. Comparison of means:

The comparison was done using:

- ☒ **The student "t" test** for comparison of means of two independent groups.
- ☒ **Mc Nemar's Chi- square test (X²)**: Used to find the association between row and column variables.

C. Validity of a screening test:

Validity of (DW-MRI) for diagnosis of pancreatic mass was tested by sensitivity, specificity, Positive Predictive Value (PPV) and Negative Predictive Value (NPV).

Receiver operating characteristic (ROC curve) analysis was used to find out the over all predictivity of parameter in and to find out the best cut-off value with detection of sensitivity and specificity at this cut-off value.

1. **Sensitivity**: is the ability of the test to detect true positive cases.
2. **Specificity**: is the ability of the test to detect true negative cases.
3. **Positive predictive value**: is the probability of disease in a patient with positive test result.
4. **Negative predictive value**: is the probability of not having the disease in a subject with negative test result.

Level of significance:

For all above mentioned statistical tests done, the threshold of significance was fixed at 5% level (P-value).

- P value of > 0.05 indicates non-significant results.
- P value of < 0.05 indicates significant results.

The smaller the P value obtained the more significant are the results.

3. RESULTS:

Table 1: Age and sex distribution :

Age	Sex				Percent (%)	
	Male		Female		N	%
	N	%	N	%		
30 - < 40	1	3.3	2	6.7	3	10
40 - < 50	5	16.7	4	13.3	9	30
50 - < 60	6	20	7	23.3	13	43
60 - < 70	1	3.3	4	13.3	5	17
Total	13	43.3	17	56.7	30	100
X²	1.823					
P value	0.611					

Age grouping of the studied group was 10% from 30 to 40 years old, 30% from 40 to 50 years old, 43% from 50 to 60 years old and 17% from 60 to 70 years old.

Sex distribution of the studied group showed that 56.7% were female (17 patients) and 43.3% were male (13 patients).

The most common age group was from 50-60 years (13 patients = 43%). Also pancreatic masses are common in females (57%) than males (43%).

Table 2: Clinical symptoms :

Symptoms	No. of patients	Percent (%)
Abdominal pain	22	73.3%
Jaundice	16	53.3%
Anorexia and weight loss	9	30%
Fever	4	13.3%
Abdominal pain and Jaundice	9	30%
Abdominal pain & Weight loss	7	23.3%

Among different patients' groups: 22 (73.3%) of patients complained of abdominal pain, 16 (53.3%) of them complained of jaundice, 9 (30%) of them complained of anorexia and weight loss, 4 (13.3%) of them complained of fever, 9 (30%) of them complained of abdominal pain and jaundice and 7 (23.3%) of them complained of abdominal pain and weight loss.

Abdominal pain was the commonest presenting symptom in all studied patients as it was detected in more than half of the studied patients (12/30; 73.3%).

Table 3: Laboratory findings (tumor markers) :

Tumor marker	Malignant (n=16) (54%)	Benign (n=14) (46%)	T test	P value
CA 19 – 9 (N. 37 U/mL)	587 ±10	30.5 ±6.6	75.631	0.001*
CEA (N. 2.5 ng/mL)	4.8 ±3	1.1 ±2.3	3.753	0.001*

Table 4: Site of pancreatic lesions :

Site	Benign (n=14) (46%)		Malignant (n=16) (54%)	
	No of patient	Percent	No of patient	Percent
Head	7	23%	10	33%
Body	4	13.3%	3	10%
Tail	2	6.7%	1	3.3%
Body & Tail	1	3.3%	2	6.7%

X2	1.209
P value	0.750

CA 19-9 (carbohydrate antigen)
CEA (carcinoembryonic antigen)

The cutoff level of CA19-9 and CEA markers were set at 37U/mL and 2.5 ng/mL, respectively. In patients with confirmed pancreatic carcinoma, the marker levels were elevated and exceeded the cutoff values and it was higher in patient with irresectable tumors rather than those with resectable tumors whereas their levels were low in the benign lesions.

The most of pancreatic lesions were located in the head of the pancreas (7 benign lesions = 23% and 10 malignant lesions = 33%).

Table 5: Histopathological & clinical follow up results of the 30 patients in our study

Lesion	N	%	Lesion	N	%
Group I: Benign	14	46.7	Group II: Malignant	16	53.3
1) Pancreatic pseudocyst	7	23.3	1) Adenocarcinoma	14	46.7
2) Pancreatic abscess	2	6.7	2) Lymphoma	1	3.3
3) Simple cyst	3	10.0	3) Metastasis	1	3.3
4) Focal pancreatitis	2	6.7			

The most common benign lesion was pancreatic pseudocyst (7 patients= 23.3%), while the most common malignant lesion was pancreatic adenocarcinoma (14 patients = 46.7%).

Histopathological studies were performed for 21 patients either by postsurgical biopsy (14 cases) or by core biopsy (4 cases) and ultrasound guided fine needle aspiration is performed for 3 cases (some cases of pancreatic pseudocyst).

Table 6: MRI findings of pancreatic lesions :

Lesion	No	T1Wis	T2Wis	Post contrast
Pancreatic pseudocyst	7	All cases were hypointense	All cases were hyperintense	-
Pancreatic abscess	2	Hypointense	Hyperintense	-
Simple cyst	3	Hypointense	Hyperintense	-

Focal pancreatitis	2	Isointense	Isointense	Homogenous enhancement pattern
Adenocarcinoma	14	All cases were hypointense	(11) Hyperintense (2) Isointense (1) Hypointense	Poor early enhancement relative to background pancreas
Lymphoma	1	Hypointense	Hyperintense	Enhanced rim
Metastasis	1	Hypointense	Hyperintense	Homogenous enhancement
Total	30	(28) Hypointense (2) Isointense	(25) Hyperintense (1) Hypointense (4) Isointense	

Twenty-five cases of the 30 patients showed increased intensity at T2-weighted images. Most of malignant cases (13 out of 16 cases) showed low or equal intensity on T1- and high intensity on T2-weighted images compared to normal pancreatic parenchyma.

Table 7: Comparison between ADC value of benign and malignant pancreatic masses :

	Benign (14)	Malignant (16)	t-test	P-value
Mean ADC value $\times 10^{-3}$ s/mm² (b=500 s/mm²)	3.006 (2.31-3.56)	1.34 (1.15-1.52)	-4.5	0.001***
Mean ADC value $\times 10^{-3}$ s/mm² (b=1000 s/mm²)	2.49 (2.12-3.01)	1.24 (1.01-1.49)	-2.09	0.002**

There was a statistical significant difference between mean ADC values of benign and malignant pancreatic masses. The mean ADC values of benign and malignant masses done by b-1000 were statistically significant higher than done by b-500. One male patient was diagnosed by DWI as malignant lesion as it has restricted diffusion and low ADC values but histopathological results diagnosed it as benign lesion (focal pancreatitis).

Table 8: Validity of DW-MRI in the diagnosis of pancreatic masses by b value 500 s/mm² :

	Malignant group	Benign group	Total
+Ve:Positive	True +ve 15	False +ve 3	18
-Ve:Negative	False -ve 1	True -ve 11	12
Total	16	14	30

+ve: positive
-ve: negative

DWI images by b value 500 s/mm² detect pancreatic masses with sensitivity of 93.8 %, specificity of 78.6%, positive predictive value (PPV) of 88.88% and negative predictive value of 100 %

Table 9: Validity of DW-MRI in the diagnosis of pancreatic masses by b value 1000 s/mm² :

	Malignant group	Benign group	Total
+Ve	True +ve 16	False +ve 2	18
-Ve	False -ve 0	True -ve 12	12
Total	16	14	30

+ve: positive
-ve: negative

DWI images by b value 1000 s/mm² detect pancreatic masses with sensitivity of 100 %, specificity of 83.33%, positive predictive value (PPV) of 93.33% and negative predictive value of 100 % .

Table 10: Cut off value between benign and malignant pancreatic masses by b value 1000 s/mm² :

	Cut off	sensitivity	Specificity	PPV	NPV	P value
(b=1000 s/mm²)	<1.5	100.0	83.33	88.88	100.0	0.001***

PPV: positive predictive value
 NPV: negative predictive value

The cut-off value for the differentiation of benign from malignant pancreatic masses was $1.5 \times 10^{-3} \text{ s/mm}^2$ with the sensitivity, specificity, PPV, NPV & p value were 100%, 83.33%, 100%, 93.33%, and <0.001 respectively .




Figure 1: Female patient 36 years old presented with abdominal pain giving history of acute pancreatitis two weeks ago.

Laboratory results show: Serum amylase and lipase were high on first 48h then return to normal values, Normal liver & kidney function tests and no elevation in total or direct bilirubin.

MRI revealed:

(A&B): Axial T1&T2 WIs reveal a well defined cystic lesion displaying low T1&high T2 signal with irregular border measuring about 3x2.5cm and showing internal septation at the region of body of pancreas (arrows).

(C&D): Axial DW image & ADC map image with b-value of 500 s/mm² show low and high signal intensity respectively of the lesion denoting free diffusion (arrows). **ADC Value = 2.94×10^{-3} mm²/sec.**

(E&F): Axial DW image & ADC map image with b-value of 1000 s/mm² show low and high signal intensity respectively of the lesion denoting free diffusion (arrows). **ADC Value = 2.41×10^{-3} mm²/sec.**

Diagnosis:

Overall MRI finding are consistent with pancreatic pseudocyst.

UNDER PEER REVIEW




Figure 2: Male patient 55 years old presented with jaundice, weight loss and abdominal pain.

Laboratory results show: Moderate increase in total & direct bilirubin with increased Direct/Total bilirubin (D/T)ratio, moderate increase in urine bilirubin and impaired liver function tests (increased AST, ALT & alkaline phosphatase).

MRI revealed:

(A&B): Axial T1&T2 WIs reveal an ill defined lesion at the region of head of pancreas displaying heterogeneous hypointense T1&isointense T2 signal measuring about 4.0cmx3.5cm and encasing pancreatic vessels (arrows).

(C&D): Axial DW & ADC map image with b-value of 500 s/mm² show high and low signal intensity respectively of the lesion denoting restricted diffusion (arrows). ADC Value = 1.33x10⁻³ mm²/sec

(E&F): Axial DW & ADC map image with b-value of 1000 s/mm² show high and low signal intensity respectively of the lesion denoting restricted diffusion (arrows). ADC Value = 1.21x10⁻³ mm²/sec

Associated findings: Dilated pancreatic duct with abrupt end. There are multiple enlarged para-aortic lymph nodes being hypointense on T1WIs and isointense on T2WIs and show restricted diffusion in the DWIs, largest one measures 10x8mm (black arrows)

Diagnosis:

Radiological: Pancreatic head carcinoma

Pathological: Pancreatic ductal adenocarcinoma

4. DISCUSSION:

The goal of evaluating pancreatic lesions is to differentiate between benign and malignant pancreatic lesions and to determine if malignant masses are resectable or irresectable so that patients with pancreatic cancer can receive adequate treatment at an earlier stage to reduce possible mortality and morbidity of the disease, while avoiding unnecessary surgery in patients with benign nodules.

Radiologic findings play a critical role in the characterization of pancreatic lesions as benign or malignant. The sensitivity and specificity, as well as the detection of extrapancreatic extension, are generally superior with MRI when compared with other imaging techniques .

Our study included 30 patients with pancreatic lesions. Their ages ranged from 30 to 70 years with a mean age 51 years. The most common age group was from 50-60 years. That was going with the same findings as Ichikawa et al. [9]

In the present study, the most common clinical presentation was abdominal pain, less commonly jaundice, anorexia, weight loss, fever or accidentally discovered pancreatic mass during sonographic examinations. That was going with the same findings as Inan et al. [10]

In the current study, the pancreatic lesions were more common in females (17 cases) than males (13 cases). ~~That~~ This is similar with the was going with the same findings by as Kartalis et al. [11]

In our study, the pancreatic masses were more common in the head of the pancreas (17 cases). ~~That was going with the same findings as~~ This is in agreement with the finding by Wang et al. [12]

~~In our study, the pancreatic masses were more common in the head of the pancreas (17 cases). That was going with the same findings as~~ Wang et al. [12]

EXPUNGE :[2U]Comment

In our DWI study, b values were 0, 500& 1000 sec/mm², with the application of parallel imaging and breath triggering technique, a satisfying image quality has been achievable on a 1.5-T scanner within an acceptable acquisition time with an exclusion rate of 0% of patients.

Pancreatic pseudocyst and pancreatic cyst tend to be hypointense relative to normal pancreas on T1-weighted images and hyperintense on T2-weighted images [13]. **These results coincide with our results, as there were 7 cases had pseudocyst and three cases had simple cysts were hypointense relative to pancreatic parenchyma in T1-weighted images and hyperintense on T2-weighted images.**

Signal intensities of cystic lesions were high on diffusion weighted images with lower b values; however, with a higher b-value (b = 1,000 s/mm²) signal intensities of pseudocysts were isointense to the pancreas [11]. **These results coincide with our results, as there were 7 cases had pseudocyst were isointense compared with the pancreas on DWI by b-value 1000 s/mm².**

Signal intensities of simple cysts were hypointense on diffusion-weighted images due to low cellularity which provides a larger extra-cellular space for diffusion of water molecules [11]. **That was going hand in hand with our present study, as there were three cases of simple cysts were hypointense relative to pancreas on DWI by b-value 1000 s/mm² and display T2-shine through effect DWI by b-value 500 s/mm².**

Pancreatic abscesses remained hyperintense. Therefore, the hyperintensity of abscesses, on 1000 s/mm² b factor images cannot be totally attributed to the T2 shine-through effect. Diffusion can be quantitatively evaluated by ADC, which is free of the T2 shine-through effect. The high signal on diffusion weighted images is due to the reduced diffusion in abscesses, because abscesses have a viscous content, they have restricted diffusion with high ADCs value [14]. **That was in agreement with our study, where two cases of pancreatic abscess were hyperintense compared with pancreas.**

Pancreatic adenocarcinoma is the most common malignant neoplasm of the pancreas and continues to be a leading cause of death in the world. Unfortunately, the majority of early and curable pancreatic adenocarcinomas are clinically silent and are associated with mild and nonspecific symptoms [15].

The best way to improve survival in these cases is with early detection and accurate staging [12].

Body MR imaging and diffusion weighted imaging have been increasingly used in the detection and characterization of pancreatic adenocarcinoma [16].

Pancreatic adenocarcinomas are typically hypointense relative to normal pancreatic tissue on T1-weighted images. The normal pancreas appears hyperintense with this sequence and of variable intensity on T2-weighted images [17]. **That was going hand in hand with our present study, as there were 14 cases of pancreatic adenocarcinomas hypointense in T1-weighted images and in 11 out of 14 cases hyperintense in T2-weighted images.**

However, small or non-contour-deforming pancreatic adenocarcinomas may lack classic imaging features and may not be detected on conventional MR images [12].

Therefore, The additional use of diffusion weighted imaging may allow earlier detection of pancreatic adenocarcinoma, since these neoplasms have increased signal intensity on diffusion weighted images with high b-values (b > 500 sec/mm²) and relatively low ADC values because of the restricted diffusion associated with fibrosis [12]. **That was going with the results of our study where 13 out of 14 cases of pancreatic adenocarcinomas were high signal intensity seen at diffusion-weighted images and low signal intensity seen at ADC map images with low ADC values.**

Kartalis et al. [11] presented a study in 36 patients with pancreatic lesions (12 malignant and 24 benign) and 39 patients without lesions were included. MRI-c and DWI (free breathing, b-values 0 and 500 s/mm²) were performed prospectively and consecutively in a 1.5-T system. The analysis was retrospectively performed blinded by two radiologists in consensus. The sensitivity, specificity, accuracy, and positive and negative predictive values of DWI and MRI-c were 92, 97, 96, 85, 98% and 100, 97, 97, 86, 100%, respectively. Mean ADC values of malignant lesions were significantly lower than those of benign lesions.

Ichikawa et al. [9] presented a study in 26 patients with pancreatic adenocarcinoma were included in the study. All patients and subjects underwent DWI (free breathing, b values 1000 s/mm²) reported a sensitivity of 96.2% and a specificity of 98.6%.

Inan et al. [10] presented a study in 42 patients with different cystic lesion of pancreas were included in the study. All patients and subjects underwent DWI (free breathing, b values 500 and 1000 s/mm²) reported a sensitivity of 70% and a specificity of 90%.

In our study, the sensitivity, Specificity, PPV, NPV & p value were 92.8%, 83.33%, 92.8%, 83.33% respectively for b value 500 s/mm².

Also, the sensitivity, Specificity, PPV, NPV & p value were 100%, 83.33%, 93.33%, 100%, and <0.001, respectively for b value 1000 s/mm².

5. CONCLUSION

MRI plays an important role in the diagnosis of different pancreatic lesions and can assess the neoplastic pancreatic lesions with accurate detection of extension, nodal involvement and hepatic metastatic lesions. It also has a major role in differentiation between benign and malignant pancreatic lesions by the aids of DWI.

Ethical consideration:

1. Written informed consent was obtained from all patients' guardians after full explanation of the benefits and risks of procedure.
2. Any unexpected risks appeared during the course of the research were cleared to all parents.

PART OF NOT :[3U]Comment
DISCUSSION

EXPUNGE

3. The privacy of the participants and confidentiality of data were guaranteed during the various phases of the study. The results will be used as scientific material only and will not be used by any legal authorities.

REFERENCES

1. Siegelman ES and Adusumilli S: MRI of the bile ducts, gall bladder and pancreas. In: Siegelman ES. Body MRI. Philadelphia: Elsevier Saunders 2005; PP. 63-128.
2. Vaughn D, Jabra A and Fishman E: Pancreatic disease in children and young adults: Evaluation with CT. *RadioGraphics* 1998; 24:1171-1187.
3. Masud R, Waheed A, Aurangzeb, et al: Role of radiology in pancreatic disorders Pakistan Armed Forces Journal 2009; 2:1-2.
4. Olivia MR, Mortele KJ, Erturk SM, et al: Magnetic resonance imaging of the pancreas. *Applied Radiology* 2006; 23: 671-685.
5. Ly JN and Miller FH: MR imaging of the pancreas, a practical approach, *Radiologic Clinics of North America*. 2002; 40(6):1289-1306.
6. Sandrasegaran K, Lin C, Akisik FM, et al: State-of-the-Art Pancreatic MRI. *AJR* 2010; 195:42-53.
7. Vitellas KM, Keogan M, Spritzer C, et al: MR Cholangiopancreatography of Bile and Pancreatic Duct Abnormalities with Emphasis on the Single-Shot Fast Spin-Echo Technique. *Scientific Exhibit* 2000; 20:939-957.
8. Niwa T, Ueno M, Ohkawa S, et al: Advanced pancreatic cancer: the use of the apparent diffusion coefficient to predict response to chemotherapy. *The British Journal of Radiology* 2009; 82:28-34.
9. Ichikawa T, Erturk SM, Motosugi U, et al: High-b Value Diffusion-Weighted MRI for Detecting Pancreatic Adenocarcinoma. *AJR* 2007; 188:409-414.
10. Inan N, Arslan A, Akansel G, et al: Diffusion-weighted imaging in the differential diagnosis of cystic lesions of the pancreas. *AJR* 2008; 191: 1115-1121.
11. Kartalis N, Lindholm T, Aspelin P, et al: Diffusion-weighted magnetic resonance imaging of pancreas tumours. *European Journal of Radiology* 2009; 19(8): 1981-1990.
12. Wang Y, Miller FH, Chen ZE, et al: Diffusion-weight MRI of solid and cystic lesions of the pancreas. *Radio-Graphics* 2011; 31 (3): 47-64.
13. Hruban RH, Iacobuzio-Donahue CA, Poultsides GA, et al: Histopathologic basis for the favorable survival after resection of intraductal papillary mucinous neoplasm associated invasive adenocarcinoma of the pancreas. *Ann Surg* 2010; 251 (3):470-6.
14. Chan JH, Tsui EY, Luk SH, et al: Diffusion weighted MRI of the liver: distinguishing hepatic abscess from cystic or necrotic tumor. *Abdom Imaging* 2007; 26: 161-165.
15. Megibow AJ: Are we really closer to predicting the development of pancreatic cancer? *Radiology* 2010; 254 (3): 642-646.
16. Park HS, Lee JM, Choi HK, et al: Preoperative evaluation of pancreatic cancer: comparison of gadolinium-enhanced dynamic MRI with MR cholangiopancreatography versus MDCT. *J Magn Reson Imaging* 2009; 30 (3): 586- 595.
17. Miller F, Rini N and Keppke A: MRI of adenocarcinoma of the pancreas. *American Journal of Roentgenology* 2006; 187: W365-372.

ABBREVIATIONS

ADC: Apparent diffusion coefficient
DWI: Diffusion-weighted MRI
DW-MRI: Diffusion Weighted Magnetic Resonance Imaging
ERCP: Endoscopic retrograde cholangiopancreatography
MRCP: Magnetic resonance cholangiopancreatography
MRI: Magnetic resonance imaging
NPV: Negative predictive value

PPV: Positive predictive value

UNDER PEER REVIEW

